

Morphometric Discriminant Analysis of isolated chondrichthyan scales for palaeoecological inferences: the Middle Triassic of the Iberian Chain (Spain) as a case of study

H. Ferrón¹, C. Pla¹, C. Martínez-Pérez^{1,2}, M.J. Escudero-Mozo^{3,4}, H. Botella^{1*}

¹Departamento de Geología, Universidad de Valencia, Avda. Dr. Moliner, 50, 46100 Burjassot, Spain.

²School of Earth Sciences, University of Bristol, Wills Memorial Building, Queen's Road, Bristol BS8 1RJ, United Kingdom.

³Departamento de Estratigrafía, Universidad Complutense de Madrid, C/José Antonio Novais 12, 28040 Madrid, Spain

⁴Instituto de Geociencias IGEO (CSIC,UCM), C/José Antonio Novais 12, 28040 Madrid, Spain

e-mail addresses: humfeji@alumni.uv.es (H.F.), M.Cristina.Pla@uv.es (C.P.), Carlos.Martinez-Perez@bristol.ac.uk (C.M.-P.); mjescudero@geo.ucm.es (M.J.E.-M.)
Hector.Botella@uv.es (H.B., *corresponding author)

Received: 12 December 2012 / Accepted: 4 December 2013 / Available online: 25 February 2014

Abstract

Palaeontological studies on exoskeletal disarticulated remains of chondrichthyans have focused on teeth and only less interest has been paid to scales due their limited taxonomic and systematic significance. However, classical works linking the morphology and the function of the squamation in extant sharks suggest that, despite their limited taxonomic value, the study of isolated scales can be a useful tool for palaeoenvironmental and palaeoecological inferences. Following this idea, we have analyzed the fossil record of shark scales from two Middle Triassic sections of the Iberian Chain (Spain), identifying different functional types by means of a morphometric discriminant analysis. From a total of 1136 isolated chondrichthyan scales, 25% were identified as abrasion resistant scales, 62% as drag reduction scales and 13% as scales of generalized functions. The elevated proportion of abrasion resistant scales suggests that this chondrichthyan palaeocommunity was highly dominated by benthic sharks that lived over a hard sea floor. However, one of the stratigraphical levels studied (He-20), presents statistically significant differences from the others, showing a lower percentage of abrasion resistant scales and a larger percentage of drag reduction scales. This level can be linked with storm episodes that could introduce remains of benthic-pelagic or pelagic forms in the inner platform. Finally, partial correlation analysis between relative abundances of functional scale types and tooth-based taxa from the same sections provide positive correlation between teeth of *Hybodus* and *Pseudodolaticus* and drag reduction scales, and teeth of *Prolatodon* and abrasion strength scales.

Keywords: Triassic, Chondrichthyes, scales, teeth, Iberian Chain

Resumen

Los estudios paleontológicos de restos desarticulados de condriactios se han centrado en los dientes, no prestando prácticamente interés al estudio de sus escamas debido a su limitada importancia taxonómica y sistemática. Sin embargo, algunos trabajos clásicos que han relacionado su morfología y función en base al estudio de la escamación de tiburones actuales, sugieren que, a pesar de su limitado valor taxonómico, el estudio de las escamas aisladas puede ser una herramienta útil para obtener inferencias paleoambientales y paleoecológicas. Siguiendo estas ideas, se ha analizado el registro fósil de escamas de tiburón de dos secciones del Triásico Medio de la Cordillera Ibérica (España), identificando diferentes tipos funcionales por medio de un análisis morfométrico discriminante. De un total de 1.136 escamas aisladas de condriactios, el 25% de ellas fueron identificadas como escamas resistentes a la abrasión, el 62% como de reducción de la fricción con el agua y el 13% de las escamas como de función generalizada. La elevada proporción de las escamas resistentes a la abrasión sugiere que esta paleocomunidad de condriactios estaba claramente dominada por tiburones bentónicos que habitaban sobre un sustrato rocoso. Sin embargo, uno de los niveles estratigráficos estudiados (He-20), presenta diferencias estadísticamente significativas con los demás, mostrando un porcentaje más bajo de escamas resistentes a la abrasión y un porcentaje mayor de las escamas de reducción de la fricción con el agua. Este nivel se relaciona con episodios de tormenta que pudieron haber introducido los restos de formas bento-pelágicas o pelágicas en la plataforma interna. Por último, el análisis de correlación parcial entre la abundancia relativa de los tipos funcionales de escamas y taxones en basados dientes de las mismas secciones proporcionan una correlación positiva entre los dientes de *Hybodus* y *Pseudodolaticus* y escamas de reducción de la fricción con el agua, y dientes de *Prolatodon* y escamas resistencia a la abrasión.

Palabras clave: Triásico, condriactios, escamas, dientes, Cordillera Ibérica

1. Introduction

Due to the cartilaginous nature of the chondrichthyan endoskeleton, the fossil record of this group consists mainly of disarticulated remains such as teeth, scales and fin spines. Within these elements teeth are by far the most informative in terms of taxonomy, systematics or autecology. Most of the extinct species of Chondrichthyes are described on the basis of isolated teeth (see for example monographs of Capetta, 1987, 2012; Ginter *et al.*, 2010 and references therein). In contrast, isolated chondrichthyan scales that commonly occur together with disarticulated teeth, provide limited taxonomic information (especially Mesozoic and Cenozoic taxa, see Reif, 1985a; Karatajute-Talimaa, 1998; Leidner and Thies, 1999; Thies and Leidner, 2011) for three main reasons: the presence of a high morphological diversity of scales in different regions of the body, which has been documented in both extant and fossil specimens (see Reif, 1973, 1974, 1985a); the evolution of some scales of similar, or even, identical morphologies in distantly related species (Muñoz-Chápuli, 1985); and the occurrence of ontogenetic variability (Reif, 1973, 1978). As a consequence, palaeontological studies of disarticulated remains of chondrichthyans have focused on teeth and very little attention has been paid to scales. However we propose that, despite the lack of taxonomic and systematic significance, isolated scales could provide useful information regarding the palaeoecological and palaeoenvironmental conditions. Our proposal is based on the classic works of Reif (1982; 1985a) where the relationship between the shape and function of scales of extant sharks was pointed out. Reif (1982, 1985a) differentiated scales into five different functions: abrasion resistance, defense, drag reduction, bioluminescence and generalized functions, and identified eight characteristic morphologies associated with these functional types. Abrasion resistant scales are found in demersal sharks, which inhabit rocky or coralline substrates and, generally, in small body regions that are often subject to abrasion in all other sharks, such as the mouth area. Scales of this type are knob-like and smooth (Morphology 1) or strongly ornamented (Morphology 2). Both types very frequently show scratch marks. Defensive scales are common in demersal sharks inhabiting muddy or sandy substrates, and protect them against ectoparasites and the settlement of epibionts. Scales of this functional type are thorn-shaped with the cusps pointing in an upward-posterior direction commonly accompanied by mucus (Morphology 3). Drag reduction scales cover most of the skin surface in fast swimming pelagic sharks. Scales of this functional type have riblets aligned in the direction of fluid flow (Morphology 4). The geometry and arrangement seem to play an important role in the drag reduction although the underlying mechanisms are not well understood yet (e.g. Bechert *et al.*, 2000; Douglas-Dean, 2011; Raschi and Musick, 1986; Reif and Dinkelacker, 1982; Reif, 1985a). Scales associated to bioluminescence have evolved in some mesopelagic sharks,

enabling the skin to carry photophores and permitting light to pass between them (Reif, 1985b). This functional type is represented by three different morphologies: square-shaped with concave facets (Morphology 5), bristle-shaped (Morphology 6) and thorn-shaped (Morphology 7). Scales with ridges and lateral cusps well developed but shorter than the principal cusp fulfill generalized functions and are found in almost all sharks (Morphology 8). Other functions related to hatching and feeding has been documented for shark scales (Grover, 1974; Reif, 1974; Southall and Sims, 2003), but they are very atypical and have not been found among our material.

In this study we firstly characterize the morphology of the five functional types proposed by Reif (1985a) using traditional morphometrics and discriminant analysis. Secondly we assign isolated placoid scales from synchronous levels of two stratigraphical sections (Middle Triassic) of the Iberian Chain (Spain), to these “functional” morphologies. The differential abundance of the functional types allow us to evaluate the structure of the chondrichthyan paleocommunity in terms of the relative dominance of more pelagic or benthic sharks and the properties of the physical environment they inhabited, such as the dominant substrate type.

2. Material provenance

The Middle Triassic carbonates in facies Muschelkalk (Anisian-Ladinian in age) of the eastern Iberian Chains, is represented by two transgressive-regressive cycles showed by two carbonate units (ramps) mainly constituted by epicontinental shallow marine environments. The Cañete Fm. (López-Gómez and Arche, 1992) corresponds to the upper carbonate unit and represents the second and main marine transgression of the Middle Triassic in the southeast area of the Iberian Chains. Litologically this unit can be subdivided in two parts: a lower one, strongly dolomitized and constituted by grey massive dolomites; and an upper part, less dolomitized and mainly composed by thin-bedded bioclastic limestones, planar stromatolitic lamination, tempestite and marls levels constituting shallowing-upwards sequences (shallow inner ramp-tidal flat). The studied samples correspond to sediments of the upper part of the Cañete Fm, which has been dated as Ladinian based on ammonites, foraminifera and conodonts (López-Gómez *et al.*, 1998). A total of 1136 isolated scales were obtained after the dissolution of Triassic carbonate rocks with 10% acetic acid from the Bugarra and Henarejos sections of the Iberian Chain (Spain). Both sections expose the dolomitic and limestone sediments of the Cañete Formation.

The Henarejos section is located 1 km south-east of the village of Henarejos (Province of Cuenca). Ladinian molluscs have been reported by Marquez-Aliaga (1985) and López *et al.* (1987). The Bugarra section is close to the village Bugarra (Province of Valencia). Márquez-Aliaga *et al.* (1984) studied the stratigraphy and the invertebrate paleontological aspects of this section (Fig. 1). All specimens studied herein come




Fig. 1.- A. Geographical setting of the studied area with indication of the studied sections. B. Stratigraphical column of Bugarra and Henarejos sections with a possible litho- and biostratigraphical correlation between them and with indication of levels that yielded the material described in this work (modified from Escudero *et al.*, 2012).

from the uppermost member of the dolomites and limestones of the Cañete Formation (see Fig. 1) and are kept in the Museum of Geology of the University of Valencia (MGUV).

3. Methodology

In order to characterize the morphologies of each functional type proposed by Reif (1985a) we performed classical morphometric analysis of scales with known functions in extant sharks. Six variables (for explanation see Table 1) were

measured on the dorsal surface of the scale crown from specimens figured in Reif (1985a) using ImageJ software. We used a total of 58 scales belonging to the eight scale morphologies. Each morphology was treated as a group. Discriminant analysis was performed to obtain maximum separation among the eight scale morphologies using SPSS Predictive Analytics Software Statistics (PASW) version 18.0. Three variables (LEN, WID and RID) were log-transformed to allow non-linear combinations between them as sums or subtractions of logarithms in the discriminant functions (see below). Sub-




Fig. 2.- Scatter plot based on the first two discriminant functions, showing the eight scale morphologies and the five functional types.

sequently the fossil specimens were included in the discriminant analysis as unknown specimens and were assigned to one of the morphologies of functional types described by Reif (1985a) based on the similarity of their centroid values. Differential abundances of these functional types were statistically analyzed by Pearson's Chi-square test and Z-test using PASW software with the purpose of detecting differences between stratigraphic levels of the studied sections. Finally, we used Partial Correlation Analysis to compare the relative abundances of scale functional types vs. the relative abundances of tooth-based genera in the different stratigraphic levels (dates for tooth occurrences were taken from previous studies, see below). Partial Correlation Analysis allows the study of the relationship between two quantitative variables controlling the possible effect of another one that could mask correlations. Thus, the analysis was repeated four times controlling, in each case, the abundance of a concrete tooth-based genus.

CIR.	Circularity = 4Π (area/perimeter ²)
ANG.	Angle created between the two most lateral ridges / 2
PA	Presence or absence of ridges
LEN.	Maximum length of the scale
WID.	Maximum width of the scale
RID.	Average length of ridges

Table 1. Explanation and coded designations of measured scale characters.

Variable	Function					
	1	2	3	4	5	6
CIR.	-0.967	-0.304	0.179	-0.458	-0.168	0.187
ANG.	1.303	1.960	5.761	1.489	0.748	0.946
PA	7.460	-8.598	7.560	5.703	0.657	-7.984
Log (LEN.)	0.609	-0.389	0.656	-1.191	-1.061	-0.349
Log (WID.)	-0.391	0.175	-0.691	1.694	0.108	-0.229
Log (RID.)	-6.266	11.025	-1.864	-4.311	0.362	8.077

Table 2. Standarized canonical discriminant function coefficients.

4. Results and Discussion

4.1. Discriminant analysis using scales of extant sharks.

Canonical variate analysis generated six discriminant functions. Coefficients, eigenvalues, proportion of explained variance and canonical correlation are presented in Tables 2 and 3. The first canonical discriminant function explains 53.6 % of the total variance while the second one accounts for 25.8% of the total variance (79.5% cumulative variance). The plot of the two canonical variables illustrates a good separation between the eight morphologies proposed by Reif (1985a) (Fig. 2). The discriminant analysis correctly classifies 100% of original cases, while the percentage of cross-validated grouped cases correctly classifies 93.1 % (i.e. 54 of the 58 scales) (Table 4). In this analysis, scales of morphologies 1, 2, 4 and 7 are the most correctly classified followed by scales of morphology 8. The least correctly classified are the scales of morphologies 3 and 6 (Table 4).

These results show that classical morphometric analysis is able to discriminate between the five functional types of chondrichthyan scales proposed by Reif (1985a) using six variables in the discriminant functions. In addition, we also tested other alternatives to the use of the classical morphometric analysis, like geometric morphometrics or Fourier analysis, but the results were not as good as expected. Due to the high morphological variability in scales, it was very difficult to set homologous points in the eight scale morphologies and, therefore, it was not possible to apply geometric morphometrics in an acceptable way. On the other hand, Fourier analysis only takes into account characters reflected in the contour of the scales without considering other features, as for example those related to the ridges, which might be useful to differentiate between groups. Although the results of this method are good, with the percentage of cross-validated grouped cases correctly classified ranging from 42.2-80% depending on the number of harmonics and the type of analysis

Function	Eigenvalue	% of Variance	Cumulative % of Variance	Canonical Correlation
1	23.079	53.6	53.6	0.979
2	11.115	25.8	79.5	0.958
3	5.124	11.9	91.4	0.915
4	2.910	6.8	98.1	0.863
5	0.639	1.5	99.6	0.624
6	0.168	0.4	100.0	0.379

Table 3. Eigenvalues, proportion of explained variance and canonical correlation of the discriminant functions.

(polar or elliptic), they are not as accurate as those obtained by classical morphometric analysis (Appendix 1 shows results obtained using Fourier analysis).

4.2. Discriminant analysis including fossil specimens.

Once the discriminant analysis was established, the 1136 fossil scales from the Henarejos and Bugarra sections were included as unknowns. Following the predictions obtained, fossil scales were assigned to four morphologies belonging to three of Reif’s functional types (Fig. 3). 289 fossil specimens (25 %) were identified as abrasion resistant, 705 (62 %) as drag reducing and 142 (13 %) were identified as scales of generalized functions. None of the fossil scales were assigned to either bioluminescence type (morphologies 5, 6 and 7) or to the defense type (morphology 3). Abundances of each functional type by section and stratigraphic level are shown in Table 5.

When it is taken into consideration that in pelagic sharks scales for abrasion resistance are restricted only to small areas of the body (e.g. surrounding the mouth) representing less than 5% of the total body surface, the high proportion of scales with a protective function against abrasion present

	Predicted group								TOTAL
	Morphology 1	Morphology 2	Morphology 3	Morphology 4	Morphology 5	Morphology 6	Morphology 7	Morphology 8	
Morphology 1	5 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	5 (100.0)
Morphology 2	0 (0.0)	5 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	5 (100.0)
Morphology 3	0 (0.0)	1 (20.0)	4 (80.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	5 (100.0)
Morphology 4	0 (0.0)	0 (0.0)	0 (0.0)	15 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	15 (100.0)
Morphology 5	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	7 (87.5)	1 (12.5)	0 (0.0)	0 (0.0)	8 (100.0)
Morphology 6	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	4 (80.0)	1 (20.0)	0 (0.0)	5 (100.0)
Morphology 7	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	5 (100.0)	0 (0.0)	5 (100.0)
Morphology 8	0 (0.0)	0 (0.0)	0 (0.0)	1 (10.0)	0 (0.0)	0 (0.0)	0 (0.0)	9 (90.0)	10 (100.0)

Table 4. Count (Non-italic numbers) and percentages (Italic numbers) of cross-validated grouped cases correctly classified.




Fig. 3.- Morphologies of fossil scales from Henarejos and Bugarra sections. A: MGUV-24.804. Morphology 1 (Abrasion resistant scales) in upper-frontal view. B: MGUV-24.799. Morphology 2 (Abrasion resistant scales) in frontal view. C: MGUV-24.798. Morphology 4 (Drag reduction scales) in frontal view. D: MGUV- 24.803. Morphology 8 (Scales with generalized functions) in frontal view. Scales: A, C, 500 μm .; B, 100 μm .; D, 200 μm .

in our association become interesting for the interpretation of the data. It could be reasonably related with the presence of a high number of benthic sharks, in which (especially in those living on hard sea floor) abrasion resistant scales cover not only the mouth area but also half of the ventral part of the body and the ventral and anterior areas of the pectoral and pelvic fins (Reif, 1985a). Thus, our results indicate that the Middle Triassic chondrichthyan community from the Iberian Chain was dominated by benthic sharks. This suggests a palaeoenvironmental interpretation as an area of shallow marine waters, probably dominated by an abrasive substrate (such as a near-shore hard sea floor environment or some type of lagoon). Noticeably, this interpretation is in accordance with sedimentological data from previous studies in the area (e.g. Marquez-Aliaga *et al.* 1984; López-Gómez *et al.*, 1993), providing an independent line of evidences that validate our analysis.

4.3. Pearson's Chi-square test and Z-test.

Pearson's Chi-square test showed significant differences between levels of the Bugarra and Henarejos sections ($p = 0.000$ Table 5). Z-test allowed us to detect between what levels and in which concrete functional types significant differences occur (Table 6). The test show significant differences between level He-20 and the remaining levels. The percentage of abrasion resistant scales is significantly lower than in

all others levels and the percentage of drag reduction scales is significantly higher than in four of the other seven levels (Bu 1-33, Bu 1-26, Bu1-26d and He-18). This suggests a change in the composition of the chondrichthyan community with an increase of the pelagic or benthic-pelagic forms. In fact the fraction of abrasion resistant scales (5.2%) is close to the expectation for pelagic sharks. Furthermore teeth of supposed benthic taxa (see below) do not occur in this level. These changes could be explained by (1) punctuated changes in the bathymetry of the basin related to transgressive-regressive pulses that produce a deepening and a displacement of the coastline] or (2) by storm episodes that introduce these communities from deeper waters into the inner platform, where sediments of level He-20 were deposited. Sedimentological dates more likely support the second scheme. According with this dates all the levels of Henarejos section correspond with shallow marine environments within the inner platform, without evidence of any punctual deepening. Moreover, in the top levels of the stratigraphical sequence different tempestite levels are recorded, just above the He-20 level.

4.4. Correlation with fossil teeth.

Previous studies (Pla *et al.*, 2009; 2013) on isolated teeth from the same samples that the scales of this study identified the following tooth based species: *Palaeobates angustissimus* (Agassiz, 1838), *Pseudodalatias henarejensis* Botella, Plasencia, Márquez-Aliaga, Cuny and Dorka, 2010; *Hybodus bugarensis* Pla, Márquez-Aliaga and Botella, in press; *Prolatodon bucheri* (Cuny, Rieppel and Sander, 2001); *Prolatodon contrarius* (Johns, Barnes and Orchard, 1997); *Hybodus plicatilis* Agassiz, 1838; and *Lissodus aff. lepagei* (Fig. 4). Partial correlation analysis between the relative abundances of functional types of scales and abundances of tooth-based taxa at generic level indicates positive correlation between teeth of *Hybodus* and *Pseudodalatias* and drag reduction scales, and teeth of *Prolatodon* and abrasion resistant scales. A negative correlation was present between teeth of *Prolatodon* and drag reduction scales (Table 7). Once more, these results are in agreement with the palaeobiological interpretations based on the tooth morphology (see Pla *et al.*, 2013). Thus, dentitions of *Prolatodon bucheri* and *Prolatodon contrarius* can be identified as belonging to a grasping-crushing feeding strategy (following the terminology of Cappetta, 1986, 1987) with cuspidate anterior teeth and flat lateral teeth. This dental type (present generally in sharks of benthic habits) corresponds to a trophic adaptation for durophagy and indicates crustaceans, ostracods, or shelled invertebrates such as gastropods and bivalves as the preferred prey. In consequence, the obtained positive correlation between *Prolatodon* and abrasion strength scales is expected. In addition, *Pseudodalatias* exhibited a cutting-clutching dentition extremely similar to those of some extant neoselachian Dalatiidae (Gray, 1851). The feeding preferences of dalatiids consist of "parasitic" bites excising portions of flesh from large-size oceanic ani-

A		Level							
		He - 14		He - 18		He - 19		He - 20	
Functional type		Count	%	Count	%	Count	%	Count	%
	Abrasion strength	32	22.9	90	46.2	31	16.4	10	5.2
	Drag reduction	96	68.6	96	49.2	137	72.5	152	79.6
	Generalized functions	12	8.6	9	4.6	21	11.1	29	15.2
TOTAL		140	100	195	100	189	100	191	100

B		Level							
		Bu 1 - 33		Bu - pl		Bu 1 -26		Bu 1 - 26d	
Functional type		Count	%	Count	%	Count	%	Count	%
	Abrasion strength	13	39.4	18	21.7	10	41.7	85	30.2
	Drag reduction	16	48.5	60	72.3	10	41.7	138	49.1
	Generalized functions	4	12.1	5	6.0	4	16.7	58	20.6
TOTAL		33	100	83	100	24	100	281	100

Test of independence of functional types abundances and stratigraphic levels: $\chi^2=139.301$; f.d. = 14; Sig. = 0.000.

Table 5. Count of scales (Non-italic numbers) and differential abundance (Italic numbers) of each functional type by levels and sections (A: Henarejos; B: Bugarra).

		Level			
		He - 14 (A)	He - 18 (B)	He - 19 (C)	He - 20 (D)
Functional type	Abrasion strength	D	A. C. D	D	
	Drag reduction	B		B	B
	Generalized functions				B

		Level			
		Bu 1 - 33 (A)	Bu - pl (B)	Bu 1 -26 (C)	Bu 1 - 26d (D)
Functional type	Abrasion strength				
	Drag reduction		C. D		
	Generalized functions				B

		Level							
		Bu 1 -33 (A)	Bu - pl (B)	Bu 1 -26 (C)	Bu 1-26d (D)	He - 14 (E)	He - 18 (F)	He - 19 (G)	He - 20 (H)
Functional type	Abrasion strength	H	H	H	G.H	H	B.D.E.G.H	H	
	Drag reduction		D.F			D.F		D.F	A.C.D.F
	Generalized functions				E.F				F

Table 6. Z-test results based on two-sided tests with a significance level 0.05. For each significant pair, the key of the category with the smaller column proportion appears under the category with the larger column proportion.

Table 7. Partial correlation analysis results between fossil teeth and functional types of scales. Significant correlations are identified with a single asterisk (A: Control variable *Prolatodon*; B: Control variable *Lissodus*; C: Control variable *Pseudodaltias*; D: Control variable *Hybodus*).

A			Teeth genera		
			<i>Hybodus</i>	<i>Pseudodaltias</i>	<i>Lissodus</i>
Functional type	Abrasion strength	Correlation	-0.142	0.214	-0.171
		Significance	0.761	0.645	0.714
	Drag reduction	Correlation	-0.080	0.182	-0.225
		Significance	0.864	0.697	0.628
	Generalized functions	Correlation	0.358	-0.622	0.607
		Significance	0.431	0.136	0.148
B			Teeth genera		
			<i>Hybodus</i>	<i>Pseudodaltias</i>	<i>Polyacrodus</i>
Functional type	Abrasion strength	Correlation	-0.515	-0.269	0.775*
		Significance	0.237	0.560	0.041*
	Drag reduction	Correlation	0.431	0.4510.310	-0.864*0.012*
		Significance	0.335		
	Generalized functions	Correlation	0.303	-0.557	0.242
		Significance	0.509	0.175	0.601
C			Teeth genera		
			<i>Hybodus</i>	<i>Lissodus</i>	<i>Polyacrodus</i>
Functional type	Abrasion strength	Correlation	-0.668	-0.041	0.760*
		Significance	0.101	0.931	0.047*
	Drag reduction	Correlation	0.793*	-0.125	-0.825*
		Significance	0.033*	0.789	0.022*
	Generalized functions	Correlation	-0.198	0.519	-0.014
		Significance	0.670	0.233	0.977
D			Teeth genera		
			<i>Pseudodaltias</i>	<i>Lissodus</i>	<i>Polyacrodus</i>
Functional type	Abrasion strength	Correlation	-0.549	-0.043	0.673
		Significance	0.202	0.928	0.098
	Drag reduction	Correlation	0.809*	-0.249	-0.833*
		Significance	0.027*	0.590	0.020*
	Generalized functions	Correlation	-0.690	0.676	0.473
		Significance	0.086	0.095	0.284

mals, including other sharks, marine mammals and bony fishes (Gasparini and Sazima, 1996; Soto and Mincarone, 2001; Heithaus, 2004; Heithaus and Vaudo, 2012). This trophic behavior is favoured by sharks with a benthic-pelagic swimming mode that look for prey in the water column. Therefore, the positive correlation between teeth of *Pseudodaltias* and drag reduction scales was also predictable. Finally, the multicuspitate teeth of hybodonts are considered to be adapted for a “grasping and swallowing” strategy (clutching- or tearing-type *sensu* Cappetta, 1987) that could include prey such as other small fishes, soft-bodied animals and arthropods. This strategy is present in both pelagic (as in some *Isurus*) and benthic sharks (such as Scyliorhinidae or Squatinidae, Cappetta, 1987). The positive correlation found between *Hybodus* teeth, which almost all belong to the species *H. plicatilis*, and scales of the drag reduction type indicate that this widespread shark – known from Germany (Agassiz, 1843), Switzerland (Meyer, 1849; Rieppel, 1981; Scheinpflug, 1984) Spain (Pla

et al., 2009; 2013) and Saudi Arabia (Vickers *et al.*, 1999) was a benthic-pelagic to pelagic swimmer.

It is important to remark, that the good statistical correlation found between the abrasion resistant scales and durophag tooth-based taxa (that putatively belonging to benthic sharks), and the high correlation between cutting teeth and drag reduction scales (that could belong to more pelagic shark), strongly supports the idea that no taphonomic bias affect significantly the relative abundance of the different type scales, and hence our final results.

5. Conclusions

Classical morphometric analysis has been able to discriminate among the eight morphologies, included in five functional types, of chondrichthyan scales proposed by Reif (1985a). A high percentage of cross-validated grouped cases were correctly classified (93.1 %). This strongly supports a




Fig. 4.- Fossil teeth from Henarejos and Bugarra sections (from Pla *et al.*, 2013). A: MGVU 25791. *Palaeobates angustissimus*, occlusal view, Bugarra section. B: MGVU 25854. *Lissodus* aff. *L. lepagei*, lingual view, Bugarra section. C: MGVU 25822. *Prolatodon contrarius*, labial view, Henarejos section. D: MGVU 25796. *Prolatodon bucheri*, labial view, Bugarra section. E: MGVU 25868. *Pseudodalatias henarejensis*, lower tooth in labial view, Henarejos section. F: MGVU 25869. *Pseudodalatias henarejensis*, upper tooth in lingual view, Henarejos section. G: MGVU 25831. *Hybodus bugarensis*, labial view, Henarejos section. H: MGVU 25837. *Hybodus plicatilis*, labial view, Bugarra section. Scales: 200 µm.

high possibility of correct identification when the discriminant functions are applied to isolated fossil scales.

A total number of 1136 isolated chondrichthyan scales were collected from two Middle Triassic sections (Henarejos and Bugarra) of the Iberian Chain. They were included in the discriminant analysis resulting in the identification of 25% as abrasion resistant, 62% as drag reduction and 13% as scales of generalized functions. Neither the scales of the bioluminescent nor the defensive type defined by Reif (1985a) could be recognized. The elevated proportion of abrasion resistant scales indicates that the chondrichthyan palaeocommunity from the Middle Triassic of the Iberian Chain was dominated by sharks adapted to a benthic lifestyle. This is in agreement with palaeobiological interpretations provided by previous analyses of isolated chondrichthyan teeth from the same sections (Pla *et al.*, 2013) which

show that in terms of diversity, the chondrichthyan fauna was dominated by durophagous sharks most of them with grasping-crushing dentitions. Dentitions of this type imply feeding preferences including crustaceans, ostracods and shelled invertebrates (gastropods and bivalves), and they are currently present in sharks which live in close relationship with the substrate bottom looking for their prey. Thus, in our opinion, the high percentage of abrasion resistant scales (i.e. of benthic sharks) could be likely related with a palaeoenvironment of shallow marine waters, mostly dominated by a rough substrate, such as a rocky shore platform. This interpretation based on isolated scales is in agreement with sedimentological and other paleontological studies in the area (see e.g. Márquez-Aliaga *et al.*, 1984; Márquez-Aliaga and López-Gómez, 1989; López-Gómez and Arche, 1992; López-Gómez *et al.*, 1993 among others).

The statistical analyses results in differences between level He-20 (Henarejos section) and all other levels concerning the relative abundance of the functional types of scales. The percentage of the abrasion resistant type in this single level is significantly lower but that of drag reduction scales significantly higher. This can be linked with punctuated storm episodes that transported remains of allochthonous chondrichthyan community with an increase in pelagic forms. Finally, the partial correlation analysis between the relative abundances of functional types of scales and tooth-based taxa allowed the detection of a positive correlation between teeth of *Hybodus* and *Pseudodalatias* and drag reduction scales, and teeth of *Prolatodon* and scales of the abrasion resistant. In summary, our analyses show that, despite their limited taxonomic value, isolated scales can be a useful tool for palaeoenvironmental and palaeoecological inferences of the chondrichthyan palaeocommunity. The good concordance of the results obtained from morphometric discriminant analysis of chondrichthyan scales from the Middle Triassic of the Iberian Chains with inferences based on isolated teeth and sedimentological data suggest that this methodology can be extrapolated for studies in other localities.

Acknowledgments

This work has been supported by the Program 458.10/2007 and /2008 from Conselleria de Cultura, Educació i Esports (Generalitat Valenciana) and by Grant BFU2008-00643 to D.R.G. from the Spanish Ministerio de Ciencia e Innovación. We thank Dr. Soledad de Esteban (Institut Català de Paleontologia) and an anonymous reviewer their helpful comments that have improved considerably the original manuscript; and Dr. John Cunningham (University of Bristol) for review and improve the *English*. CMP benefits from a postdoctoral contract Marie Curie FP7-People IEF 2011-29968.

References

- Agassiz, L. (1833–44): *Recherches sur les poissons fossiles*, 3. Imprimerie de Petitpierre, Neuchâtel 32: 390 p.
- Botella, H., Plasencia, P., Márquez-Aliaga, A., Cuny, G., Dorka, M. (2009): *Pseudodalatias henarejensis* nov. sp. A new Pseudodalatiid (Elasmobranchii) from the Middle Triassic of Spain. *Journal of Vertebrate Paleontology* 29(4), 1006-1012. doi: 10.1671/039.029.0425
- Beichert, D. W., Bruse, M., Hage, W. (2000): Experiments with three-dimensional riblets as an idealized model of shark skin. *Experiments in Fluids* 28(5), 403-412. doi: 10.1007/s003480050400
- Cappetta, H. (1986): Types dentaires adaptatifs chez les sélaciens actuels et post-paléozoïques. *Palaeovertebrata* 16, 57-76.
- Cappetta, H. (1987): *Chondrichthyes II, Mesozoic and Cenozoic Elasmobranchii. Handbook of Paleoichthyology*. Verlag Dr. Friedrich Pfeil, München: 193 p.
- Cappetta, H. (2012): *Chondrichthyes, Mesozoic and Cenozoic Elasmobranchii: Teeth. Handbook of Paleoichthyology*. Verlag Dr. Friedrich Pfeil, München: 521 p.
- Cuny, G., Rieppel, O., Sander, P.M. (2001): The shark fauna from the Middle Triassic (Anisian) of North-Western Nevada. *Zoological Journal of the Linnean Society* 13, 285-30. doi: 10.1111/j.1096-3642.2001.tb00627.x
- Douglas-Dean, B. (2011): *The effect of shark skin inspired riblet geometries on drag in rectangular duct flow*. Master thesis. Ohio: 86 p.
- Escudero M.J., Pla, C., Botella, H., Márquez-Aliaga, A. (2012): Situación estratigráfica de tiburones del Triásico Medio de la Cordillera Ibérica. En: XXVIII Jornadas de la Sociedad Española de Paleontología y Simposios de los Proyectos nº 587 y 569 del PICG. Valencia y Sóller, 1-6 de octubre de 2012. Homenaje a Guillem Colom Casanovas (1900-1993). Libro de Resúmenes (eds. Liao, J. C., Gámez Vintaned, J.A.; Valenzuela-Ríos, J.I. & García-Fórner, A.) Universitat de València, Valencia; Sociedad Española de Paleontología, Madrid, 217-220.
- Gasparini, J. L., Sazima, I. (1996): A stranded melonheaded whale, *Peponocephala electra*, in southeastern Brazil, with comments on wounds from the cookiecutter shark *Isistius brasiliensis*. *Marine Mammal Science* 12, 308-312. doi: 10.1111/j.1748-7692.1996.tb00582.x
- Ginter, M., Hampe, O., Duffin, C. J. (2010): *Chondrichthyes. Paleozoic Elasmobranchii: Teeth. Handbook of Paleoichthyology*. Verlag Dr. Friedrich Pfeil, München: 168 p.
- Gray, J. E. (1851): *List of the specimens of fish in the collection of the British Museum. Part 1. Chondropterygii*. British Museum (Natural History), London: 160 p.
- Grover, C. A. (1974): Juvenile denticles of the swell shark *Cephaloscyllium ventriosum*: function in hatching. *Canadian Journal of Zoology* 52, 359–363. doi: 10.1139/z74-043
- Heithaus, M. R. (2004): Predator–prey interactions. In: J.C. Carrier, J.A. Musick, M.R. Heithaus (eds), *Biology of Sharks and their Relatives* CRC Press, Boca Raton, FL, USA: 487–521.
- Heithaus M.R., Vaudo, J.J. (2012): Predator–prey interactions. In: J.C. Carrier, J.A. Musick, M.R. Heithaus (eds), *Biology of Sharks and their Relatives* CRC Press, Boca Raton, FL, USA: 505–546.
- Johns, M. J., Barnes, C. R., Orchard M. J. (1997): Taxonomy and Biostratigraphy of Middle and Late Triassic elasmobranch ichthyoliths from northeastern British Columbia. *Geological Survey of Canada* 502, 1-235.
- Karatajuté -Talimaa, V. (1998): Determination methods for the exoskeletal remains of early vertebrates. *Mitteilungen aus dem Museum für Naturkunde in Berlin, Geowissenschaftliche Reihe* 1, 21–52. doi:10.1002/mmng.19980010103
- Leidner, A., Thies, D. (1999): Placoid scales and oral teeth of Late Jurassic elasmobranchs from Europe. *Mesozoic Fishes 2 - Systematics and Fossil Record*: 29-40.
- López-Gómez, J., Arche, A. (1992): Las unidades litoestratigráficas del Pérmico y Triásico Inferior y Medio en el sector SE de la Cordillera Ibérica. *Estudios Geológicos* 48, 123-143.
- López-Gómez, J., Márquez-Aliaga, A., Arche, A., Goy, A. (1987): La facies Muschelkalk de Henarejos (Cuenca): sedimentología y fauna del tramo superior. *Cuadernos de Geología Ibérica* 11, 665–676.
- López-Gómez, J., Mas, R., Arche, A. (1993): The evolution of the Middle Triassic (Muschelkalk) carbonate ramp in the SE Iberian Ranges, Eastern Spain: sequence stratigraphy, dolomitization processes and dynamic controls. *Sedimentary Geology* 87, 165-193.
- López-Gómez, J., Arche, A., Calvet, F., Goy, A. (1998): Epicontinental marine carbonate sediments of the Middle and Upper Triassic in the westernmost part of the Tethys Sea, Iberian Peninsula. *Zentralblatt für Geologie und Paläontologie* 9-10, 1033-1084.
- Márquez-Aliaga, A. 1985. *Bivalvos del Triásico Medio del Sector Meridional de la Cordillera Ibérica y de los Catalánides*. Colección Tesis Doctorales. Editorial de la Universidad Complutense de Madrid 40: 429 pp.
- Márquez-Aliaga, A., De Santisteban, C., Márquez, L. (1984): Triásico Medio de Bugarra. *Estudios Geológicos* 40, 365-374.
- Márquez-Aliaga, A., López Gómez, J. (1989): Paleontología y ambientes

sedimentarios del Triásico medio, Muschelkalk, de la Cordillera Ibérica 1: Cuencas y Valencia. España. *Estudios geológicos* 45, 387-398.

Márquez-Aliaga, A., Valenzuela-Ríos, J.I., Plasencia, P., Ros, S. (2004): Los fósiles del Muschelkalk (Triásico Medio) en el sector oriental de la Península Ibérica. In: E. Baquedano, S. Rubio (eds.), *Miscelánea en homenaje a Emiliano Aguirre. II: Paleontología*. Museo Arqueológico Regional de Alcalá de Henares: 276-291.

Meyer, H. von (1849): Fossile Fische aus dem Muschelkalk von Jena, Querfurt und Esperstädt. *Palaeontographica* 1, 195–208.

Muñoz-Chápuli, R. (1985): Sobre la clasificación tipológica del esqueleto dérmico de escualos. *Miscelánea Zoológica* 9, 396-400.

Pla, C., Marquéz-Aliaga, A., Botella, H. (2013): The chondrichthyan fauna from the Middle Triassic (Ladinian) of the Iberian Range (Spain). *Journal of Vertebrate Paleontology* 33, 770-785.

Pla, C., Plasencia, P., and Botella, H. (2009): Estudio preliminar de los condriictios del Ladinense (Triásico Medio) de la sección de Bugarra (Valencia, España). *Paleolusitana* 1, 383-389.

Raschi, W., Musick, J. (1986): *Hydrodynamic aspects of shark scales*. NASA Contractor Report 3963: 123 p.

Reif, W. E. (1973): Ontogenese des Hautskelettes von *Heterodontus falcifer* (Selachii) aus dem Untertithon. *Stuttgarter Beitrage zur Naturkunde* 7, 1–16.

Reif, W. E. (1974): Morphogenese und Musterbildung des Hautzähnen-Skelettes von *Heterodontus*. *Lethaia* 7, 25–42. doi:10.1111/j.1502-3931.1974.tb00882.x

Reif, W. E. (1978): Types of morphogenesis of the dermal skeleton in fossil sharks. *Paläontologische Zeitschrift* 52, 235–257. doi:10.1007/BF03006733

Reif, W. E. (1982): Morphogenesis and function of the squamation in sharks. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 164, 172–183.

Reif, W. E. (1985a): Squamation and Ecology of Sharks. *Courier Forschungsinstitut Senckenberg* 78, 1-255.

Reif, W. E. (1985b): Function of scales and photophores in mesopelagic luminescent sharks. *Acta Zoologica* 66, 111–118. doi:10.1111/j.1463-6395.1985.tb00829.x

Reif, W. E., Dinkelacker, A. (1982): Hydrodynamics of the squamation in fast swimming sharks. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 164, 184–187.

Rieppel, O. (1981): The Hybodont sharks from the Middle Triassic of Monte San Giorgio, Switzerland. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 161, 324–353.

Scheinpflug, R. (1984): Wirbeltierfunde im mainfränkischen Hauptmuschelkalk. *Aufschluss* 35, 21–36.

Soto, J., Mincarone, M. (2001): First record of kitefinshark, *Dalatias licha* (Bonnaterre, 1788) (Chondrichthyes, Dalatiidae), in the south Atlantic. *Mare Magnum* 1, 26-26.

Southall, E.J., Sims, D.W. (2003): Shark skin: a function of feeding. *Proceedings of the Royal Society of London B (Suppl.)* 270, S47-S49. doi: 10.1098/rsbl.2003.0006 1471-2954

Thies, D., Leidner, A. (2011): Sharks and guitarfishes (Elasmobranchii) from the Late Jurassic of Europe. *Palaeodiversity* 4, 63-184.

Vickers-Rich, P., Rich, T.-H., Rieppel, O., Thulbom, R. A., McClure, H. A. (1999): A Middle Triassic Vertebrata Fauna from the Jilh Formation, Saudi Arabia. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 213(2), 201–232.

APPENDIX

Percentages of original and cross-validated grouped cases correctly classified obtained using Fourier analysis (Polar and elliptical) with different number of harmonics (20, 40, 60 and 100).

Number of harmonics	Original cases			
	20	40	60	100
Polar Fourier Analysis	82 %	88 %	88 %	87 %
Elliptical Fourier Analysis	80 %	88 %	86 %	53,3 %

Number of harmonics	Cross-validated cases			
	20	40	60	100
Polar Fourier Analysis	76 %	70 %	72 %	80 %
Elliptical Fourier Analysis	64 %	68 %	66 %	42,2 %