

Supplementary data I. Geographic and faunistic data of the 100 localities used in the analysis. (1) Nearctic region; (2) Neotropical region; (3) Palaearctic region; (4) Afrotropic region; (5) Indomalaysian region.^a Cl. Climate zone (see Hernández Fernández and Peláez-Campomanes 2003). / Datos faunísticos y geográficos de las 100 localidades usadas en el análisis. (1) Neártico; (2) Neotrópico; (3) Paleártico; (4) Afrotrópico (5) Indomalaya.^a Cl. Zona climática (ver Hernández Fernández y Peláez-Campomanes 2003).

N	^a Cl.	Biogeographic Region	Locality	Country	Latitude	Longitude	Altitude	References	
								Fauna	Climate
1	I	2	Paramaribo	Suriname	5° 51' N	55° 10' W	3 m	Eisenberg, 1989	Snow, 1976
2	I	2	Puerto Limón	Costa Rica	10° 00' N	083° 03' W	-3 m	Eisenberg, 1989; Eisenberg and Redford, 1999	Canty <i>et al.</i> , 2008
3	I	2	Sao Paulo	Brasil	23° 37' S	46° 39' W	807 m	Eisenberg, 1989; Eisenberg and Redford, 1999	Rudloff, 1981
4	I	2	Tumaco	Colombia	1° 49' N	78° 47' W	4 m	Eisenberg, 1989	Snow, 1976
5	I	2	Uapes	Brasil	0° 08' S	67° 05' W	83 m	Eisenberg, 1989; Eisenberg and Redford, 1999	Meteorological Office, 1958
6	I	4	Greenville	Liberia	5° 04' N	9° 05' W	10 m	Kingdon, 1971, 1979, 1982ab, 1997; Dorst and Dandelot, 1973; Corbet, 1978; Nowak, 1999; Skinner and Chimimba, 2005; WWF, 2006	Meteorological Office, 1983
7	I	4	Kribi	Cameroon	2° 57' N	9° 54' E	624 m	Kingdon, 1971, 1974ab, 1977, 1979, 1982ab; Nowak, 1999; Dorst and Dandelot, 1973	Bultot and Griffiths, 1972
8	I	4	Yangambi	Democratic Republic of the Congo	0° 49' N	24° 29' E	487 m	Kingdon, 1971, 1974ab, 1977, 1979, 1982ab; Nowak, 1999; Dorst and Dandelot, 1973	Bultot and Griffiths, 1972
9	I	5	Medan	Indonesia	3° 40' N	98° 35' E	14 m	Corbet and Hill, 1992	Sukanto, 1969
10	I	5	Silchar	India	24° 49' N	92° 48' E	29 m	Corbet and Hill, 1992	Meteorological Office, 1966
11	II	2	Acapulco	Mexico	16° 50' N	99° 56' W	3 m	Eisenberg, 1989; Eisenberg and Redford, 1999	Meteorological Office, 1958
12	II	2	Brasilia	Brasil	15° 47' S	47° 56' W	1158 m	Eisenberg, 1989; Eisenberg and Redford, 1999	Mühr, 2007
13	II	2	Pto Ayacucho	Venezuela	5° 41' S	67° 38' W	99 m	Eisenberg, 1989	Snow, 1976
14	II	2	Roque Sáenz Peña	Argentina	26° 49' S	60° 27' W	92 m	Redford and Eisenberg, 1992	Prohaska, 1976
15	II	4	Moundou	Chad	8° 37' N	16° 04' E	420 m	Kingdon, 1971, 1974ab, 1977, 1979, 1982ab; Nowak, 1999; Dorst and Dandelot, 1973	Griffiths, 1972a
16	II	4	Mtwarra	Tanzania	10° 16' S	40° 16' E	113 m	Kingdon, 1971, 1974, 1977, 1979, 1982ab	Griffiths, 1972b
17	II	4	Ziguinchor	Senegal	12° 35' N	16° 16' W	10 m	Kingdon, 1971, 1974ab, 1977, 1979, 1982ab; Nowak, 1999; Dorst and Dandelot, 1973	Griffiths, 1972a

Supplementary data I. Geographic and faunistic data of the 100 localities used in the analysis. (1) Nearctic region; (2) Neotropical region; (3) Palaearctic region; (4) Afrotropic region; (5) Indomalaysian region. ^a Cl. Climate zone (see Hernández Fernández and Peláez-Campomanes 2003). / Datos faunísticos y geográficos de las 100 localidades usadas en el análisis. (1) Neártico; (2) Neotrópico; (3) Paleártico; (4) Afrotrópico (5) Indomalaya. ^a Cl. Zona climática (ver Hernández Fernández y Peláez-Campomanes 2003).

N	^a Cl.	Biogeographic Region	Locality	Country	Latitude	Longitude	Altitude	References	
								Fauna	Climate
18	II		5	Patna	25° 37' N	85° 10' E	53 m	Corbet and Hill, 1992	Rao, 1981
19	II		5	Phnom Penh	11° 33' N	104° 55' E	12 m	Corbet and Hill, 1992	Nieuwolt, 1981
20	II		5	Trivandrum	8° 29' N	18° 32' E	64 m	Corbet and Hill, 1992	Rao, 1981
21	II/III		1	Río Verde	21° 56' N	99° 59' W	987 m	Hall, 1981	Meteorological Office, 1980
22	II/III		1	Santiago	23° 28' N	109° 43' W	125 m	Hall, 1981	Canty <i>et al.</i> , 2008
23	II/III		2	Catamarca	28° 26' S	65° 46' W	547 m	Redford and Eisenberg, 1992	Prohaska, 1976
24	II/III		2	Las Piedras	11° 42' N	70° 12' W	15 m	Eisenberg, 1989	Snow, 1976
25	II/III		2	Remanso	9° 41' S	42° 04' W	411 m	Eisenberg, 1989; Eisenberg and Redford, 1999	Rudloff, 1981
26	II/III		4	Gaborone	24° 41' S	25° 55' E	983 m	Kingdon 1971, 1979, 1982ab, 1997; Dorst and Dandelot, 1973; Corbet, 1978; Nowak, 1999; Skinner and Chimimba, 2005	Meteorological Office, 1983
27	II/III		4	Tulear	23° 23' S	43° 44' E	9 m	Garbutt, 2007	Griffiths and Ranaivoson, 1972
28	II/III		4	Voi	3° 24' S	38° 34' E	560 m	Kingdon, 1971, 1974ab, 1977, 1979, 1982ab	Griffiths, 1972b
29	II/III		4	Zinder	13° 48' N	8° 59' E	510 m	Kingdon, 1971, 1974ab, 1977, 1979, 1982ab; Nowak, 1999; Dorst and Dandelot, 1973	Griffiths, 1972a
30	II/III		5	Jaipur	26° 49' N	75° 48' E	390 m	Corbet and Hill, 1992	Rao, 1981
31	III		1	El Paso	31° 48' N	106° 24' W	1194 m	Hall, 1981	Meteorological Office, 1980
32	III		1	Phoenix	33° 26' N	112° 01' W	340 m	Hall, 1981	Court, 1974
33	III		2	Arica	18° 28' S	70° 22' W	29 m	Redford and Eisenberg, 1992	Miller, 1976
34	III		2	Mendoza	32° 53' S	68° 49' W	800 m	Eisenberg, 1989; Eisenberg and Redford, 1992	Meteorological Office, 1958
35	III		3	Assuan	23° 58' N	33° 40' E	194 m	Corbet, 1978	Estienne and Godard, 1970
36	III		3	Jacobabad	28° 17' N	68° 29' E	57 m	Corbet and Hill, 1992	Meteorological Office, 1966

Supplementary data I. Geographic and faunistic data of the 100 localities used in the analysis. (1) Nearctic region; (2) Neotropical region; (3) Palaearctic region; (4) Afrotropic region; (5) Indomalaysian region. ^a Cl. Climate zone (see Hernández Fernández and Peláez-Campomanes 2003). / Datos faunísticos y geográficos de las 100 localidades usadas en el análisis. (1) Neártico; (2) Neotrópico; (3) Paleártico; (4) Afrotrópico (5) Indomalaya. ^a Cl. Zona climática (ver Hernández Fernández y Peláez-Campomanes 2003).

N	^a Cl.	Biogeographic Region	Locality	Country	Latitude	Longitude	Altitude	References	
								Fauna	Climate
37	III	3	Muscat	Oman	23° 37' N	58° 35' E	4 m	Corbet, 1978	Taha <i>et al.</i> , 1981
38	III	3	Smara	Western Sahara	26° 44' N	11° 26' W	140 m	Corbet, 1978	Font Tullot, 1955
39	III	4	Galcaio	Somalia	6° 46' N	47° 26' E	240 m	Kingdon, 1971, 1979, 1982ab, 1997; Dorst and Dandelot, 1973; Corbet, 1978; Nowak, 1999; Skinner and Chimimba, 2005; WWF, 2006	Meteorological Office 1983
40	III	4	Luderitz Bay	Namibia	26° 38' S	15° 06' E	23 m	Kingdon, 1971, 1974ab, 1977, 1979, 1982ab; Nowak, 1999; Dorst and Dandelot, 1973	Schulze, 1972
41	IV	1	Fresno	USA	36° 46' N	119° 42' W	100 m	Hall, 1981	Court, 1974
42	IV	1	San Diego	USA	32° 44' N	117° 10' W	4 m	Hall, 1981; Smithsonian National Museum of Natural History, 2006	Meteorological Office, 1980
43	IV	2	Santiago de Chile	Chile	33° 27' S	70° 42' W	520 m	Redford and Eisenberg, 1992	Miller, 1976
44	IV	3	Aleppo	Syria	36° 11' N	37° 13' E	395 m	Corbet, 1978	Taha <i>et al.</i> , 1981
45	IV	3	Esfahan	Iran	32° 37' N	51° 40' E	1598 m	Corbet, 1978	Taha <i>et al.</i> , 1981
46	IV	3	Potenza	Italy	40° 38' N	15° 46' E	823 m	Mitchell-Jones <i>et al.</i> , 1999	Meteorological Office, 1972
47	IV	3	Sanlúcar de Barrameda	Spain	36° 47' N	6° 21' W	30 m	Corbet, 1978	Elías and Ruiz, 1977
48	IV	3	Tripoli	Greece	37° 31' N	22° 21' E	661 m	Mitchell-Jones <i>et al.</i> , 1999	Meteorological Office, 1972
49	IV	3	Tunis	Tunisia	36° 50' N	10° 14' E	3 m	Corbet, 1978	Meteorological Office, 1983
50	IV	4	Cape Town	South Africa	33° 54' S	18° 32' E	17 m	Skinner and Chimimba, 2005	Schulze, 1972
51	V	1	Jacksonville	USA	30° 25' N	81° 39' W	7 m	Hall, 1981; Smithsonian National Museum of Natural History, 2006	Meteorological Office, 1980
52	V	1	New Orleans	USA	29° 57' N	90° 04' W	3 m	Hall, 1981	Court, 1974
53	V	2	Corrientes	Argentina	27° 28' S	58° 49' W	60 m	Redford and Eisenberg, 1992	Prohaska, 1976
54	V	2	Montevideo	Uruguay	34° 52' S	56° 12' W	22 m	Redford and Eisenberg, 1992	Meteorological Office, 1958

Supplementary data I. Geographic and faunistic data of the 100 localities used in the analysis. (1) Nearctic region; (2) Neotropical region; (3) Palaearctic region; (4) Afrotropic region; (5) Indomalaysian region. ^a Cl. Climate zone (see Hernández Fernández and Peláez-Campomanes 2003). / Datos faunísticos y geográficos de las 100 localidades usadas en el análisis. (1) Neártico; (2) Neotrópico; (3) Paleártico; (4) Afrotrópico (5) Indomalaya. ^a Cl. Zona climática (ver Hernández Fernández y Peláez-Campomanes 2003).

N	^a Cl.	Biogeographic Region	Locality	Country	Latitude	Longitude	Altitude	References	
								Fauna	Climate
55	V	2	Puerto Montt	Chile	41° 28' S	72° 57' W	13 m	Redford and Eisenberg, 1992	Miller, 1976
56	V	3	Fuzhou	China	26° 05' N	119° 18' E	88 m	Corbet and Hill, 1992	Watts, 1969
57	V	3	Kagoshima	Japan	31° 34' N	130° 33' E	4 m	Corbet, 1978	Arakawa and Taga, 1969
58	V	3	Pingnan	China	23° 23' N	110° 03' E	39 m	Corbet and Hill, 1992	Canty <i>et al.</i> , 2008
59	V	3	Shaoguan	China	55° 45' N	37° 34' E	156 m	Corbet and Hill, 1992	Canty <i>et al.</i> , 2008
60	V	4	East London	South Africa	33° 02' S	27° 52' E	125 m	Skinner and Smithers, 1990; Skinner and Chimimba, 2005	Schulze, 1972
61	VI	1	Cleveland	USA	41° 24' N	81° 51' W	237 m	Hall, 1981	Court, 1974
62	VI	1	Colorado Springs	USA	38° 49' N	104° 43' W	1882 m	Hall, 1981	Meteorological Office, 1980
63	VI	1	Prince Rupert	Canada	54° 17' N	136° 23' W	16 m	Hall, 1981	Hare and Hay, 1974
64	VI	1	St Louis	USA	38° 45' N	90° 23' W	163 m	Hall, 1981; Smithsonian National Museum of Natural History, 2006	Meteorological Office, 1980
65	VI	2	Evangelistas	Chile	52° 24' S	75° 06' W	55 m	Redford and Eisenberg, 1992	Miller, 1976
66	VI	3	Belgrado	Serbia	44° 48' N	20° 28' E	132 m	Mitchell-Jones <i>et al.</i> , 1999	Meteorological Office, 1972
67	VI	3	Blagoveschenk s	Russia	50° 15' N	106° 30' E	142 m	Corbet, 1978	Meteorological Office, 1966
68	VI	3	Moscow	Russia	55° 45' N	37° 34' E	156 m	Mitchell-Jones <i>et al.</i> , 1999	Meteorological Office, 1972
69	VI	3	Tsingtao	China	36° 04' N	120° 19' E	77 m	Corbet, 1978	Watts, 1969
70	VI	3	Vlissingen	Netherlands	51° 27' N	3° 36' E	12 m	Corbet, 1978	Arléy, 1970
71	VII	1	Medicine Lake	USA	48° 29' N	104° 27' W	595 m	Hall, 1981; Smithsonian National Museum of Natural History, 2006	Meteorological Office, 1980
72	VII	1	Rapid City	USA	44° 01' N	103° 03' W	965 m	Hall, 1981	Court, 1974
73	VII	1	Santa Fé	USA	35° 40' N	105° 55' W	2195 m	Smithsonian National Museum of Natural History, 2006	Meteorological Office, 1980
74	VII	1	Winnemucca	USA	40° 54' N	117° 48' W	1434 m	Hall, 1981	Meteorological Office, 1980

Supplementary data I. Geographic and faunistic data of the 100 localities used in the analysis. (1) Nearctic region; (2) Neotropical region; (3) Palaearctic region; (4) Afrotropic region; (5) Indomalaysian region. ^a Cl. Climate zone (see Hernández Fernández and Peláez-Campomanes 2003). / Datos faunísticos y geográficos de las 100 localidades usadas en el análisis. (1) Neártico; (2) Neotrópico; (3) Paleártico; (4) Afrotrópico (5) Indomalaya. ^a Cl. Zona climática (ver Hernández Fernández y Peláez-Campomanes 2003).

N	^a Cl.	Biogeographic Region	Locality	Country	Latitude	Longitude	Altitude	References	
								Fauna	Climate
75	VII	2	Maquinchao	Argentina	41° 15' S	68° 43' W	888 m	Redford and Eisenberg, 1992	Canty <i>et al.</i> , 2008
76	VII	2	Puerto Sta Cruz	Argentina	50° 01' S	68° 32' W	12 m	Redford and Eisenberg, 1992	Prohaska, 1976
77	VII	3	Almaty	Kazakhstan	43° 16' N	76° 53' E	775 m	Corbet, 1978	Meteorological Office, 1966
78	VII	3	Fort Shevchenko	Kazakhstan	44° 33' N	50° 17' E	23m m	Corbet, 1978	Lydolph, 1977
79	VII	3	Paotou	China	40° 34' N	109° 50' E	1044 m	Corbet, 1978	Watts, 1969
80	VII	3	Urumqi	China	43° 47' N	87° 37' E	912 m	Corbet, 1978	Watts, 1969
81	VIII	1	Edmonton	Canada	53° 34' N	113° 31 W	676 m	Hall, 1981; Smithsonian National Museum of Natural History, 2006	Meteorological Office, 1980
82	VIII	1	Fairbanks	USA	64° 49' N	147° 52' W	133 m	Hall, 1981	Hare and Hay, 1974
83	VIII	1	Fort Smith	Canada	60° 01' N	11° 58' W	62 m	Hall, 1981	Hare and Hay, 1974
84	VIII	1	Gaspé	Canada	48° 50' N	64° 29' W	28 m	Hall, 1981; Smithsonian National Museum of Natural History, 2006	Meteorological Office, 1980
85	VIII	1	Smooky falls	Canada	46° 27' N	79° 55' W	227 m	Hall, 1981; Smithsonian National Museum of Natural History, 2006	Meteorological Office, 1980
86	VIII	3	Erbogachen	Russia	61° 16' N	108°01' E	287 m	Corbet, 1978	Lydolph, 1977
87	VIII	3	Kajaani	Finland	64° 17' N	27° 41' E	134 m	Corbet, 1978	Werner, 1970
88	VIII	3	Nikolayevsk-on-Amur	Russia	53° 08' N	140° 45' E	21 m	Corbet, 1978	Meteorological Office, 1966
89	VIII	3	Petropavlovsk	Russia	52° 53' N	158° 42' E	87 m	Corbet, 1978	Meteorological Office, 1966
90	VIII	3	Serov	Russia	59° 36' N	60° 32' E	132 m	Corbet, 1978	Lydolph, 1977
91	IX	1	Baker Lake	Canada	64° 18' N	96° 00' W	9 m	Hall, 1981; Smithsonian National Museum of Natural History, 2006	Meteorological Office, 1980
92	IX	1	Barrow	USA	71° 18' N	156° 47' W	7 m	Hall, 1981	Hare and Hay, 1974
93	IX	1	Cape Hope Advances	Canada	61° 05' N	69° 33' W	73 m	Hall, 1981; Smithsonian National Museum of Natural History, 2006	Meteorological Office, 1980

Supplementary data I. Geographic and faunistic data of the 100 localities used in the analysis. (1) Nearctic region; (2) Neotropical region; (3) Palaearctic region; (4) Afrotropic region; (5) Indomalaysian region. ^a Cl. Climate zone (see Hernández Fernández and Peláez-Campomanes 2003). / Datos faunísticos y geográficos de las 100 localidades usadas en el análisis. (1) Neártico; (2) Neotrópico; (3) Paleártico; (4) Afrotrópico (5) Indomalaya. ^a Cl. Zona climática (ver Hernández Fernández y Peláez-Campomanes 2003).

N	^a Cl.	Biogeographic Region	Locality	Country	Latitude	Longitude	Altitude	References	
								Fauna	Climate
94	IX	1	Coopermine	Canada	67° 49' N	115° 05' W	9 m	Hall, 1981; Smithsonian National Museum of Natural History, 2006	Meteorological Office, 1980
95	IX	1	Port Harrison	Canada	58° 27' N	78° 08' W	6 m	Hall, 1981	Hare and Hay, 1974
96	IX	3	Bulun	Russia	70° 45' N	127° 47' E	35 m	Corbet, 1978	Meteorological Office, 1966
97	IX	3	Malye-Karmakuly	Russia	72° 23' N	55° 44' E	16 m	Corbet, 1978	Lydolph, 1977
98	IX	3	Mys Chelyuskin	Russia	77° 43' N	104° 17' E	6 m	Corbet, 1978	Lydolph, 1977
99	IX	3	Mys Scmidtta	Russia	68° 55' N	179° 17' E	6 m	Corbet, 1978	Lydolph, 1977
100	IX	3	Nizhne-Kolymsk	Russia	68° 32' N	160° 59' E	5 m	Corbet, 1978	Meteorological Office, 1966

SUPPLEMENTARY DATA I REFERENCES

- Arakawa, H., Taga, S. (1969): Climate of Japan. In: H. Arakawa (ed.), *Climates of Northern and Eastern Asia. World survey of climatology, Vol. 8*. Elsevier Co, Amsterdam: 119-158.
- Arléry, R. (1970): The climate of France, Belgium, The Netherlands and Luxembourg. In: C.C. Wallen (ed.), *Climates of Northern and Western Europe. World survey of climatology. Vol. 5*. Elsevier Co, Amsterdam: 135-194.
- Bultot, F., Griffiths, J.F. (1972): The Equatorial Wet zone. In: J.F. Griffiths (ed.), *Climates of Africa. World survey of climatology. Vol. 10*. Elsevier Co, Amsterdam: 259-312.
- Canty, J.L., Canty, W.S., Breitbart, S., Frischling, D.A. (2008): Weatherbase. www.weatherbase.com
- Corbet, G.B. (1978): *The Mammals of the Palearctic Region: a Taxonomic Review*. British Museum (Natural History) and Cornell University Press, London, UK and Ithaca, USA: 314 p.
- Corbet, G.B., Hill, J.E. (1992): *The Mammals of the Indomalayan Region: A Systematic Review*. Natural History Museum Publications, Oxford University Press, Oxford: 496 p.
- Court, A. (1974): The climate of the conterminous United States. In: R.A. Bryson, F.K. Hare (eds.), *Climates of North America. World survey of climatology. Vol. 11*. Elsevier Co, Amsterdam: 193-343.
- Dorst, J., Dandelot, P. (1973): *A Field Guide to the Larger Mammals of Africa*. Collins, London: 287 p.
- Eisenberg, J.F. (1989): *Mammals of the Neotropics. The Northern Neotropics. Vol. 1*. The University of Chicago Press, Chicago: 550 p.
- Eisenberg, J.F., Redford, K.H. (1999): *Mammals of the Neotropics. The Central Neotropics. Vol. 3*. The University of Chicago Press, Chicago: 624 p.
- Elías, F., Ruiz, L. (1977): *Agroclimatología de España*. INIA, Ministerio de Agricultura, Madrid: 29 p.
- Estienne, P., Godard, A. (1970): *Climatologie*. Librairie Armand Colin, Paris: 367 p.
- Font Tullot, Y. (1955): *El clima del Sáhara (con especial referencia a la zona española)*. Consejo Superior de Investigaciones Científicas, Madrid: 120 p.
- Garbutt, N. (2007): *Mammals of Madagascar. A Complete Guide*. A&C Black, London: 304p.
- Griffiths, J.F. (1972a): Semi-arid zones. In: J.F. Griffiths (ed.), *Climates of Africa. World survey of climatology, Vol. 10*. Elsevier Co, Amsterdam: 193-220.

- Griffiths, J.F. (1972b): Eastern Africa. In: J.F. Griffiths (ed.), *Climates of Africa. World survey of climatology, Vol. 10*. Elsevier Co, Amsterdam: 313-348.
- Griffiths, J.F., Ranaivoson, R. (1972): Madagascar. In: J.F. Griffiths (ed.), *Climates of Africa. World survey of climatology, Vol. 10*. Elsevier Co, Amsterdam: 461-499.
- Hall, E.R. (1981): *The mammals of North America (Volume I & II)*. John Wiley and Sons, New York: 1241 p.
- Hare, F.K., Hay, J.E. (1974): The climate of Canada and Alaska. In: R.A. Bryson, F.K. Hare (eds.), *Climates of North America. World Survey of Climatology, Vol. 11*. Elsevier Co, Amsterdam: 49-192.
- Kingdon, J. (1971): *East African mammals. An atlas of evolution in Africa, Vol. 1*. Academia Press, London: 446 p.
- Kingdon, J. (1974a): *East African mammals. An atlas of evolution in Africa, Vol. 2. Part A (insectivores and bats)*. Academic Press, London: 392 p.
- Kingdon, J. (1974b): *East African mammals. An atlas of evolution in Africa, Vol. 2. Part B (hares and rodents)*. Academic Press, London: 762 p.
- Kingdon, J. (1977): *East African mammals. An atlas of evolution in Africa, Vol. 3. Part A (Carnivores)*. Academic Press, London: 476 p.
- Kingdon, J. (1979): *East African mammals. An atlas of evolution in Africa, Vol. 3. Part B (large mammals)*. Academic Press, London: 436 p.
- Kingdon, J. (1982a): *East African mammals. An atlas of evolution in Africa, Vol. 3. Part C (bovids)*. Academic Press, London: 393 p.
- Kingdon, J. (1982b): *East African mammals. An atlas of evolution in Africa, Vol. 3. Part D (bovids)*. Academic Press, London: 746 p.
- Kingdon, J. (1997): *The Kingdon Field Guide to African Mammals*. A&C Black, London: 476 p.
- Lydolph, P.E. (1977): Climates of the Soviet Union. In: H.E. Landsberg (ed.), *Climates of the Soviet Union. World survey of climatology, Vol. 7*. Elsevier Co, Amsterdam: 1-443.
- Meteorological Office (1958): *Tables of temperature, relative humidity and precipitation for the World. Part II: Central and South America, the West Indies and Bermuda*. Her Majesty's Stationery Office, London
- Meteorological Office (1966): *Tables of temperature, relative humidity and precipitation for the world. Part V: Asia*. Her Majesty's Stationery Office, London
- Meteorological Office (1972): *Tables of temperature, relative humidity, precipitation and sunshine for the world. Part III: Europe and Azores*. Her Majesty's Stationery Office, London
- Meteorological Office (1980): *Tables of temperature, relative humidity, precipitation and sunshine for the World. Part I: North America and Greenland (including Hawaii and*

- Bermuda). Her Majesty's Stationery Office, London
- Meteorological Office (1983): *Tables of temperature, relative humidity, precipitation and sunshine for the World. Part IV: Africa, the Atlantic Ocean South of 35°N and the Indian Ocean.* Her Majesty's Stationery Office, London
- Miller, A. (1976): The climate of Chile. In: W. Schwerdtfeger (ed.), *Climates of Central and South America. World survey of climatology, Vol. 12.* Elsevier Co, Amsterdam: 113-145.
- Mitchell-Jones, A., Amori, G., Bogdanowicz, W., Krystufek, B., Reijnders, P. (1999): *The Atlas of European Mammals.* T. & A. D. Poyser, London: 484 p.
- Mühr, B. (2007): *Klimadiagramme weltweit.* www.klimadiagramme.de/index
- Nieuwolt, S. (1981): The Climates of Continental Southeast Asia. In: K. Takahashi, Y. Arakawa (eds.), *The climates of southern and western Asia. World survey of climatology. Vol. 9.* Elsevier Co, Amsterdam: 1-66.
- Nowak, R.M. (1999): *Walker's Mammals of the World. Vols. 1 y 2.* Johns Hopkins University Press, Baltimore: 2015 p.
- Prohaska, F. (1976): The climate of Argentina, Paraguay and Uruguay. In: W. Schwerdtfeger (ed.), *Climates of Central and South America. World survey of climatology. Vol. 12.* Elsevier Co, Amsterdam: 13-112
- Rao, Y.O. (1981): The climate of the Indian subcontinent. In: K. Takahashi, H. Arakawa (eds.), *Climates of Southern and Western Asia. World survey of climatology, Vol. 9.* Elsevier Co, Amsterdam: 67-182.
- Redford, K.H., Eisenberg, J.F. (1992): *Mammals of the Neotropics. The Southern Cone.* Vol. 2. The University of Chicago Press, Chicago: 460 p.
- Rudloff, W. (1981): *World-Climates, with tables of climatic data and practical suggestions.* Wissenschaftliche Verlagsgesellschaft, Stuttgart: 632 p.
- Schulze, B.R. (1972): South Africa. In: J.F. Griffiths (ed.), *Climates of Africa. World survey of climatology, Vol. 10.* Elsevier Co, Amsterdam: 501-586.
- Skinner, D., Chimimba, C.T. (2005): *The Mammals of the Southern African Subregion.* Cambridge University Press, Cambridge: 814 p.
- Skinner, J.D., Smithers, R.H.N. (1990): *The Mammals of the Southern African Subregion.* University of Pretoria, Pretoria: 771 p.
- Smithsonian National Museum of Natural History (2006): *North American Mammals.* www.mnh.si.edu/mna/main.cfm
- Snow, J.W. (1976): The climate of Northern South America. In: W. Schwerdtfeger (ed.), *Climates of Central and South America. World survey of climatology, Vol. 12.* Elsevier Co,

Amsterdam: 295-403.

Sukanto, M. (1969): Climate of Indonesia. In: H. Arakawa (ed.), *Climates of Northern and Eastern Asia. World survey of climatology, Vol. 8*. Elsevier Co, Amsterdam: 215-230 p.

Taha, M.F., Harb, S.A., Nagib, M.K., Tantawy, A.H. (1981): The climate of the Near East. In: K. Takahashi, H. Arakawa (eds.), *Climates of Southern and Western Asia. World survey of climatology, Vol. 9*. Elsevier Co, Amsterdam: 183-255.

Watts, I.E.M. (1969): Climates of China and Korea. In: H. Arakawa (ed.), *Climates of Northern and Eastern Asia. World survey of climatology, Vol. 8*. Elsevier Co, Amsterdam: 1-118 p.

Werner, T. (1970): The climate of Scandinavia. In: C.C. Wallen (ed.), *Climates of Northern and Western Europe. World survey of climatology, Vol. 5*. Elsevier Co, Amsterdam: 23-80.

World Wildlife Fund (2006): *WildFinder: Online database of species distributions*. www.worldwildlife.org/WildFinder

Supplementary data II. Faunal lists for the six fossil site included in the analysis.
 Listado faunístico para los seis yacimientos incluidos en el análisis

DEPOSITIONAL UNIT	MIDDLE UNIT			LOWER UNIT		BODY WEIGHT (LN)
	D	E	F	G		
LOCAL ZONE	E1	PA	AOPV	S	P5	P3
FOSSIL SITE						
<i>Euprox furcatus</i>					X	10,62
<i>Heteroprox larteti</i>					X	10,60
<i>H. moralesi</i>			X	X	X	10,96
<i>Procervulus dichotomus</i>	X					10,82
<i>Micromeryx flourensianus</i>			X		X	8,48
<i>M. sp. cf. M. flourensianus</i>				X		8,48
<i>Palaeomeryx magnus</i>			X		X	12,38
<i>Triceromeryx pachecoi</i>	X	X				11,92
<i>Dorcatherium crassum</i>		X				10,09
<i>Tethytragus</i> sp.	X	X	X	X	X	10,53
<i>T. langai</i>					X	10,53
<i>Conohyus simorrensis</i>			X	X		11,70
<i>Bunolistriodon lockharti</i>	X	X	X			11,73
<i>Listriodon splendens</i>					X	11,87
<i>Alicornops simorrensis</i>					X	14,64
<i>Cainotherium miocaenicum</i>	X	X				7,15
<i>Hispanotherium matritensis</i>	X	X				14,09
<i>Hoploaceratherium tetractylum</i>			X		X	14,11
<i>Lartetotherium sansaniense</i>					X	13,85
<i>Prosantorhinus douvillei</i>				X		14,26
<i>Rhinocerotidae</i> indet			X			14,07
<i>Anchitherium</i> sp.					X	11,41
<i>A. alberdidae</i>		X				11,35
<i>A. cursor</i>			X	X		11,12
<i>A. matritensis</i>	X		X			11,42
<i>A. procerum</i>					X	11,79
<i>Chalicotherium grande</i>					X	12,76
<i>Gomphotherium angustidens</i>			X	X	X	14,64
<i>Amphechinus</i> sp.				X		6,92
<i>A. cf. intermedius</i>					X	6,92
<i>Crocidosoricinae</i> indet					X	2,30
<i>Soricidae</i> indet	X		X			2,20
<i>Galerix</i> sp.			X		X	4,45
<i>G. exilis</i>				X		4,04
<i>Lagopsis verus</i>					X	5,86
<i>L. penai</i>	X	X	X	X		5,70
<i>Prolagus</i> sp.				X	X	4,50
<i>Democricetodon</i> sp.	X	X		X		3,71
<i>D. larteti</i>			X	X		3,71
<i>Megacricetodon collongensis</i>	X	X	X			2,40
<i>Megacricetodon crusafonti</i>					X	3,04
<i>Megacricetodon gersii</i>					X	2,83
<i>Megacricetodon rafaeli</i>					X	2,48
<i>Megacricetodon</i> sp. cf. <i>M. collongensis</i>					X	2,40
<i>Cricetodon soriae</i>					X	4,86
<i>Microdyromys monspeliensis</i>					X	3,09
<i>Microdyromys koenigswaldi</i>		X		X	X	3,09
<i>Microdyromys</i> sp.	X					2,20
<i>Simplomys simplicidens</i>	X	X	X			3,48
<i>Armantomys aragonensis</i>	X	X	X			4,39
<i>Armantomys jasperi</i>		X				3,50
<i>Armantomys tricristatus</i>				X	X	4,65
<i>Heteroxerus grivensis</i>			X	X		4,98
<i>Heteroxerus rubricati</i>	X	X			X	4,39
<i>Atlantoxerus blacki</i>	X	X				5,72

Supplementary data IIIa. Eigenvalues and correlation tables between the body size spectra categories and the discriminant functions obtained for global and biogeographic realms analyses

Global Discriminant Analysis

Eigenvalues				
Function	Eigenvalue	% of Variance	Cumulative %	Canonical Correlation
1	2,361	66,2	66,2	0,838
2	0,606	17	83,2	0,614
3	0,303	8,5	91,7	0,483
4	0,181	5,1	96,7	0,392
5	0,076	2,1	98,9	0,266
6	0,038	1,1	99,9	0,191
7	0,002	0,1	100	0,05

Standarized Canonical Coefficients

	Function						
	1	2	3	4	5	6	7
CAT. A	1,685	0,581	1,883	0,762	0,627	1,663	2,578
CAT. B	0,680	0,631	0,928	0,807	0,191	1,925	1,750
CAT. C	0,691	-0,076	1,105	0,611	1,140	1,341	2,008
CAT. D	0,151	0,042	1,341	0,305	-0,190	0,966	1,753
CAT. E	1,105	-0,375	0,467	-0,001	-0,024	1,290	1,133
CAT. F	0,552	0,445	0,095	-0,593	0,489	0,605	1,099
CAT. G	0,671	-0,096	-0,199	0,753	-0,053	0,283	1,111

Afrotropic Discriminant Analysis

Eigenvalues				
Function	Eigenvalue	% of Variance	Cumulative %	Canonical Correlation
1	10,478	73,3	73,3	0,955
2	3,325	23,3	96,6	0,877
3	0,492	3,4	100	0,574

Standarized Canonical Coefficients

	Function		
	1	2	3
CAT. A	2,362	1,176	0,103
CAT. B	4,008	3,579	1,036
CAT. C	1,838	0,724	0,961
CAT. D	4,442	2,681	0,921
CAT. E	0,596	-0,090	0,835
CAT. F	-0,369	1,558	0,275

Supplementary data IIIa. Eigenvalues and correlation tables between the body size spectra categories and the discriminant functions obtained for global and biogeographic realms analyses

Indomalaysian Discriminant Analysis

Eigenvalues				
Function	Eigenvalue	% of Variance	Cumulative %	Canonical Correlation
1	442,601	100	100	0,999

Standardized Canonical Coefficients	
	Function
	1
CAT. A	23,081
CAT. B	18,246
CAT. C	-9,156

Paleotropic Discriminant Analysis

Eigenvalues				
Function	Eigenvalue	% of Variance	Cumulative %	Canonical Correlation
1	1,821	58,6	58,6	0,803
2	0,856	27,6	86,2	0,679
3	0,429	13,8	100	0,548

Standardized Canonical Coefficients			
	Function		
	1	2	3
CAT. A	1,259	2,377	0,629
CAT. B	0,082	2,434	-0,132
CAT. C	0,208	2,479	0,978
CAT. D	0,611	2,195	0,197
CAT. E	-0,580	1,245	0,663
CAT. F	0,337	0,449	1,557
CAT. G	-0,792	-0,099	-0,689

Supplementary data IIb. Eigenvalues and correlation tables between the cenogram variables and the discriminant functions obtained for global and biogeographic realms analyses

Global Discriminant Analysis

Eigenvalues				
Function	Eigenvalue	% of Variance	Cumulative %	Canonical Correlation
1	4,495	63,1	63,1	0,904
2	1,33	18,7	81,7	0,756
3	0,619	8,7	90,4	0,618
4	0,325	4,6	95	0,495
5	0,203	2,8	97,8	0,41
6	0,079	1,1	98,9	0,27
7	0,037	0,5	99,4	0,188
8	0,03	0,4	99,9	0,172
9	0,01	0,1	100	0,099

Standarized Canonical Coefficients

	Function								
	1	2	3	4	5	6	7	8	9
W1	0,511	-0,449	-0,227	0,523	0,559	0,253	0,223	0,100	-0,256
W2	-0,018	-0,972	0,608	-0,196	-1,178	-0,031	0,486	0,051	-0,662
W5	-0,015	0,296	0,014	0,271	0,303	0,250	0,447	0,188	0,137
P1	0,015	0,053	0,173	0,033	0,075	-0,372	-0,038	0,447	-0,074
P5	1,455	1,017	0,011	0,069	-0,444	0,206	1,083	0,109	0,699
P5-P1	-1,099	-0,678	-0,219	-0,222	0,325	0,077	-0,308	0,422	-0,162
G	-0,966	0,112	-0,551	-0,422	1,232	-0,170	0,262	0,727	0,312
mG	-0,677	0,265	0,741	-0,233	0,291	-0,957	1,357	0,690	-0,287
MG	0,835	0,133	-0,262	0,950	-1,150	1,311	-0,554	-0,581	0,680
WG	0,241	0,991	0,060	-0,493	0,499	0,338	-0,078	-0,435	-0,129
WmG	0,298	-0,450	0,779	-0,073	0,419	0,129	-0,130	-0,069	0,399
WMG	-0,146	0,367	-0,190	0,242	-0,166	0,552	-0,285	0,690	-0,442

Afrotropic Discriminant Analysis

Eigenvalues				
Function	Eigenvalue	% of Variance	Cumulative %	Canonical Correlation
1	204,547	85,5	85,5	0,998
2	27,908	11,7	97,2	0,983
3	6,716	2,8	100	0,933

Standarized Canonical Coefficients

	Function		
	1	2	3
W1	1,530	-1,417	0,031
W2	21,974	1,133	-2,185
W5	-4,590	-4,904	0,304
P1	-14,198	-12,018	3,507
P5	1,222	16,715	-0,839
G	7,554	5,647	1,239
mG	-0,874	6,409	4,639
MG	4,357	-6,061	-4,735

Supplementary data IIb. Eigenvalues and correlation tables between the cenogram variables and the discriminant functions obtained for global and biogeographic realms analyses

Indomalaysian Discriminant Analysis

Eigenvalues				
Function	Eigenvalue	% of Variance	Cumulative %	Canonical Correlation
1	0,573	100	100	0,604

Standarized Canonical Coefficients	
	Function
	1
W1	0,415
W2	0,678
W5	-0,764

Paleotropic Discriminant Analysis

Eigenvalues				
Function	Eigenvalue	% of Variance	Cumulative %	Canonical Correlation
1	13,813	68	68	0,966
2	4,24	20,9	88,9	0,9
3	2,263	11,1	100	0,833

Standarized Canonical Coefficients			
	Function		
	1	2	3
W1	3,739	1,888	1,593
W2	-1,168	-0,085	1,490
W5	2,736	3,552	0,888
P1	-1,518	3,482	-1,557
P5	1,595	-6,404	-0,453
G	1,574	0,935	1,841
mG	4,510	-0,172	1,040
MG	-5,352	0,182	-2,508
WG	2,551	3,769	0,509
WmG	2,910	1,341	0,619
WMG	-1,062	0,658	-0,310