

Cousinia khowiensis (Asteraceae, Cardueae), a new species from north-west of Iran

Farideh Attar

Central herbarium of Tehran University, School of Biology, College of Science, University of Tehran, Tehran, Iran ✉

Mansour Mirtadzadini

Department of Biology, Faculty of Science, Shahid Bahonar University of Kerman, Kerman, Iran ✉

Neda AtazadehDepartment of Biology, Faculty of Science, Yasouj University, Yasouj, Iran ✉✉ 

Corresponding author

<https://doi.org/10.5209/mbot.101482>

Received: 6 March 2025 / Accepted: 29 September 2025 / Published online: 24 February 2026

Abstract. A new endemic species (Asteraceae, Cardueae) is described as *Cousinia khowiensis* on the basis of specimens collected in West Azarbaijan Province, Iran. The distribution area of the new species belongs to the distribution range of the species of sect. *Cousinia* in the northwest of Iran. It is also morphologically similar to sect. *Cousinia*. These similarities are mainly in some characters such as the size and shape of the involucre, the number of flowers, the height of the plant and the texture of the leaves. The main differences between the new species and the members of sect. *Cousinia* in Iran are: the shape of the leaves, the number of phyllaries, the size of the capitula and the colour of the corolla. *Cousinia khowiensis* is compared with the closest species, including *C. microcephala*, *C. sivasica*, *C. woronowii* and *C. calolepis*. *C. sivasica* and *C. woronowii* of sect. *Cousinia* are distributed in Turkey, and *C. calolepis* belongs to sect. *Stenocephalae*. The holotype and isotype specimens of *Cousinia khowiensis* are illustrated, together with close-ups of various parts of the new species. An identification key for all taxa of sect. *Cousinia* in Iran as well as *C. sivasica* and *C. woronowii*, which is distributed in Turkey, has been provided. The number of species in the sect. *Cousinia* in Iran, including the new species, is 12. The relationship of *Cousinia khowiensis* and the related taxa was presented separately. In addition, a distribution map showing the distribution range of *C. khowiensis* and the related species is presented here.

Keywords: *Cousinia*, sect. *Cousinia*, flora of Iran, endemic.

How to cite: Attar, F., Mirtadzadini, M. & Atazadeh, N. 2026. *Cousinia khowiensis* (Asteraceae, Cardueae), a new species from north-west of Iran. *Mediterr. Bot.* 47(1), e101482. <https://doi.org/10.5209/mbot.101482>

1. Introduction

The genus *Cousinia* Cass. belongs to the tribe Cardueae (Asteraceae) and is one of the most diverse genera in Central and Southwest Asia with about 665 species (Atazadeh & Ghahremaninejad, 2025; POWO, 2025), after *Senecio* L. (c. 1500 species) and *Vernonia* Schreb. (c. 1000 species) (Tscherneva, 1962; Rechinger, 1972, 1979; Frodin, 2004; Attar & Ghahreman, 2006; Susanna & Garcia-Jacas, 2007; Attar & Djavadi, 2010; Mehregan & Assadi, 2016; Minaeifar *et al.*, 2016; Rastegar *et al.*, 2017, 2018; Attar & Amini Rad, 2019; Atazadeh *et al.*, 2021). The greatest diversity and distribution of this genus is found in the mountainous regions of Iran, Afghanistan and Turkmenistan, with almost 400 species in Southwest Asia, 379 of which endemic (Rechinger, 1986; Knapp, 1987; Attar & Djavadi, 2010; Atazadeh *et al.*, 2020).

In Iran, *Cousinia* is mainly distributed in the Kopet Dagh, Elborz, Zagros Mountains and in the north-western regions of the country (Attar & Djavadi, 2010). In the Flora Iranica (Rechinger, 1972, 1979), *Cousinia* was artificially divided into 71 sections, 45 of which are recorded in Iran (Rechinger, 1986; Attar & Ghahreman, 2002; Mehregan & Assadi, 2010; Attar *et al.*, 2024), including the sect. *Cousinia* with about 67 species of which 11 species are distributed in the northwest of Iran (Rechinger, 1970). Only a few species are distributed in the adjacent countries of Turkey and the Caucasus region (Tscherneva, 1962; Huber-Morath, 1975).

The members of the sect. *Cousinia* are more or less small plants with branched stems, with almost leathery or rarely herbaceous leaves, decurrent or not decurrent stem leaves, small or medium-sized capitula, with (5-) 10-45 flowers and a yellow or pink to purple corolla. The involucre is ovoid-oblong to broadly ovoid, rarely globose, densely

arachnoid-woolly to glabrous, with small and narrow or subulate exappendiculate, erect, spreading or recurved and falcate phyllaries.

According to the Flora Iranica (Rechinger, 1972, 1979), the genus *Cousinia* comprises about 220 species in the flora of Iran. However, in recent studies (Djavadi & Attar, 2006; Djavadi *et al.*, 2007; Attar *et al.*, 2007, 2016; Assadi, 2009, 2010, 2011; Attar & Mirtadzadini, 2009; Attar & Djavadi, 2010; Mehregan & Assadi, 2010; Mehregan *et al.*, 2010; Attar, 2011; Mirtadzadini & Attar, 2014; Rastegar *et al.*, 2018; Attar & Amini Rad, 2019; Attar *et al.*, 2022; Attar *et al.*, 2024) several additional species of *Cousinia* have been described and recorded in the flora of Iran. Therefore, the number of species of the genus is now estimated to be about 271, which still makes it the second largest genus after *Astragalus* L. in the flora of Iran (Rastegar *et al.*, 2018).

Two interesting specimens were collected in the province of West Azarbaijan as part of the *Cousinia* study project. These specimens were included in the *Cousinia* section due to their morphological similarities and their distribution range. After careful examination, it was determined that these specimens have distinct characteristics and can therefore be described as a new species. This newly described species, characterized by several characters including ovoid capitula with less than 20 flowers, longly decurrent leaves and exappendiculate phyllaries, distinguished as the only member of sect. *Cousinia* with a purple corolla in the flora of Iran.

2. Material and Methods

The specimens of the new species were discovered during a trip to northwestern Iran and identified on the basis of morphological characters. We have carried out thorough and accurate botanical work in the province of West Azarbaijan to identify the new species. After the field investigations, we collected two specimens from the existing individuals of the new species at this location and examined them as holotype and isotype. To confirm the identity, the specimens were compared with other specimens and/or images in virtual herbaria such as TUH, TARI, IRAN, K, B, M, E and W (herbaria acronyms follow Thiers 2024+). In addition, various relevant literature sources were consulted, including Flora Iranica (Rechinger, 1972, 1979), Flora of the USSR (Tscherneva, 1962), Flora of Turkey (Huber-Morath, 1975) and related publications.

3. Results and Discussion

Cousinia khoiensis Attar, Mirtadz. & Atazadeh, *sp. nov.* (Figures 1–3)

Type: Iran, West Azerbaijan province: Khoi, Valadian, 13.06.1997 Ghahreman & Attar, TUH20583 (Holotype & Isotype).

Perennial, ca. 20 cm high; rootstock woody with densely cottony hairs and the remnants of old petioles at the base. Stem grayish, covered with milky-white arachnoid hairs, divaricately branched from the base, leafy throughout, Lower branches

longer than the uppers. Basal leaves ovate-elliptic, 3–5 cm long and 1–3 cm wide, green, densely white arachnoid on both surfaces terminating to spines with length 5 mm; petiole 6 mm long; stem leaves ovate-elliptic, 2.5–3 cm long and 1–1.5 cm wide, densely white arachnoid on both surfaces, longly decurrent. Capitula 15–20 flowered, peduncle 3–5 mm long; involucre ovoid, arachnoid, ca. 10 mm in diameter; phyllaries 32–44, exappendiculate; the outer lanceolate-linear, spreading, 3–4 mm long and 1 mm wide; the median similar to the outer but longer, spreading, 6–7 mm long and 1 mm wide, narrow and acuminate toward apex; the inner 4 mm long and 1 mm wide, erect, imbricate, dark purple on the outer surface, obtuse; the innermost longer than the inner, brownish, imbricate. Corolla purple, 11–12 mm long, limb 5–6 mm, tube 6 mm; anthers purple, glabrous, concolor. Receptacle bristles smooth. Achenes immature.

Etymology: The epithet refers to the type locality, the city of Khoi in Western Azerbaijan Province.

Phenology: Flowering: June–August

Taxonomic note

The new species is found in the northwestern Iran within the distribution range of the species of sect. *Cousinia* and is morphologically similar to them. These similarities are mainly in the size and shape of the involucre, the number of flowers, the height of the plant and the texture of the leaves. Of them, *C. microcephala* C.A.Mey. is the species most similar to the new species and whose distribution area coincides with the area of the new species. However, *C. microcephala* and all species of this section in this area have yellow corollas, whereas the corolla of the new species is purple. A little further away, in Turkey, species similar to the new species are distributed, whose corollas are yellow and red suffused. *C. sivasica* Hub.-Mor., which is distributed in the center of Turkey, resembles the new species in some morphological characters, such as the shape of the capitula and the arrangement of the phyllaries. *C. woronowii* Bornm. is the other similar species occurring in northeastern Turkey and about 250 km away from the new species. Although it has leaves with deeper incisions and deflexed (not straight) phyllaries, this species and the new species have the purple corolla. Totally, this species is morphologically most closely related to the new species. Also, *C. khoiensis* due to the small involucre and purple or pink corollas has similarities with the species of sect. *Stenocephalae* Bunge, especially *C. calolepis* Boiss. However, the members of this section are distributed in the north and north-east of Iran and far from the distribution range of the new species. They also have narrower involucre with fewer florets. A comparison of the morphological characters of the new species *C. khoiensis* and its closely related species is given in Table 1. In addition, a distribution map is also provided here showing the distribution range of *C. khoiensis* and related species (Figure 4). An identification key for all taxa of sect. *Cousinia* in Iran as well as *C. sivasica* and *C. woronowii*, which is distributed in Turkey, has been provided:



Holotype

**Central Herbarium of Tehran University (TUH)
Flora of Iran**

No: 20583

Family: Asteraceae

Name: *Cousinia Khoiensis* Attar, Mirtadz. & Atazadeh, sp. nov.

Loc.: Iran. Azarbaijan: Khoi: Valadian.

Alt.: 1320 m

Date: 13.06.1997

Note: 38° 29' 15" N 45° 11' 44" E

Leg.: Ghahreman & Attar

Det.:

Figure 1. Holotype of *Cousinia khoiensis*.



Figure 3. Morphological details of *Cousinia kholensis*. A: Stem; B: Involucre and phyllaries arrangement; C: flowers; D: Innermost phyllaries. A, B and C, Scale bars= 0.1 mm; D Scale bars=0.2 mm.

Table 1. Morphological comparison of *C. kholensis*, *C. microcephala*, *C. sivasica*, *C. woronowii* and *C. calolepis*.

Characters	<i>C. kholensis</i>	<i>C. microcephala</i>	<i>C. sivasica</i>	<i>C. woronowii</i>	<i>C. calolepis</i>
Plant height (cm)	20	15-50	10-20	25-40	30-35
Stem color	grayish	light brownish	light brownish	light brownish	brownish
Leaves texture	herbaceous	leathery	leathery	leathery-herbaceous	leathery
Leaves shape	ovate-elliptic	lanceolate/pinnatisect	lanceolate/pinnatisect	linear-lanceolate	lanceolate
No of flowers	15-20	9-15	20-30	25-40	6-11
No of phyllaries	32-44	35-40	70-120	100-120	44-56
Phyllaries arrangement	spreading	strongly imbricate	spreading	spreading	imbricate-slightly recurved at tip
Capitula shape	ovoid	ovoid-strongly constricted above	ovoid	ovoid	cylindrical
Capitula size (mm)	13 × 8-10	9-11× 6-8	25-27× 18-20	23-25× 15-20	10× 5-8
Involucre diameter (mm)	8-10	6-8	18-20	15-20	5
Corolla color	purple	yellow	red/yellow	purple	purple
Corolla length (mm)	11-12	11-12	12-15	13-15	10-11
Innermost phyllaries color/indumentum on Adaxial side	white/smooth	yellow/ papillose-scabrous	white/smooth	white/smooth	purple/ papillose




Figure 4. Distribution map of *Cousinia khoiensis* (red), *C. microcephala* (blue), *C. sivasica* (pink), *C. woronowii* (light blue) and *C. calolepis* (green).

Diagnostic key to the new species and related species

1. Phyllaries subulate, recurved and falcate2
- 1' Phyllaries straight, spreading or imbricate4
2. Phyllaries 65–70 *C. tenuifolia* C.A.Mey. in DC
- 2' Phyllaries more than 100 (rarely up to 160)3
3. involucre densely arachnoid *C. wilhelmiae* Rech.f.
- 3' involucre glabrescent *C. lomakinii* C. Winkl.
4. Leaves shortly or longly decurrent5
- 4' Leaves not decurrent10
5. Involucre glabrescent. *C. chlorocephala* C.A.Mey. in DC
- 5' Involucre arachnoid6
6. Phyllaries 70–120; flowers 20–40; corolla 12–15 mm7
- 6' Phyllaries 30–408
7. Flowers 20–30; corolla, yellow or red-suffused; leaves segments linear *C. sivasica* Hub.-Mor.
- 7' Flowers 25–40; corolla red to purple; leaves segments triangular *C. woronowii* Bornm.
8. Flowers more or less 10; phyllaries more or less 60 *C. azerbaijanica* Djavadi, Attar & Najafi
- 8' Flowers more than 10; phyllaries up to 459
9. Corolla yellow; phyllaries 40–45; leaves arachnoid, green, concolored *C. tabriziana* Bunge
- 9' Corolla purple; phyllaries 32–44; leaves discolored, densely white-tomentose below *C. khoiensis* Attar, Mirtadz. & Atazadeh

10. Flowers 30–40 *C. hamosa* C.A.Mey. in DC
- 10' Flowers up to 2011
11. Phyllaries imbricate; involucre constricted above *C. microcephala* C.A.Mey. in DC
- 11' Phyllaries spreading-recurved; involucre not constricted above12
12. Involucre glabrescent *C. seidlitzii* C.A.Mey. in DC
- 12' Involucre arachnoid13
13. Leaves segments broad triangular; capitula 9–12 mm in diam. *C. hohenackeri* Fisch. & C.A.Mey.
- 13' Leaves segments linear; capitula 12–15 mm in diam. *C. urumiensis* Bornm.

Appendix. Other examined specimens

Cousinia calolepis Boiss.

Iran. Qazvin, Alamut, between Gazorkhan and Khoshkechal, 2500 m asl, Ghahreman & Mozaffarian TUH9972; Qazvin, Alamut, 30 km after Qazvin toward Rajai-Shahr, 2350 m asl, Saber & Zarrei, TUH34378; Qazvin, Alamut, Gazorkhan, TUH27626.

Cousinia microcephala C.A.Mey.

Iran. East Azerbaijan, Sufian to Shabestar, 1400 m asl, K. H. Reichinger 43600 (K2037283)
Iran. East Azerbaijan, Sufian region, Attar, Maroofi & Zamani TUH38313; Iran. East Azerbaijan, Shabestar, Sharafkhaneh, Ahmad-Khanbeygi TUH38318;
Iran. East Azerbaijan, Sufian to Shabestar,

Ahmad-Khanbeygi TUH38317; Iran. East Azerbaijan, Marand to Evoghli, 1250 m asl, Ghahreman & Mozaffarian TUH9772. Iran, Azarbaijan, Khoi, near Seidchadschi, Szovitz (G-DC, type specimen).

Cousinia sivasica Hub.-Mor.

Turkey. B6 Sivas, 36 km S Kangal, 1600 m asl. 2.8.1976, F. Sorger s.n. (W). B6 Malatya, 36 km S Kangal, E Kürecik, W der Passhöhe, 1800 m asl. 1.9.1977, F. Sorger s.n. (W).

Cousinia woronowii Bornm.

Turkey. B9 Erzurum, c. 30 km from Horasan to Tahir, Davis W47300.

References

- Assadi, M. 2009. Four new species of the genus *Cousinia* Cass. (Asteraceae) from Iran. Iranian Journal of Botany. 15: 36–44.
- Assadi, M. 2010. A Taxonomic Revision of *Cousinia* Cass. Sect. *Chaetocephalae* (Asteraceae). Iranian Journal of Botany. 16: 191–196.
- Assadi, M. 2011. Two new species of the genus *Cousinia* (Asteraceae) from Iran. Iranian Journal of Botany. 17: 6–9. doi: 10.22092/ijb.2011.101529
- Atazadeh, N., & Ghahremaninejad, F. 2025. Pollen Diversity in the Tribe Cardueae (Asteraceae) and Its Taxonomic Significance. Microscopy Research and Technique. 88: 1422–1450. doi: 10.1002/jemt.24778
- Atazadeh, N., Sheidai, M., Attar, F. & Koohdar, F. 2020: Species delimitation in the genus *Cousinia* Cass. (Family Asteraceae), sections *Cynaroideae* Bunge and *Platyacanthae* Rech. f.: morphometry and molecular analysis. Caryologia. 73(3): 3–12. doi: 10.13128/caryologia-275
- Atazadeh, N., Sheidai, M., Attar, F. & Koohdar, F. 2021. Molecular phylogeny and morphometric analyses in the genus *Cousinia* Cass. (Family Asteraceae), sections *Cynaroideae* Bunge and *Platyacanthae* Rech. Caryologia 74(4): 39–50. doi: 10.13128/caryologia-1114
- Attar, F. 2011. New taxa of *Cousinia* Cass., sect. *Stenocephalae* Bunge (Asteraceae) from Iran. Journal of Taxonomy and Biosystematics. 3: 59–68.
- Attar, F. & Amini Rad, M. 2019. *Cousinia sharifii*, a new species of *Cousinia* sect. *Cynaroideae* (Asteraceae) from NW Iran. Phytotaxa 395: 287–295. <https://doi.org/10.11646/phytotaxa.395.4.4>
- Attar, F. & Djavadi, S.B. 2010. A taxonomic revision of *Cousinia*, sect. *Cynaroides* (Asteraceae, Cardueae) in the flora of Iran. Iranian Journal of Botany. 16 (1): 130–184.
- Attar, F., Maroofi, H., Mirtadzadini, M. & Sotoodeh, A. 2022. Two new species of genus *Cousinia* (Asteraceae, Cardueae) from W and NE Iran. Nova Biologica Reperta. 9 (2): 144–152. <https://doi.org/10.52547/nbr.9.2.144>
- Attar, F., Mirtadzadini, M., Rastegar, A. & Atazadeh, N. 2024. *Cousinia meyanensis* (Asteraceae), a new species from northeastern Iran. Ann. Bot. Fenn. 61: 97–101. doi: 10.5735/085.061.0114
- Attar, F. & Ghahreman A. 2002. New taxa of genus *Cousinia* (Compositae) from Iran. Iranian Journal of Botany. 9 (2): 161–169.
- Attar, F. & Ghahreman A. 2006. A synopsis of sect. *Cynaroides* (*Cousinia*, Compositae), distribution patterns and diversity centers. Rostaniha 7(2): 315–342.
- Attar, F., Ghahreman, A., Mahdigholi, K. & Sheidai, M. 2007. A synopsis of *Cousinia* sect. *Serratuloides* (Asteraceae) in Iran. Novon. 17: 1–7. doi: 10.3417/1055-3177(2007)17[1:ASOCSS]2.0.CO;2
- Attar, F. & Mirtadzadini, M. 2009. Two new species of *Cousinia* Cass. sect. *Pugioniferae* Bunge (Asteraceae, Cardueae), from east and southeast of Iran. Iranian Journal of Botany. 15: 146–152.
- Attar, F., Mozaffarian, V. & Mirtadzadini, M. 2016. *Cousinia brevicaulis* sp. nov. (sect. *Lasiandra*) for the flora of Iran. Nord. J. Bot. 34: 174–177. doi: 10.1111/njb.00818
- Djavadi, S.B. & Attar, F. 2006. Two new species of *Cousinia* Cass. of sect. *Stenocephalae* Bunge from Iran. Feddes Repertorium: Zeitschrift für botanische Taxonomie und Geobotanik. 117: 453–458.
- Djavadi, S.B., Attar, F. & Najafi, Z. 2007. A new species of *Cousinia* sect. *Cousinia* from NW Iran. Iranian Journal of Botany. 13 (1): 43–46.
- Frodin, D.G. 2004. History and concepts of big plant genera. Taxon. 53(3): 753–776. doi:10.2307/4135449
- Huber-Morath, A. 1975. *Cousinia*. In: Davis, P.H. (Ed.). Flora of Turkey and the East Aegean Islands, vol. 5: 329–353. Edinburgh University Press, Edinburgh.
- Knapp, H.D. 1987. On the distribution of the genus *Cousinia* (Compositae). Plant Syst. Evol. 155: 15–25.
- Mehregan, I. & Assadi, M. 2010. *Cousinia* sect. *Argenteae* (Asteraceae, Cardueae), a new section including a new species from NE Iran. Willdenowia. 39: 265–271. <https://doi.org/10.3372/wi.39.39205>
- Mehregan, I. & Assadi, M. 2016. A synopsis of *Cousinia* sect. *Pseudactinia* (Cardueae, Asteraceae) including a new species from NE Iran. Phytotaxa. 257(3): 271. <https://doi.org/10.11646/phytotaxa.257.3.5>
- Mehregan, I., Djavadi, S.B. & Pahlevani, A.H. 2010. *Cousinia* (sect. *Hausknechtianae*) *karkasensis*, a new species from Karkas Mts. in central Iran. Iranian Journal of Botany. 16: 200–203.
- Minaeifar, A., Sheidai, M., Attar, F., Noormohammadi, Z. & Ghasemzadeh-Baraki, S. 2016. Biosystematic study in the genus *Cousinia* Cass. (Asteraceae), sect. *Cousinia*. Biochem. Syst. Ecol. 69: 252–260. doi: 10.1016/j.bse.2016.10.008
- Mirtadzadini, M. & Attar, F. 2014. *Cousinia kermanensis* and *C. pulcherantha* spp. nov. (Compositae, sect. *Congestae* and sect. *Leiocaulae*) from southeastern Iran. Nord. J. Bot. 32: 279–284. doi: 10.1111/njb.00159
- Rastegar, A., Ahmad, S.A. & Attar, F. 2017. *Cousinia azmarensis* (Asteraceae, Cardueae), a New Species from Kurdistan, Iraq. Harvard Papers in Botany. 22(1): 71–73. doi: 10.3100/hpib.v22iss1.2017.n8
- Rastegar, A., Attar, F. & Mirtadzadini, M. 2018. Novelties in Iranian *Cousinia* sect. *Cynaroideae* (Asteraceae, Cardueae): new taxa and taxonomic

- notes. *Phytotaxa*. 343: 227. doi: 10.11646/phytotaxa.343.3.3
- Rechinger, K. H. 1970. *Flora Iranica* 90 (Cynareae). Pp. 87–95. Akademische Druck- u. Verlagsanstalt, Graz.
- Rechinger, K.H. 1972. *Flora Iranica* 90. Compositae-Cynareae I. Cousinia. Natural History Museum, Vienna.
- Rechinger, K.H. 1979. Cousinia. In: Rechinger K.H. (Ed.). *Flora Iranica: Compositae III-Cynareae*, vol. 139°. Pp. 108–153. Natural History Museum, Vienna.
- Rechinger, K.H. 1986. Cousinia: morphology, taxonomy, distribution and phytogeographical implication. *Proc. R. Soc. Edin.* B 89: 45–58.
- Susanna, A. & Garcia-Jacas, N. 2007. Tribe Cardueae Cass. 1819. In: Kadereit J.W., Jeffrey C. (Ed.), *The families and genera of vascular plants, Flowering plant, Eudicots, Asterales*, vol. 8: 123–147. Springer Verlag, Berlin.
- Tscherneva, OV. [Cherneva OV]. 1962. Compositae-Cousinia in: Shishkin BK, Bobrov EG, eds. *Flora SSSR*. 27: 108–357. Moscow and Leningrad [English translation: Enfield. 1993].

Websites

- POWO. 2025. Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. <http://www.plantsoftheworldonline.org>