

Supplementary Material

Hurtado, P., Aragón, G., Martínez, I., Mayrhofer, H. & Prieto, M. 2023. The epiphytic lichens on *Fagus sylvatica* in beech forests of Europe: towards an open and dynamic checklist. *Mediterr. Bot.* 44, e84299.

<https://doi.org/10.5209/mbot.84299>

S1. List of primary studies included in this review.

Adamčík, S., Aude, E., Bässler, C., Christensen, M., van Dort, K., Fritz, Ö., Glejdura, S., Heilmann-Clausen, J., Holec, J., Jančovičová, S., Kunca, V., Lackovičová, A., Lüth, M. & Ódor, P. 2016. Fungi and lichens recorded during the Cryptogam Symposium on Natural Beech Forests, Slovakia 2011. *Czech Mycol.* 68: 1–40.

Amo, G. & Burgaz, A.R. 2004. *Sclerophora peronella* new to the Iberian Peninsula. *Lichenologist* 36: 265–267.

Aragón, G., Abuja, L., Belinchón, R. & Martínez, I. 2015. Edge type determines the intensity of forest edge effect on epiphytic communities. *Eur. J. Forest Res.* 134: 443–451.

Aragón, G., Martínez, I. & García, A. 2012. Loss of epiphytic diversity along a latitudinal gradient in southern Europe. *Sci. Total Environ.* 426: 188–195.

Bartoli, A., Castello, M. & Nimis, P.L. 1991. Phytogeographical outline of epiphytic lichens in southern Calabria (Italy). *Bot. Chron.* 10: 936–943.

Benesperi, R., Brunialti, G., Samari Fappiano, A., Frati, L., Giordani, P., Loppi, S., Paoli, L., Ravera, S., Buzio, P., Benco, C., Martellos, S. & Tretiach, M. 2007. Contributo alla conoscenza della flora lichenica dell'Abetone, Appennino Tosco-Emiliano. *Inf. Bot. It.* 39: 43–51.

Berger, F., Breuss, O., Maliček, J. & Türk, R. 2018. Lichens in the primeval forest areas 'Großer Urwald' and 'Kleiner Urwald' (Rothwald, 'Dürrenstein Wilderness Area', Lower Austria, Austria). *Herzogia* 31: 716–731.

Bilovitz, P.O. 2014. The importance of old deciduous trees and wooden fences for lichen diversity - an example from the Teichalm area (Eastern Alps). *Herzogia* 27: 199–204.

Bilovitz, P.O., Arup, U. & Mayrhofer, H. 2010. Contribution to the lichen biota of Slovenia XII. Some lichens from Logarska dolina. *Hladnikia* 26: 15–20.

Bilovitz, P.O., Batič, F. & Mayrhofer, H. 2011. Epiphytic lichen mycota of the virgin forest reserve Rajhenavski Rog (Slovenia). *Herzogia* 24: 315–324.

Bilovitz, P.O., Stešević, D. & Mayrhofer, H. 2010. Epiphytic lichens and lichenicolous fungi from the northern part of Montenegro. *Herzogia* 23: 249–256.

- Boch, S., Fischer, M. & Prati, D. 2015. To eat or not to eat - relationship of lichen herbivory by snails with secondary compounds and field frequency of lichens. *J. Plant. Ecol.* 8: 642–650.
- Boch, S., Müller, J., Prati, D., Blaser, S. & Fischer, M. 2013. Up in the tree - The overlooked richness of bryophytes and lichens in tree crowns. *PLoS ONE* 8: e84913.
- Boch, S., Prati, D. & Fischer, M. 2016. Gastropods slow down succession and maintain diversity in cryptogam communities. *Ecology* 97: 2184–2191.
- Burgaz, A.R., Fuertes, E. & Escudero, A. 1994. Ecology of cryptogamic epiphytes and their communities in deciduous forests in Mediterranean Spain. *Vegetatio* 112: 73–86.
- Christensen, S.N. 1994. Lichens from Croatia. *Acta Bot. Croat.* 53: 101–113.
- Clerc, P. & Truong, C. 2008. The non-sorediate and non-isidiate *Parmelina* species (lichenized ascomycetes, Parmeliaceae) in Switzerland - *Parmelina atricha* (Nyl.) P. Clerc reinstated in the European lichen flora. *Sauteria* 15: 175–194.
- Çobanoğlu, G., Yavuz, M., Costache, I., Radu, I., Açıkgöz, B. & Baloni, L. 2009. Epiphytic and terricolous lichens diversity in Cozia National Park (Romania). *Muzeul Olteniei Craiova. Oltenia* 25: 17–22.
- Costache, I., Yavuz, M., Çobanoğlu, G., Radutoiu, D. & Radu, I. 2007. Preliminary data about the Romanian-Turkish collaboration in the study of the lichens from Cozia Mount. *Ann. Univ. Craiova (Ser. Biol.)* 12: 15–20.
- Czarnota, P. 2015. *Lecania cuprea* and *Micarea pycnidiophora* (lichenized Ascomycota) new to Poland. *Acta Soc. Bot. Poloniae* 84: 303–307.
- Czarnota, P., Kison, H.U. & Seelemann, A. 2014. Remarkable records of lichens and lichenicolous fungi from the Harz National Park (Lower Saxony and Saxony-Anhalt, Germany). *Herzogia* 27: 67–82.
- Czarnota, P. & Kukwa, M. 2001. Lichens of the genera *Lepraria* and *Leproloma* from the Gorce Mts (Western Carpathians, Poland) and note on lichenicolous fungus *Paranectria oropensis* found on *Leproloma membranaceum*. *Pol. Bot. J.*: 199–206.
- Czarnota, P. & Kukwa, M. 2009. Contribution to the knowledge of some poorly known lichens in Poland. III. *Trapelia corticola* and the genus *Vezeadaea*. *Folia Crypt. Estonica* 46: 25–31.
- Dietrich, M. & Scheidegger, C. 1996. The importance of sorediate crustose lichens in the epiphytic lichen flora of the Swiss Plateau and the Pre-Alps. *Lichenologist* 28: 245–256.
- Dymytrova, L. 2011. Notes on the genus *Scoliciosporum* (Lecanorales, Ascomycota) in Ukraine. *Pol. Bot. J.* 56: 61–75.
- Dymytrova, L., Breuss, O. & Kondratyuk, S.Y. 2011. *Agonimia borysthenica*, a new lichen species (Verrucariales) from Ukraine. *Osterr. Z. Pilz.* 20: 25–28.

Dymytrova, L., Nadyeina, O., Hobi, M.L. & Scheidegger, C. 2014. Topographic and forest-stand variables determining epiphytic lichen diversity in the primeval beech forest in the Ukrainian Carpathians. *Biodivers. Conserv.* 23: 1367–1394.

Dymytrova, L., Nadyeina, O., Naumovych, A., Keller, C. & Scheidegger, C. 2013. Primeval beech forests of Ukrainian Carpathians are sanctuaries for rare and endangered epiphytic lichens. *Herzogia* 26: 73–89.

Ellegård Larsen, H.M., Hansen, E.S., Nord-Larsen, T. & Rasmussen, H.N. 2020. Deciduous trees as lichen phorophytes: biodiversity and colonization patterns under common garden conditions. *Lichenologist* 52: 221–232.

Erdmann, G., Otte, V., Langel, R., Scheu, S. & Maraun, M. 2007. The trophic structure of bark-living oribatid mite communities analysed with stable isotopes (^{15}N , ^{13}C) indicates strong niche differentiation. *Exp. Appl. Acarol.* 41: 1–10.

Ertz, D., Diederich, P., Brand, A.M., van den Boom, P. & Sérusiaux, E. 2008. New or interesting lichens and lichenicolous fungi from Belgium, Luxembourg and northern France. *XI. Bull. Soc. Natur. Luxembourg.* 109: 35–51.

Fałtynowicz, W. 2003. The lichens, lichenicolous and allied fungi of Poland. An annotated checklist. In: Mirek, Z. (Ed.). *Biodiversity of Poland* 6: 1–435. W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków.

Fernández-Marín, B., Becerril, J.M. & García-Plazaola, J.I. 2010. Unravelling the roles of desiccation-induced xanthophyll cycle activity in darkness: a case study in *Lobaria pulmonaria*. *Planta* 231: 1335–1342.

Flakus, A. & Kukwa, M. 2009. Additions to the biota of lichenized fungi of Poland. *Acta Mycol.* 44: 249–257.

Frati, L., Brunialti, G., Landi, S., Filigheddu, R. & Bagella, S. 2021. Exploring the biodiversity of key groups in coppice forests (Central Italy): the relationship among vascular plants, epiphytic lichens, and wood-decaying fungi. *Plant Biosyst.* 156(4): 835–846.

Friedel, A., Oheimb, G.V., Dengler, J. & Härdtle, W. 2006. Species diversity and species composition of epiphytic bryophytes and lichens - a comparison of managed and unmanaged beech forests in NE Germany. *Feddes Repert.* 177: 172–185.

Fritz, Ö., Brunet, J. & Caldiz, M. 2009. Interacting effects of tree characteristics on the occurrence of rare epiphytes in a Swedish boreal forest. *Bryologist* 112: 488–505.

Fritz, Ö., Gustafsson, L. & Larsson, K. 2008. Does forest continuity matter in conservation? – A study of epiphytic lichens and bryophytes in beech forests of southern Sweden. *Biol. Conserv.* 141: 655–668.

Fritz, Ö., Niklasson, M. & Churski, M. 2009. Tree age is a key factor for the conservation of epiphytic lichens and bryophytes in beech forests. *Appl. Veg. Sci.* 12: 93–106.

- Fuertes, E., Burgaz, A.R. & Escudero, A. 1996. Pre-climax epiphyte communities of bryophytes and lichens in Mediterranean forests from the Central Plateau (Spain). *Vegetatio* 123: 139–151.
- Gómez, A. & Hladun, N.L. 1981. Datos para la flora líquénica de Catalunya: epífitos de *Fagus sylvatica* L. *Butll. Inst. Catalana Hist. Nat.* 46: 83–94.
- Groner, U. & LaGreca, S. 1997. The 'Mediterranean' *Ramalina panizzei* north of the Alps: morphological, chemical and rDNA sequence data. *Lichenologist* 29: 441–454.
- Guzow-Krzemińska, B., Halda, J.P. & Czarnota, P. 2012. A new *Agonimia* from Europe with a flabelliform thallus. *Lichenologist* 44: 55–66.
- Hafellner, J. & Kashta, L. 2003. Miscellaneous records of lichens and lichenicolous fungi from Albania. *Herzogia* 16: 135–142.
- Hafellner, J. & Türk, R. 2016. Die lichenisierten Pilze Österreichs – eine neue Checkliste der bisher nachgewiesenen Taxa mit Angaben zu Verbreitung und Substratökologie [The lichenized fungi of Austria – a new checklist of the taxa so far recorded, with data to distribution and substrate ecology]. *Stapfia* 104: 1–216.
- Hallingbäck, T. 1989. Occurrence and ecology of the lichen *Lobaria scrobiculata* in southern Sweden. *Lichenologist* 21: 331–341.
- Heinrichs, S., Ammer, Ch., Mund, M., Boch, S., Budde, S., Fischer, M., Müller, J., Schöning, I., Schulze, E.D., Schmidt, W., Weckesser, M. & Schall, P. 2019. Landscape-scale mixtures of three species are more effective than stand-scale mixtures for biodiversity of vascular plants, bryophytes and lichens. *Forests* 10: 73.
- Hofmeister, J., Hošek, J., Malíček, J., Palice, Z., Syrovátková, L., Steinová, J. & Černajová, I. 2016. Large beech (*Fagus sylvatica*) trees as 'lifeboats' for lichen diversity in central European forests. *Biodivers. Conserv.* 25: 1073–1090.
- Hurtado, P., Prieto, M., de Bello, F., Aragón, G., López-Angulo, J., Giordani, P., Díaz-Peña, E., Vicente, R., Merinero, S., Košuthová, A., Benesperi, R., Bianchi, E., Mayrhofer, H., Nascimbene, J., Grube, M., Wedin, M., Westberg, M. & Martínez, I. 2020. Contrasting environmental drivers determine biodiversity patterns in epiphytic lichen communities along a European gradient. *Microorganisms* 8:1913.
- Hurtado, P., Prieto, M., Aragón, G., Escudero, A. & Martínez, I. 2019. Critical predictors of functional, phylogenetic, and taxonomic diversity are geographically structured in lichen epiphytic communities. *J. Ecol.* 107: 2303–2316.
- Hurtado, P., Prieto, M., Martínez-Vilalta, J., Giordani, P., Aragón, G., López-Angulo, J., Košuthová, A., Merinero, S., Díaz-Peña, E.M., Rosas, T., Benesperi, R., Bianchi, E., Grube, M., Mayrhofer, H., Nascimbene, J., Wedin, M., Westberg, M. & Martínez, I. 2020. Disentangling functional trait variation and covariation in epiphytic lichens along a continent-wide latitudinal gradient. *P. Roy Soc. B* 287: 20192862.

- Jabłońska, A., Oset, M. & Kukwa, M. 2009. The lichen family Parmeliaceae in Poland. I. The genus *Parmotrema*. *Acta Mycol.* 44: 211–222.
- Jönsson, A.M. 1998. Bark lesions on beech (*Fagus sylvatica*) and their relation to epiphytes and site variables in Scania, South Sweden. *Scandinavian J. Forest Res.* 13: 297–305.
- Kaufmann, S., Hauck, M. & Leuschner, C. 2018. Effects of natural forest dynamics on vascular plant, bryophyte, and lichen diversity in primeval *Fagus sylvatica* forests and comparison with production forests. *J. Ecol.* 106: 2421–2434.
- Kaufmann, S., Weinrich, T., Hauck, M. & Leuschner, C. 2019. Vertical variation in epiphytic cryptogam species richness and composition in a primeval *Fagus sylvatica* forest. *J. Veg. Sci.* 30: 881–892.
- Khodosovtsev, O., Dymytrova, L., Nadyeina, A., Naumovych, G., Khodosovtseva, Y. & Scheidegger, C. 2013. A contribution to beech forest-associated epiphytic lichen-forming and lichenicolous fungi in Crimean Mts (Ukraine). *Fl. Med.* 23: 57–68.
- Kossowska, M., Fałtynowicz, W., Dimos-Zych, M., Fałtynowicz, H., Patejuk, K., Piegdoń, A., Buksakowska, M. & Jarema, A. 2018. Additions to the lichen biota of the Sudety Mountains. I. Records from the Karkonosze Mountains. *Acta Mycol* 53: 11–13.
- Kowalewska, A. & Kukwa, M. 2003. Addition to the Polish lichen flora. *Graphis Scripta* 14: 11–17.
- Kowalewska, A., Kukwa, M., Ostrowska, I., Jabłońska, A., Oset, M. & Szok, J. 2008. The lichens of the *Cladonia pyxidata-chlorophaea* group and allied species in Poland. *Herzogia* 21: 61–78.
- Kubiak, D. 2005. Lichens and lichenicolous fungi of Olsztyn town (NE Poland). *Acta Mycol.* 40: 293–332.
- Kukwa, M. 2009. The lichen genus *Ochrolechia* in Poland III with a key and notes on some taxa. *Herzogia* 22: 43–66.
- Kukwa, M. & Diederich, P. 2007. New records of leprarioid lichens from Luxembourg and France, with the first report of fertile *Lecanora rouxii*. *Bull. Soc. Natur. Luxemb.* 108: 15–19.
- Kukwa, M., Łubek, A., Szymczyk, R. & Zalewska, A. 2012. Seven lichen species new to Poland. *Mycotaxon* 120: 105–118.
- Kukwa, M. & Śliwa, L. 2005. The genus *Lepraria* (lichenized Ascomycota) in the Beskid Sdecki Mts (West Carpathians, S Poland). *Pol. Bot. J.* 50: 163–184.
- Labak, I., Ozimec, S., Dumbović, V. & Topić, J. 2011. Contribution to the knowledge of lichens of Papuk Nature Park (Slavonia, eastern Croatia). *Nat. Croat.* 20: 35–52.

- Lackovičoca, A. & Guttová, A. (2005) Genus *Dimerella* (Coenogoniaceae, lichenized Ascomycota) in Slovakia. *Acta Bot. Croat.* 64: 289–301.
- Liška, J., Dětinský, R. & Palice, Z. 1996. Importance of the Šumava Mts. for the biodiversity of lichens in the Czech Republic. *Silva Gabreta* 1: 71–81.
- Llop, E. 2012-2013. La diversitat líquènica de les fagedes olositàniques, i un estudi comparatiu de la diversitat líquènica epifítica dels alzinars i les fagedes de la Garrotxa. *Butll. Inst. Catalana Hist. Nat.* 77: 61–74.
- Loppi, S., Bonini, I. & de Dominicis, V. 1999. Epiphytic lichens and bryophytes of forest ecosystems in Tuscany. *Cryptogamie Mycol.* 20: 127–135.
- Machacova, K., Maier, M., Svobodova, K., Lang, F. & Urban, O. 2017. Cryptogamic stem covers may contribute to nitrous oxide consumption by mature beech trees. *Sci. Rep.* 7: 13243.
- Malíček, J., Berger, F., Palice, Z. & Vondrák, J. 2017. Corticolous sorediate *Lecanora* species (Lecanoraceae, Ascomycota) containing atranorin in Europe. *Lichenologist* 49: 431–455.
- Malíček, J., Bouda, F., Liška, J., Palice, Z. & Peksa, O. 2015. Contribution to the lichen biota of the Romanian Carpathians. *Herzogia* 28: 713–735.
- Malíček, J. & Mayrhofer, H. 2017. Additions to the lichen diversity of Macedonia (FYROM). *Herzogia* 30: 431–444.
- Malíček, J. & Palice, Z. 2013. Lichens of the virgin forest Reserve Žofínský Prales (Czech Republic) and surrounding woodlands. *Herzogia* 26: 253–292.
- Malíček, J., Palice, Z., Acton, A., Berger, F., Bouda, F., Sanderson, N. & Vondrák, J. 2018. Uholka primeval forest in the Ukrainian Carpathians – a keynote area for diversity of forest lichens in Europe. *Herzogia* 31: 140–171.
- Malíček, J., Palice, Z. & Vondrák, J. 2015. New lichen records and rediscoveries from the Czech Republic and Slovakia. *Herzogia* 27: 257–284.
- Marti, J. & Aebli, A. 2019. Die Verbreitung von *Usnea florida* im Kanton Glarus. *Meylania* 64: 22–26.
- Martínez, I., Flores, T., Otálora, M.A., Belinchón, R., Prieto, M., Aragón, G. & Escudero, A. 2012. Multiple-scale environmental modulation of lichen reproduction. *Fungal Biol.* 116: 1192–1201.
- Massara, A.C., Bates, J.W. & Bell, J.N.B. 2009. Exploring causes of the decline of the lichen *Lecanora conizaeoides* in Britain: effects of experimental N and S applications. *Lichenologist* 41: 673–681.
- Mayrhofer, H., Bilovitz, P.O. & Rohrer, A. 2018. Lichenized and lichenicolous fungi from Croatia kept in the herbarium GZU. *Fritschiana* 89: 1–35.

- Mayrhofer, H. & Coppins, B.J. 2013. Remarkable records of lichenized fungi from Slovenia. *Herzogia* 26: 201–206.
- Mayrhofer, H., Czeh, D., Kobald, E.M. & Bilovitz, P.O. 2016. Catalogue of the lichenized and lichenicolous fungi of Kosovo. *Herzogia* 29: 529–554.
- Mayrhofer, H., Stešević, D., Brudermann, A., Fötschl, B.R. & Bilovitz, P.O. 2017. New or otherwise interesting lichenized and lichenicolous fungi from Montenegro II. *Fritschiana* 86: 1–30.
- Mazziotta, A., Vizentin-Bugoni, J., Tottrup, A.P., Bruun, H.H., Fritz, Ö. & Heilmann-Clausen, J. 2017. Interaction type and intimacy structure networks between forest-dwelling organisms and their host trees. *Basic Appl. Ecol.* 24: 86–97.
- Mitchell, R.J., Hewison, R.L., Beaton, J. & Douglass, J.R. 2021. Identifying substitute host tree species for epiphytes: The relative importance of tree size and species, bark and site characteristics. *Appl. Veg. Sci.* 24: e12569.
- Mrak, T., Mayrhofer, H. & Batić, F. 2004. Contributions to the lichen flora of Slovenia XI. Lichens from the vicinity of Lake Bohinj (Julian Alps). *Herzogia* 17: 107–127.
- Nadyeina, O., Dymytrova, L., Naumovych, A., Postoyalkin, S., Werth, S., Cheenacharoen, S. & Scheidegger, C. 2014. Microclimatic differentiation of gene pools in the *Lobaria pulmonaria* symbiosis in a primeval forest landscape. *Mol. Ecol.* 23: 5164–5178.
- Nadyeina, O., Grube, M. & Mayrhofer, H. 2010. A contribution to the taxonomy of the genus *Rinodina* (Physciaceae, lichenized Ascomycotina) using combined ITS and mtSSU rDNA data. *Lichenologist* 42: 521–531.
- Nascimbene, J., Marini, L. & Nimis, P.L. 2009. Influence of tree species on epiphytic macrolichens in temperate mixed forests of northern Italy. *Can. J. Forest Res.* 39: 785–791.
- Nascimbene, J., Di Cecco, V., Di Martino, L., Frascaroli, F., Giordani, P., Lelli, Ch., Vallese, Ch., Zannini, P. & Chiarucci, A. 2019. Epiphytic lichens of the sacred natural site "Bosco di Sant'Antonio" (Majella National Park-Abruzzo). *Ital. Bot.* 7: 149–156.
- Nimis, P.L. 1984. Phytosociology, ecology and phytogeography of epiphytic lichen vegetation in the Calamone Lake area. *Studia Geobot.* 4: 109–127.
- Nimis, P.L. 1985. Contributi alle conoscenze floristiche sui licheni d'Italia I. Florula lichenica del M. Ventasso (Appennino Reggiano). *Webbia* 39: 141–161.
- Obermayer, W. & Mayrhofer, H. 2007. Hunting for *Cetrelia chicitae* (lichenized Ascomycetes) in the eastern European Alps. *Phyton* 47: 231–290.

- Öckinger, E., Niklasson, M. & Nilsson, S.G. 2005. Is local distribution of the epiphytic lichen *Lobaria pulmonaria* limited by dispersal capacity or habitat quality? *Biodivers. Conserv.* 14: 759–773.
- Oran, S. 2011. Epiphytic lichen diversity on *Fagus orientalis* Lipsky and *Fagus sylvatica* L. in the Marmara region (Turkey). *Biol. Divers. Conserv.* 4: 134–143.
- Otte, V. 2005. Noteworthy lichen records for Bulgaria. *Abh. Ber. Naturkundemus. Görlitz* 77: 77–86.
- Ozimec, S., Bošković, I., Florijančić, T., Jelkić, D., Opačak, A., Puškadija, Z. & Labak, I. 2010. The lichen flora of Risnjak National Park (Croatia). *Acta Bot. Croat.* 69: 19–29.
- Packham, J.R., Thomas, P.A., Atkinson, M.D. & Degen, T. 2012. Biological flora of the British Isles: *Fagus sylvatica*. *J. Ecol.* 100: 1557–1608.
- Paoli, L., Guttová, A., Grassi, A., Lackovičová, A., Senko, D. & Loppi, S. 2014. Biological effects of airborne pollutants released during cement production assessed with lichens (SW Slovakia). *Ecol. Ind.* 40: 127–135.
- Pilegaard, K., Rasmussen, L. & Gydesen, H. 1979. Atmospheric background deposition of heavy metals in Denmark monitored by epiphytic cryptogams. *J. Appl. Ecol.* 16: 843–853.
- Pirintsos, S., Diamantopoulos, J. & Stamou, G. 1995. Analysis of the distribution of epiphytic lichens within homogeneous *Fagus sylvatica* stands along an altitudinal gradient (Mount Olympos, Greece). *Vegetatio* 116: 33–40.
- Pirintsos, S.A., Diamantopoulos, J. & Stamou, G.P. 1996. Hierarchical analysis of the relationship between spatial distribution and abundance of epiphytic lichens (Mt. Olympos-Greece). *Vegetatio* 122: 95–106.
- Pišút, I., Lackovičová, A., Guttová, A. & Palice, Z. 2007. New lichen records from Bukovské vrchy Mts (NE Slovakia). *Acta Mycol.* 42: 267–280.
- Prügger, J., Surina, B. & Mayrhofer, H. 2001. The lichens of the Ždrcle Forest Reserve. *Zb. Gozd. Lesarstva* 63: 7–25.
- Puntillo, D. 1993. Contributi alle conoscenze floristiche sui Licheni d'Italia. 6. Florula lichenica della Valle del Fiume Argentino (Calabria Italia). *Webbia* 47: 163–186.
- Puntillo, D. & Vězda, A. 1991. A new species of *Gyalideopsis* (Lichens) from Calabria (South Italy). *Webbia* 46: 159–161.
- Purvis, O.W., Tittley, I., Chimonides, P.D.J., Bamber, R., Hayes, P.A., James, P.W., Rumsey, F.J. & Read, H. 2010. Long-term biomonitoring of lichen and bryophyte biodiversity at Burnham Beeches SAC and global environmental change. *Syst. Biodivers.* 8: 193–208.

- Ravera, S., Genovesi, V., Falasca, A., Marchetti, M. & Chirici, G. 2010. Lichen diversity of old-growth forests in Molise (Central-Southern Italy). *L'Italia Forestale e Montana/Ital. J. Forest Mount. Environ.* 65: 505–517.
- Ravera, S., Genovesi, V. & Massari, G. 2006. Phytoclimatic characterization of lichen habitats in central Italy. *Nova Hedwigia* 82: 143–165.
- Rohrer, A., Bilovitz, P.O. & Mayrhofer, H. 2012. Lichenized fungi from the Jakupica Mountain range (Macedonia, FYROM). *Herzogia* 25: 167–175.
- Rose, F. & James, P. 1974. Regional studies on the British lichen flora I. The corticolous and lignicolous species of the New Forest, Hampshire. *Lichenologist* 6: 1–72.
- Roux, C. et al. 2017. Catalogue des lichens et champignons lichénicoles de France métropolitaine. Édité. Association française de lichénologie (A. F. L.), Fontainebleau.
- Ruete, A., Fritz, Ö. & Snäll, T. 2014. A model for non-equilibrium metapopulation dynamics utilizing data on species occupancy, patch ages and landscape history. *J. Ecol.* 102: 678–689.
- Schiefelbein, U. 2013. Additions to the lichenized and lichenicolous fungi of Mecklenburg-Western Pomerania (Germany). *Herzogia* 26: 65–72.
- Schiefelbein, U., Czarnota, P., Thüs, H. & Kukwa, M. 2012. The lichen biota of the Drawieński National Park (NW Poland, Western Pomerania). *Folia Crypt. Eston.* 49: 59–71.
- Schiefelbein, U. & Thell, A. 2018. Current state of *Lobaria pulmonaria* in southernmost Sweden. *Graphis Scripta* 30: 94–104.
- Sérusiaux, E., Coppins, B., Diederich, P. & Scheidegger, C. 2001. *Fellhanera gyrophorica*, a new European species with conspicuous pycnidia. *Lichenologist* 33: 285–289.
- Slavíková-Bayerová, Š. & Orange, A. 2006. Three new species of *Lepraria* (Ascomycota, Stereocaulaceae) containing fatty acids and atranorin. *Lichenologist* 38: 503–513.
- Špičková, J., Navrátil, T., Rohovec, J., Mihaljevič, M., Kubínová, P., Minařík, L. & Skřivan, P. 2010. The characteristics of rare earth elements in bulk precipitation, throughfall, foliage and lichens in the Lesní potok catchment and its vicinity, Czech Republic. *Geochem. Explor. Environ. Anal.* 10: 383–390.
- Spier, L., van Dort, K. & Fritz, Ö. 2008. A contribution to the lichen mycota of old beech forests in Bulgaria. *Mycol. Balc.* 5: 141–146.
- Strasser, E.A., Hafellner, J., Stešević, D., Geci, F. & Mayrhofer, H. 2015. Lichenized and lichenicolous fungi from the Albanian Alps (Kosovo, Montenegro). *Herzogia* 28: 520–544.

- Svoboda, D., Bouda, F., Malíček, J. & Hafellner, J. 2012. A contribution to the knowledge of lichenized and lichenicolous fungi in Albania. *Herzogia* 25: 149–165.
- Szczepańska, K. & Staniaszek-Kik, M. 2012. New lichenized fungi of the Polish Karkonosze Mountains. *Pol. Bot. J.* 57: 279–283.
- Szymczyk, R. & Zalewska, A. 2008. Lichen biota of the Grabianka River Valley in the Elbląg Upland (Wysoczyzna Elbląska). *Pol. J. Nat. Sci.* 23: 398–414.
- Szymczyk, R., Zalewska, A., Szydłowska, J. & Kukwa, M. 2015. The lichen family Parmeliaceae in Poland. IV. The genus *Punctelia*. *Herzogia* 28: 556–566.
- Thor, G. 1988. *Caloplaca lucifuga*: a new lichen species from Europe. *Lichenologist* 20: 175–178.
- Tretiach, M. 2014. *Porina pseudohibernica* sp. nov., an isidiate, epiphytic lichen from central and south-eastern Europe. *Lichenologist* 46: 617–625.
- van den Boom, P., Brand, M., Diederich, P., Aptroot, A. & Sérusiaux, E. 1994. Report of a lichenological field meeting in Luxembourg. *Bull. Soc. Natur. Luxembourg.* 95: 145–176.
- van den Boom, P. & van den Boom, B. 2006. An inventory of lichens and lichenicolous fungi in northern Belgium (area of Stramprooierbroek). *Bull. Soc. Natur. Luxembourg.* 106: 39–52.
- van der Pluijm, A. 2015. *Phaeographis smithii* terug in Nederland in wilgenbossen in de Biesbosch. *Buxbaumiella* 103: 7–13.
- Vicol, I. 2013. Chorology of the *Physcia* (Schreb.) Michaux (1803) genus in Romania. *Muzeul Olteniei Craiova. Oltenia* 29: 268–282.
- Vicol, I. 2013. Distribution of *Flavoparmelia caperata* (L.) Hale in Romania. *Analele Științifice ale Universității “Al. I. Cuza” Iași s. II a. Biologie vegetală* 59: 65–73.
- Vicol, I. 2016. Effect of old-growth forest attributes on lichen species abundances: A study performed within Ceahlău National Park (Romania). *Cryptogamie Mycol.* 36: 399–407.
- Vondrák, J. & Liška, J. 2013. Lichens and lichenicolous fungi from the Retezat Mts and overlooked records for the checklist of Romanian lichens. *Herzogia* 26: 293–305.
- Vondrák, J., Malíček, J., Šoun, J. & Pouska, V. 2015. Epiphytic lichens of Stuzica (E Slovakia) in the context of Central European old-growth forests. *Herzogia* 28: 108–130.
- Wieczorek, A. 2018. The lichen genus *Opegrapha* s.l. in Poland: morphological variability, ecology, and distribution. *Monog. Bot.* 107: 1–162.
- Zduńczyk, A. & Kukwa, M. 2014. A revision of sorediate crustose lichens containing usnic acid and chlorinated xanthenes in Poland. *Herzogia* 27: 13–40.