

HEEGAARD SPLITTINGS OF THE PAIR OF THE SOLID TORUS AND THE CORE LOOP

Chuichiro HAYASHI* and Koya SHIMOKAWA†

Abstract

We show that any Heegaard splitting of the pair of the solid torus ($\cong D^2 \times S^1$) and its core loop (an interior point $\times S^1$) is standard, using the notion of Heegaard splittings of pairs of 3-manifolds and properly imbedded graphs, which is defined in this paper.

1 Introduction

The bridge presentation of a link in the 3-sphere S^3 is introduced by H. Schubert[S] and generalized by H. Doll[D] for a link in a closed orientable 3-manifold. In [HS1], [HS2], [HS3] we generalize it for a 1-submanifold properly imbedded in a compact connected orientable 3-manifold possibly with boundary. Here we recall it.

Let $I = [0, 1]$ an interval, F a disjoint union of closed orientable surfaces. A *compression body* C is a connected orientable 3-manifold obtained from a 3-manifold W homeomorphic to a ball B or $F \times I$ by attaching 0 or several 1-handles on ∂B or $F \times \{1\}$. Set $\partial_- C = \emptyset$ when $W \cong B$, and $\partial_- C = F \times \{0\}$ when $W \cong F \times I$. Let $\partial_+ C = \partial C - \partial_- C$. In usual definitions $\partial_- C$ has no 2-sphere component, but in this paper $\partial_- C$ may have 2-sphere components. A compression body C is called a *handlebody* if $W \cong B$ and $\partial_- C = \emptyset$.

A set of arcs $\{t_1, \dots, t_n\}$ properly imbedded in a compression body C is *trivial* if there is a homeomorphism $C \cong Y \cup V$ (where Y is a ball or

*The first author was partially supported by Grant-in-Aid for Scientific Research (No. 10740039), Ministry of Education, Science and Culture.

†The second author was partially supported by Fellowship of the Japan Society for the Promotion of Science for Japanese Junior Scientists.

homeomorphic to $\partial_- C \times I$ and V is a disjoint union of 1-handles) such that each arc t_i satisfies one of the following conditions.

- (1) t_i is *vertical*, i.e., $t_i = (\text{a point}) \times I \subset \partial_- C \times I$ and $t_i \cap V = \emptyset$.
- (2) t_i is ∂_+ -*parallel*, i.e., there is an imbedded disc $D \subset C$ such that $(\cup_j t_j) \cap D = t_i \subset \partial D$ and $D \cap \partial C = \text{cl}(\partial D - t_i) \subset \partial_+ C$.

We call such a disc D as in the condition (2) a *cancelling disc* of t_i . A standard cut and paste argument allows us to take the cancelling discs to be disjoint.

It is well known that every compact connected orientable 3-manifold M has a Heegaard splitting H , i.e., $M = C_1 \cup_H C_2$, where C_1 and C_2 are compression bodies and $H = \partial_+ C_1 = \partial_+ C_2$. Let T be a properly imbedded 1-manifold in M . The Heegaard splitting H of M is a *Heegaard splitting* of (M, T) if H is transverse to T and $T_i = T \cap C_i$ is trivial in C_i for $i = 1$ and 2 . Every pair of a compact connected orientable 3-manifold and a properly imbedded 1-manifold has a Heegaard splitting as shown in [Lemma 2.1, HS1]. See also Lemma 2.1 in this paper.

A Heegaard splitting $(M, T) = (C_1, T_1) \cup_H (C_2, T_2)$ is said to be *weakly T -reducible* if there is a disc D_i properly imbedded in C_i for $i = 1$ and 2 such that $D_i \cap T = \emptyset$, ∂D_i is an essential loop in $H - T$ and $\partial D_1 \cap \partial D_2 = \emptyset$ in H . Otherwise the splitting H is *strongly T -irreducible*. The splitting H is *T -reducible* if we can take discs as above so that $\partial D_1 = \partial D_2$ instead of the condition $\partial D_1 \cap \partial D_2 = \emptyset$. Otherwise H is *T -irreducible*. If H is T -reducible, then we can see that H is weakly T -reducible by slightly isotoping D_2 near ∂D_2 to be disjoint from ∂D_1 . A Heegaard splitting $(M, T) = (C_1, T_1) \cup_H (C_2, T_2)$ is said to be *cancellable* if there is a cancelling disc D_i of an arc component t_i of T_i for $i = 1$ and 2 such that $\emptyset \neq (\partial D_1 \cap \partial D_2) \subset (T \cap H)$. We call these discs a *cancelling pair* of discs. When H is cancellable and $\partial D_1 \cap \partial D_2$ is a single point, isotoping t_i along the disc D_i for $i = 1$ and 2 , we can obtain another Heegaard splitting H' of (M, T) such that $|T \cap H'| = |T \cap H| - 2$.

A Heegaard splitting $(M, T) = (C_1, T_1) \cup_H (C_2, T_2)$ is said to be *stabilized* if there is a properly imbedded disc D_i disjoint from T_i in C_i for $i = 1$ and 2 such that ∂D_1 and ∂D_2 intersect transversely at a single point in H .

Here we give a characterization of Heegaard splittings of a pair of solid torus and its core loop. The following theorem is used in the proof

of [Theorem 1.2, HS1] to extend a result by A.J. Casson and C.McA. Gordon ([Theorem 3.1, CG]) on irreducible and weakly reducible Heegaard splittings and closed incompressible surfaces. There we use this theorem to show that the obtained closed incompressible surface does not bound a solid torus intersecting the 1-submanifold in its core loop.

Theorem 1.1 *Let V be a solid torus, T_0 a core loop of V , and H a Heegaard splitting of (V, T_0) . Then H is either stabilized or cancellable. Moreover, when $|H \cap T_0| = 2$ and $\text{genus}(H) \geq 2$, H is stabilized.*

Heegaard splittings of handlebodies can be easily classified. See [Lemma 2.7, ST].

2 Heegaard splittings of imbedded graphs in 3-manifolds

H. Goda defined bridge presentations for theta curves in [G]. Here we generalize it for pairs of compact connected orientable 3-manifolds and ‘properly imbedded’ graphs.

Let M be a compact 3-manifold possibly with non-empty boundary, and T a (possibly disconnected) graph imbedded in M . The graph T is said to be *properly imbedded* in M if $T \cap \partial M$ consists of all the valency one vertices of T . Let ∂T denote the union of these vertices.

Let C be a compression body and T a graph properly imbedded in C . The graph T is said to be *trivial* in C if there is a homeomorphism $C \cong Y \cup V$ as below, where Y is a ball or homeomorphic to $\partial_- C \times I$ and V is a disjoint union of 1-handles.

- (1) Each component t of $T \cap C$ is an arc or a graph with a single vertex besides ∂t ,
- (2) every arc component of $T \cap C$ is either vertical or ∂_+ -parallel (as in section 1),
- (3) for every component t containing a vertex of T besides ∂t there is a properly imbedded disc D such that $D \cap T = t$ and $D \cap \partial C = \partial D \subset \partial C_+$.

We can take the disc D in (3) so that it is inessential in C , by replacing a disc component D' of $D - t$ with a parallel copy of the disc

$D - D'$ slightly isotoped off of $D - \text{cl}(D')$. By performing this operation on every disc of (3), we can take the discs of (3) to be not nested.

Let H be a Heegaard splitting surface of a compact connected orientable 3-manifold M , and C_1, C_2 the compression bodies obtained by cutting M along H . Let T be a graph properly imbedded in M . We say H is a *Heegaard splitting* of the pair (M, T) if the vertices of T are disjoint from H , the graph T is transverse to H and the graph $T \cap C_i$ is trivial in C_i for $i = 1$ and 2 .

Let C be a compression body, and T a trivial graph properly imbedded in C . A *spine* of (C, T) is a (possibly disconnected when $\partial_- C \neq \emptyset$) 1-dimensional complex Q (possibly non-properly) imbedded in C such that

- (1) $\partial_+ C \cap Q = \emptyset$,
- (2) all the vertical arcs of T are disjoint from Q ,
- (3) every component t of T intersects Q at precisely one vertex of Q if t is not a vertical arc,
- (4) for every component t of T with a vertex p besides ∂t , the 1-complex Q intersects t at the vertex p and
- (5) C collapses to $\partial_- C \cup Q \cup T$.

When $T = \emptyset$, we call it a *spine* of C .

Lemma 2.1. *Let M be a compact connected orientable 3-manifold possibly with boundary, and H be a Heegaard splitting of M . Let T be a properly imbedded graph in M . Then we can move T by an ambient isotopy of M so that H gives a Heegaard splitting of (M, T) .*

Proof. The manifold M is decomposed into two compression bodies C_1 and C_2 , where $H = \partial_+ C_1 = \partial_+ C_2$. Let X_i be a spine of C_i for $i = 1$ and 2 . We can isotope T to be disjoint from $N(X_1 \cup X_2)$. Then $M - \text{int}(N(X_1) \cup N(X_2))$ is homeomorphic to $H \times I$ where $I = [0, 1]$, $H = H \times \{1/2\}$ and $H \times \{0\} \subset C_1$. Let π be the projection $H \times I \rightarrow H \times \{1/2\}$. We can take this product structure so that the singular set of $\pi(T)$ consists of double points away from the vertices of T . Let v be a vertex of T . Let S be the set consisting of the vertices of T

and the singular points on T , the over ones and the under ones, with respect to the projection map π . We can isotope T in M so that short arcs containing the over singular points are contained in C_2 and so that short arcs containing the under singular points are contained in C_1 . (Note that the points of ∂T are also vertices.) We take regular points of T , one between every pair of adjacent points of S . We take regular points of T , one more between adjacent points of S if both points are contained in C_1 or in C_2 . In addition we take two regular points of T on every circle component of T without singular points. Then we can isotope T so that $T \cap H$ consists of the above regular points and $T \cap C_i$ is trivial in C_i for $i = 1$ and 2 . ■

In general, let X be a compact 3-manifold, and T a graph properly imbedded in X . Let F be a union of compact surfaces in X . Suppose that every surface (connected component) F_0 of F satisfies (1) or (2) below. (1) F_0 is properly imbedded in X transversely to T . (2) F_0 is imbedded in X , $\partial F_0 \subset (\partial X \cup T)$, $(F_0 \cap \partial X) \subset \partial F_0$, and $F_0 \cap T$ is a union of edges of T . In addition, suppose that distinct surfaces of F may intersect each other in a union of edges of T . Then we say that F is *T -compressible* if there is a disc D such that $D \cap T = \emptyset$, $D \cap F = \partial D$ is essential in $F - T$, that is, ∂D does not bound a disc in $F - T$. We call such a disc D a *T -compressing disc* of F . If F is not T -compressible, then it is *T -incompressible*. We say that F is *T - ∂ -compressible* if there is a disc D such that $D \cap T = \emptyset$, $D \cap (F \cup \partial X) = \partial D$ and $\partial D \cap F = \alpha$ is an essential arc in $F - T$, that is, α does not cobound a disc with a subarc of $\partial F - T$ on $F - T$. We call such a disc D a *T - ∂ -compressing disc* of F . If F is not T - ∂ -compressible, then it is *T - ∂ -incompressible*.

Lemma 2.2. *Let B be a ball, and T a connected graph with a single vertex v besides ∂T . Suppose that T is properly imbedded in B and trivial in B . Let F be a union of compact surfaces in B . Suppose for every surface F_0 of F that F_0 is imbedded in B , $\partial F_0 \subset (\partial B \cup T)$, $(F_0 \cap \partial B) \subset \partial F_0$ and $F_0 \cap T$ is a union of edges of T . In addition, suppose that distinct surfaces of F may intersect each other in a union of edges of T . If F is T -incompressible and T - ∂ -incompressible, then F is a union of*

- (1) *spheres disjoint from T ,*
- (2) *properly imbedded discs disjoint from T ,*

- (3) discs such that the boundary loop ℓ of each of them intersects ∂B in an arc α and such that the arc $\text{cl}(\ell - \alpha)$ is in T ,
- (4) discs which are unions of discs of the above type (3) (see Figure 2.1) and
- (5) properly imbedded discs each of which contains a single edge of T with one of the endpoints v in its interior.




Figure 2.1

Proof. Let $N(T)$ be a small regular neighbourhood of T in B . We can take $N(T)$ so that it is a union of a ball neighbourhood $N(v)$ of the vertex v and very small tubular neighbourhoods of the edges of T , and each tubular neighbourhood intersects $\partial N(v)$ in a very small disc.

Claim 2.3. Let ρ be an arc which is a union of an arc ρ_0 in the punctured sphere $\partial N(T) \cap \partial N(v)$, and two essential arcs in the annuli $\text{cl}(\partial N(T) - (\partial N(v) \cup \partial B))$. Then ρ and an arc in ∂B together form a loop which bounds a disc properly imbedded in $\text{cl}(B - N(T))$.

Proof of Claim 2.3. We can coordinate $B = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 \leq 1\}$ so that $v = (0, 0, 0)$ and the edges of T are straight line segments connecting v and ∂B . Set $B_\epsilon = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 \leq \epsilon^2\}$ for a small positive real number ϵ . We can take the coordinates so that $N(T)$ is a union of B_ϵ and line segments connecting v and ∂B . Let D denote the disc which is the union of straight line segments connecting v and ∂B and intersecting the arc ρ_0 . Then $D \cap \text{cl}(B - N(T))$ gives a disc as desired. ■

Suppose first that F contains a surface which contains an arc $\lambda \subset T$ properly imbedded in B . We take the arc λ to be outermost in F , that is, there is a regular neighbourhood Q of v in F such that $\lambda \cap Q$ and a subarc of ∂Q cobound a disc Q' with $Q' \cap T = \lambda \cap Q'$. Then λ and an arc in ∂B together form a loop bounding a disc D in B such that $D \cap T = \lambda$. By Claim 2.3, we can take D so that D intersects F precisely in λ near λ . Since F is T -incompressible and T - ∂ -incompressible, by a standard innermost loop and outermost arc argument we can see that we can take D so that $D \cap F = \lambda$. We isotope D near λ slightly so that $F \cap D$ is an arc λ' parallel to λ . Then λ' and a subarc of $\partial F - T$ together form a loop which bounds a disc D' on $F - T$ since F is T - ∂ -incompressible. Hence the surface of F contains a subdisc of type (3). We repeat the argument as above for every outermost arc in $T \cap F$, to conclude that the surface of F is a disc of type (3) or (4).

Suppose secondly that F contains a surface which contains an edge e of T as a connected component of $T \cap F$. Then one of the endpoints v of e is in the interior of F and the other endpoint of e is in ∂F . Hence similar argument as in the previous paragraph shows that the component is of type (5).

Thus we can assume that F is disjoint from T . There is a disc R which is properly imbedded in B and contains T . Since F is T -incompressible and T - ∂ -incompressible, we can take R so that it is disjoint from F . Then we obtain two balls by cutting B along R , and F is properly imbedded in the two balls. Hence F consists of surfaces of types (1) and (2). ■

Let C be a compression body, and T a trivial graph properly imbedded in C . We cap off each sphere component of $\partial_- C$ with a 3-ball if it is disjoint from T , to obtain a new compression body denoted by \tilde{C} in the next lemma. Discs and annuli imbedded in (C, T) are called *vertical* if they are disjoint union of vertical arcs with respect to some homeomorphism $C \cong Y \cup V$ as in the definition for T to be trivial.

Lemma 2.4. *Let C be a compression body, and T a trivial graph properly imbedded in C . Let F be a union of compact surfaces in C . Suppose for every surface F_0 of F that F_0 is imbedded in C , $\partial F_0 \subset (\partial C \cup T)$, $(F_0 \cap \partial C) \subset \partial F_0$ and $F_0 \cap T$ is a union of edges of T . In addition, suppose that distinct surfaces of F may intersect in a union of edges of T . If F is T -incompressible and T - ∂ -incompressible, then F is a union*

of

- (1) *spheres disjoint from T ,*
- (2) *properly imbedded discs disjoint from T ,*
- (3) *cancelling discs of ∂_+ -parallel arc components of T ,*
- (4) *properly imbedded discs which are unions of two cancelling discs,*
- (5) *discs such that the boundary loop ℓ of each of them intersects ∂C in an arc α and such that the arc $cl(\ell - \alpha)$ is in T ,*
- (6) *discs which are unions of discs of the above type (5),*
- (7) *properly imbedded discs each of which intersects T in a single edge of T with an endpoint in its interior,*
- (8) *vertical discs each of which intersects T in two or more vertical arc components,*
- (9) *vertical annuli each of which intersects T in zero or more vertical arc components and*
- (10) *closed surfaces parallel to a component of $\partial_- \tilde{C}$ disjoint from T .*

Proof. Let t_1, \dots, t_n be the components of T such that each t_i contains a vertex of T besides ∂t_i . There are discs D_1, \dots, D_n properly imbedded in C such that D_i contains t_i for $i = 1, \dots, n$. We can take these discs so that D_i cuts off a ball B_i from C with $B_i \cap T = D_i \cap T = t_i$ for $i = 1, \dots, n$. We take parallel copies D'_1, \dots, D'_n of these discs so that D'_i cuts off a ball B'_i from C with $B'_i \cap T = t_i$ for $i = 1, \dots, n$. Since F is T -incompressible and T - ∂ -incompressible, by a standard innermost loop and outermost arc argument on $F \cap (\cup_i D'_i)$, we can deform F to be disjoint from $\cup_i D'_i$ as usual. Note that this operation does not change whether F is a union of surfaces of types (1)–(10) in the statement of this lemma or not.

By Lemma 2.2 the components of F in B'_i are surfaces of types in the statement of this lemma. By similar argument as in [Lemma 2.1, HS2] the remainder components of F are also surfaces of types in the statement of this lemma. ■

The next theorem is a generalization of Haken's theorem [Theorem in section 7, H]. The proof of the next theorem is the same as that of [Theorem 1.3, HS1]. We omit the proof. The only difference is that we use the above Lemma 2.4 in this paper instead of Lemma 2.4 in [HS1].

Let M be a compact connected orientable 3-manifold, and T a properly imbedded graph in M . The pair (M, T) is *split* if the complement $M - T$ contains an essential sphere S , that is, S does not bound a ball in $M - T$. This sphere S is called a *splitting sphere*.

Theorem 2.5. *Let H be a Heegaard splitting of (M, T) , and S a disjoint union of splitting spheres in (M, T) and T -compressing discs of ∂M . Then there is a disjoint union S' of splitting spheres and T -compressing discs such that*

- (1) S' is obtained from S by 2-surgeries and isotopy in (M, T) ,
- (2) each sphere of S' intersects H in at most one simple closed curve and
- (3) each disc of S' intersects H in precisely one simple closed curve essential on $H - T$.

Here a 2-surgery on S is such an operation as taking a disc D with $D \cap S = \partial D$ and $D \cap T = \emptyset$ and replacing a small annular neighbourhood of ∂D in S by two parallel copies of D .

3 Proof of Theorem 1.1, I

Let M be a compact 3-manifold possibly with boundary, T a graph properly imbedded in M , and H a Heegaard splitting surface of (M, T) . Let C_1 and C_2 be compression bodies obtained by cutting M along H , and set $T_i = T \cap C_i$ for $i = 1$ and 2. The splitting H is said to be *cancellable* if there is a disc D_i imbedded in C_i for $i = 1$ and 2 such that (1) $t_i = D_i \cap T_i \subset \partial D_i$ is an arc properly imbedded in C_i for $i = 1$ and 2, (2) $\text{cl}(\partial D_i - t_i) = D_i \cap \partial_+ C_i$ for $i = 1$ and 2, (3) $D_1 \cap D_2$ consists of one or two points of $T \cap H$ and (4) either t_1 and t_2 are ∂_+ -parallel arc component of T_1 and T_2 respectively, or the components of T_j which are adjacent to t_i are ∂_+ -parallel arc components of T_j for $(i, j) = (1, 2)$

or $(2, 1)$. See Figure 3.1. We call the discs D_1 and D_2 a *cancelling pair of discs*. When $D_1 \cap D_2$ consists of precisely one point, we say that H is *strongly cancellable*. We assume that H is strongly cancellable and t_1 and t_2 are ∂_+ -parallel arc components of T_1 and T_2 respectively. Then we can obtain another Heegaard splitting of (M, T) by isotoping t_2 along the disc D_2 and further isotoping it slightly into C_1 beyond H . Next assume that H is strongly cancellable, that t_2 contains precisely one vertex v of T with its valency equal to 3 and that the components of T_1 adjacent to t_2 are ∂_+ -parallel arcs. Let e_1 and e_2 be the edges of T_2 which together compose t_2 such that e_2 is adjacent to $t_1 = T \cap D_1$ on T . Then we can obtain another Heegaard splitting of (M, T) by isotoping e_2 along the disc D_2 and further isotoping it slightly into C_1 beyond H . In this operation, the vertex v is slid into the compression body C_1 along the edge e_1 . This can also be pursued by moving T by an ambient isotopy of M . We call these two operations *cancelling operations*.




Figure 3.1

Let M be a compact connected orientable 3-manifold, and T a graph properly imbedded in M . A Heegaard splitting $(M, T) = (C_1, T_1) \cup_H (C_2, T_2)$ is said to be *stabilized* if there is a properly imbedded disc D_i disjoint from T_i in C_i for $i = 1$ and 2 such that ∂D_1 and ∂D_2 intersect transversely at a single point in H .

Now we concentrate on our situation. Let (V, T_0) be the pair of the solid torus and the core loop. We take a meridian disc D of V so that D intersects T_0 transversely in exactly one point p . We take an imbedded annulus A such that $T_0 \subset \partial A$, $A \cap \partial V = \partial A - T_0$ and $A \cap D$ is an arc T' connecting the point p and a point in ∂V . Let $T = T_0 \cup T'$ the properly imbedded graph in V with the vertex p of valency 3 and the vertex ∂T

of valency 1.

Theorem 3.1 *Every Heegaard splitting H of (V, T) is stabilized or cancellable. Moreover, H is stabilized when $\text{genus}(H) \geq 2$ and when H is not strongly cancellable. In addition, if two closed surfaces of genus 1 give Heegaard splittings of (V, T) which are not strongly cancellable, then they are isotopic in (V, T) .*

We will give a proof of Theorem 3.1 later. Theorem 1.1 follows from Theorem 3.1 as below.

Proof of Theorem 1.1 assuming Theorem 3.1. Let H be a Heegaard splitting surface of the pair (V, T_0) of the solid torus and the core loop. Let C_1 and C_2 be the compression bodies obtained by cutting V along H . Assume that C_2 contains the boundary torus ∂V . We take a point, say p , in $T_0 - H$. We take a meridian disc D of V and an imbedded annulus A so that D intersects T_0 transversely in the single point p and that $T_0 \subset \partial A$, $A \cap \partial V = \partial A - T_0$ and $A \cap D$ is an arc, say T' . Let $T = T_0 \cup T'$ a graph with the vertex p of valency 3 and the vertex ∂T of valency 1. We will isotope the arc T' so that H gives a Heegaard splitting of (V, T) .

Let X_i be a spine of C_i for $i = 1$ and 2 . Then $M - \text{int}(N(X_1 \cup X_2))$ is homeomorphic to $H \times I$ where $I = [0, 1]$, $H = H \times \{1/2\}$ and $H \times \{0\} \subset C_1$. Let $\pi : H \times I \rightarrow H \times \{1/2\}$ be the projection.

We can move X_i and the arc T' by a small isotopy and take a neighbourhood so that $T \cap N(X_1 \cup X_2) = \emptyset$ and so that T is transverse to H . We can isotope T' so that the singular set of $\pi(T)$ consists of double points disjoint from the vertices p and ∂T . Suppose there is a double point x of $\pi(T') \cap \pi(T_0)$ such that the point $\pi^{-1}(x) \cap T_0$ is contained in an arc, say α , of $T_0 \cap C_2$ and the point $\pi^{-1}(x) \cap T'$ is contained in an arc, say β , of $T' \cap C_2$, and that $\pi^{-1}(x) \cap \beta$ is over $\pi^{-1}(x) \cap \alpha$ with respect to the second coordinate of the product structure $H \times I$. By retaking the point x if necessary, we assume that x is outermost along $\pi(\alpha)$, that is, there is neither such a double point nor the vertex p between x and an endpoint, say y , of $\pi(\alpha)$. Let ρ be the subarc of $\pi(\alpha)$ between x and y . We isotope T' so that $\pi(T')$ is isotoped along ρ . See Figure 3.2. By repeating this operation, we can eliminate such double points. If there is a double point z of $\pi(T') \cap \pi(T_0)$ such that the point of $\pi^{-1}(z) \cap T_0$ is contained in an arc, say γ , of $T_0 \cap C_1$ and the point $\pi^{-1}(z) \cap T'$ is

contained in an arc, say δ , of $T' \cap C_1$, and that $\pi^{-1}(z) \cap \delta$ is under $\pi^{-1}(z) \cap \gamma$, then in the same way we can eliminate z . Note that these operations do not yield such a new double point as above. Hence these repetitions terminate in finite times.




Figure 3.2

Then as in the proof of Lemma 2.1, we let S be the set consisting of the vertices p , ∂T and singular points of T' with respect to the projection map π . We take regular points of T' , one between every pair of adjacent points of S . We take regular points of T' , one more between every pair of adjacent points of S if both points are Then we isotope T' so that $T' \cap H$ consists of the above regular points and $T_i = T \cap C_i$ is trivial in C_i for $i = 1$ and 2 .

Now, H gives a Heegaard splitting of (V, T) . By Theorem 3.1, H is stabilized or cancellable. Moreover, H is stabilized when $\text{genus}(H) \geq 2$ and when H is not strongly cancellable.

If H is stabilized as a Heegaard splitting of (V, T) , then it is also stabilized as a Heegaard splitting of (V, T_0) .

Suppose that the Heegaard splitting H of (V, T) is cancellable. Let $D_1 \subset C_1$ and $D_2 \subset C_2$ be a cancelling pair of discs, and set $t_i = D_i \cap T$ for $i = 1$ and 2 . Then t_i is an arc properly imbedded in C_i . (Note that ∂C_i does not go through an edge of T_i two or more times by condition (1) of the definition of cancellability.) There are three cases.

- (1) $(t_1 \cup t_2) \subset T'$,
- (2) $t_1 \cup t_2$ is entirely contained in neither T_0 nor T' , or
- (3) $(t_1 \cup t_2) \subset T_0$.

In the cases of (1) and (2), since T' is an arc, $D_1 \cap D_2$ is a single point of $T \cap H$. Hence in both cases we can obtain a new Heegaard splitting

H' of (V, T) by a cancelling operation such that $|T' \cap H'| = |T' \cap H| - 2$ in the case (1) and $|T' \cap H'| = |T' \cap H| - 1$ in the case (2). (This holds even when t_2 is an arc component of $T_0 \cap C_2$, and when t_1 is a union of a subarc of $T_0 \cap C_1$ and a subarc t'_1 of T' . Because the component of $T' \cap C_2$ which is adjacent to the subarc t'_1 is a ∂_+ -parallel arc component by condition (4) of the definition of cancellability.) By repeating these operations, we can assume that the cases (1) and (2) never occur.

Hence the case (3) happens. Then it follows that H is also cancellable as a Heegaard splitting of (V, T_0) . (This holds even when $t_1 \cup t_2$ contains the vertex p since all we have to do is forgetting the arc T' .) When $|H \cap T_0| = 2$ and $\text{genus}(H) \geq 2$, H is not strongly cancellable as a Heegaard splitting of (V, T) . Hence H is stabilized by Theorem 3.1. ■

We will prove Theorem 3.1 in this and the next sections.

Lemma 3.2. *The Heegaard splitting H of (V, T) is stabilized if it is T -reducible.*

Proof. If H is T -reducible, then there is a sphere S disjoint from T and intersecting H in a single simple closed curve essential in $H - T$. Since V is irreducible, T is connected and $T \cap \partial V \neq \emptyset$, S bounds a ball disjoint from T . Hence H is stabilized by [Satz 3.1, W]. ■

Let C_1, C_2 be the compression bodies obtained by cutting V along the Heegaard splitting surface H , and set $T_i = T \cap C_i$ for $i = 1$ and 2 . Assume that C_2 contains the boundary torus ∂V . Then C_1 is a handlebody. We can isotope $D \cup A$ fixing on T so that $D \cup A$ intersects C_1 in a union of T_1 -incompressible and T_1 - ∂ -incompressible surfaces. Moreover, we take $D \cup A$ so that the number $|(D \cup A) \cap H|$ of the intersection loops and arcs is minimal. Then $(D \cup A) \cap C_1$ is a union of surfaces of types (2)–(7) in the statement of Lemma 2.4. The discs of type (2) are T_1 -compressing discs of $\partial_+ C_1$ since $|(D \cup A) \cap H|$ is minimal. In addition, a standard cut and paste argument shows that $(D \cup A) \cap C_2$ is T_2 -incompressible in (C_2, T_2) by the minimality of $|(D \cup A) \cap H|$.

Proposition 3.3. *The intersection $(D \cup A) \cap C_1$ is a union of surfaces of types (3)–(7) in the statement of Lemma 2.4.*

We utilize a method introduced by W. Haken in [section 7, H], and extended by W. Jaco in [Theorem II.7, J], M. Ochiai in [O] and H. Doll in [D].

Proof. We assume for a contradiction that $(D \cup A) \cap C_1$ contains a disc of type (2). Then $(D \cup A) \cap C_2$ contains a T_2 -incompressible non-disc planar surface with a boundary component disjoint from T and in $\partial_+ C_2$, and hence has a T_2 - ∂ -compressing disc D_1 by Lemma 2.4. Set $R_0 = D \cup A$. We can isotope a small neighbourhood of the arc $\alpha_1 = D_1 \cap R_0$ along the disc D_1 into C_1 . Then the resultant union of surfaces in C_2 are homeomorphic to the surfaces obtained by cutting $R_0 \cap C_2$ along the arc α_1 , and a band is attached to the surfaces $R_0 \cap C_1$ in C_1 . Let R_1 be the resultant union of surfaces in V . Note that $R_1 \cap C_2$ is T_2 -incompressible. Again, it has a T_2 - ∂ -compressing disc D_2 if it does not consist of surfaces of types (2)–(7) and (9). Thus we can perform isotopy along a T_2 - ∂ -compressing disc D_i of $R_{i-1} \cap C_2$ for $i = 1, \dots, n$, until $R_n \cap C_2$ consists of surfaces of types (2)–(7) and (9). Then the arcs $\alpha_i = D_i \cap R_{i-1} \subset R_0 \cap C_2$ divide $R_0 \cap C_2$ into surfaces of types (2)–(7) and (9). We can retake the T_2 - ∂ -compressing discs D_1, \dots, D_n so that the endpoints of $\alpha_1, \dots, \alpha_n$ are in the boundaries of the surfaces $R_0 \cap C_1$. Note that each α_i is essential in $(R_0 \cap C_2) - T_2$. The arc α_i is called a *self-connector* if it has distinct endpoints in the same component of $(R_0 \cap C_1) - T_1$. Suppose that every T_1 -compressing disc of $\partial_+ C_1$ in $R_0 \cap C_1$ has a self-connector incident to it. Then there must be an inessential arc α_i in $(R_0 \cap C_2) - T_2$ because $R_0 - T$ is a union of discs. This is a contradiction. Hence there is a T_1 -compressing disc Q among $R_0 \cap C_1$ to which no self-connector is incident. Let α_k be the first arc incident to Q . The arc α_k connects Q to another surface Q' of $R_0 \cap C_1$. Then the argument in Lemma 1 in [O] or Lemma 3.3(a) in [D] shows that we can reduce the number $|(D \cup A) \cap H|$ as below.

The ∂ -compression of $R_{i-1} \cap C_2$ along the disc D_i can be reversed by a ∂ -compression of $R_i \cap C_1$. We call it the *dual ∂ -compression*. Let D'_i denote the *dual disc* for D_i , that is, one which gives the dual ∂ -compression.

First suppose that $D_k \cap R_0 = \alpha_k$. Then an isotopy of R_0 along D_k would reduce $|R \cap H|$.

Secondly suppose that $(D_k \cap R_0) - \alpha_k \neq \emptyset$. Then there are dual ∂ -compressing discs of $R_{k-1} \cap C_1$ intersecting $\beta_k = D_k \cap H$ among D'_1, \dots, D'_{k-1} . If we isotope R_{k-1} along the discs D'_{k-1}, \dots, D'_1 in this order, then we obtain R_0 . The interior of the discs D'_1, \dots, D'_{k-2} may intersect R_{k-1} . The arc β_k connects Q and Q' . Let D'_j ($j < k$) be the

dual disc intersecting β_k and the nearest to Q . Let E be the subdisc of $\partial N(\beta_k) \cap C_1$ between Q and D'_j . We retake the dual disc D'_j so that the subdisc $D'_j \cap N(\beta_k)$ is replaced by $E \cup (Q - N(\beta_k))$, and we isotope the new D'_j slightly off of Q . We perform the above deformation of dual discs for each intersection point of dual discs and β_k , so that dual discs D'_1, \dots, D'_k are disjoint from β_k . Then we can isotope the disc R_{k-1} along the discs D'_i ($i = k - 1, \dots, 1$), and further isotope along the disc D_k , to reduce $|R \cap H|$. This is a contradiction. ■

4 Proof of Theorem 1.1, II

By Proposition 3.3, we can take $R = D \cup A$ so that it intersects C_1 in surfaces of types (3)–(7) in the statement of Lemma 2.4. Let Q be a spine of (C_1, T_1) . We can take Q to be a connected 1-complex which intersects R in vertices of Q so that $(Q \cap R) \subset T$. We think of C_1 as being a very small neighbourhood of Q . Let G be the ball obtained by cutting the solid torus V along the union R of the meridian disc D and the annulus A . Then $Q_G = Q \cap G$ is a graph.

The next lemma is essentially Proposition 2.5 in [ST], and this proof is essentially due to [F].

Lemma 4.1. *We can assume that every component of Q_G is a tree. Otherwise, H is stabilized.*

Proof. Suppose not. Then Q_G contains a (possibly knotted) simple closed curve c . Let e be an edge of Q_G contained in c . We slide an endpoint q of e along the arc $c - e$, to deform e into a loop edge. We slide q slightly on e passing through the other endpoint, and obtain a loop $\ell \subset e$ imbedded in the interior of G . We take a small neighbourhood $N(\ell)$ of ℓ in C_1 . Then $C_1 - \text{int}N(\ell)$ is a compression body, and H is a Heegaard splitting of $(V - \text{int}N(\ell), T)$. We push the sphere ∂G slightly into $\text{int}G$ and obtain a splitting sphere S of $(V - \text{int}N(\ell), T)$, that is, S does not bound a ball in $V - \text{int}N(\ell) - T$. Then by Theorem 2.5 there is a sphere S' which is disjoint from T and intersecting H in a single simple closed curve essential in $H - T$. Then H is T -reducible as a splitting of $(V - \text{int}N(\ell), T)$, hence also as a splitting of (V, T) . Thus the splitting H is stabilized by Lemma 3.2. ■

Since T_0 is a loop, $Q \cap T_0 \neq \emptyset$. In addition, $Q \cap T' \neq \emptyset$ since the arc

T' has an endpoint in $\partial V = \partial_- C_2$ and the other endpoint at the vertex $p = T_0 \cap T'$. However, it is possible that $Q \cap T' = p$. Because $R \cap C_1$ is a union of surfaces of types (3)–(7) in the statement of Lemma 2.4, each of the discs obtained by cutting D and A along T' intersects C_2 in a disc. Hence the union of surfaces $R \cap C_2$ is T_2 -incompressible. Then $R \cap C_2$ is T_2 - ∂ -compressible by Lemma 2.4. Note that $A \cap C_2$ is T_2 - ∂ -compressible even when $Q \cap T = p$. We can isotope R along T_2 - ∂ -compressing discs D_1, \dots, D_n to deform $R \cap C_2$ into a union of surfaces of types (3)–(7) and (9) in the statement of Lemma 2.4. Note that the disc of type (2) cannot appear since $R \cap C_1$ does not have the disc component of type (2) and since D_1, \dots, D_n are T_2 - ∂ -compressing discs. Set $R_0 = R$, and let R_i be the resultant union of surfaces obtained by the i -th isotopy along the disc D_i . Let γ_i be the arc $R_{i-1} \cap \partial D_i$, and δ_i the arc $\partial D_i \cap \partial_+ C_2$. We can isotope the discs D_1, \dots, D_n so that the endpoints of $\gamma_1, \dots, \gamma_n$ are in the boundary of the discs $R_0 \cap C_1$. Let Γ_i be the graph with the edges $\cup_{j=1}^i \gamma_j$, regarding the disc components of $R_0 \cap C_1$ as vertices. Let $\tilde{\Gamma}_i$ be the intersection of Γ_i and the sphere ∂G . The sphere ∂G contains two copies of D , two copies of A cut along T' , a copy of T_0 cut at p and four copies of T' . Note that the graph $\tilde{\Gamma}_i$ contains two copies of each edge γ_j . See Figure 4.1.




Figure 4.1

We concentrate on the patterns of $\tilde{\Gamma}_i$ as below.

- (1) An edge e of $\tilde{\Gamma}_i$ in A .
- (2) An edge e of $\tilde{\Gamma}_i$ in D such that e connects two points in the same copy of T' (Figure 4.2).

- (3) Two edges of $\tilde{\Gamma}_i$ in a copy of D such that they connects distinct copies of T' and such that they have an endpoint at the same point (Figure 4.3).




Figure 4.2




Figure 4.3

We allow the edge e of the above types (1) and (2) to be incident to the vertex $p = T_0 \cap T'$. Let k be the smallest number such that $\tilde{\Gamma}_k$ contains such a pattern. Note $i < k$, that $\tilde{\Gamma}_i$ is a disjoint union of arcs for $i < k$.

We perform the i th isotopy of R_{i-1} along the T_2 - ∂ -compressing disc D_i for $i \leq k$ by sliding edges of the spine Q as below. (The description of deformation is completed right after Lemma 4.3.) Let Q_i denote the spine after the i th deformation. The original spine is denoted by Q_0 . We will deform the spine so that $Q_i \cap R_i = \Gamma_i$. That is, we allow i edges of Q_i to be contained in R_i . Let Q_{G_i} denote the graph $Q_i \cap G$. The edges $Q_i \cap R_i = \Gamma_i$ appear twice in the graph Q_{G_i} .

Lemma 4.2. *We can assume that every component of Q_{G_i} is a tree for $i < k$. Otherwise, H is stabilized.*

Proof. The proof is similar to that of Lemma 4.1, and we omit the detail. Suppose that Q_{G_i} contains a cycle c . Since $\tilde{\Gamma}_i$ does not contain a cycle, the cycle c contains an edge e such that e is not contained in $\tilde{\Gamma}_i$. We slide the edge e as in the proof of Lemma 4.1, and the same argument shows that H is stabilized. ■

We can assume that the condition of Lemma 4.2 are kept after any sliding operations on edges of $Q_{G_i} - \tilde{\Gamma}_i$ in G . Otherwise, H is stabilized.

We can extend the T_2 - ∂ -compressing disc D_i so that $\delta_i \subset Q_{i-1}$, and isotope D_i so that δ_i is a sequence of edges of $Q_{G(i-1)}$, where the same edge possibly appears more than once.

Lemma 4.3. *We can deform Q_{i-1} by sliding edges of $Q_{G(i-1)} - \tilde{\Gamma}_{i-1}$ and isotope D_i so that δ_i does not intersect an edge of $Q_{G(i-1)}$ more than once.*

Proof. Let x and y be points of δ_i which are in the same vertex v of $Q_{G(i-1)}$. We can choose x and y so that the subarc ρ connecting x and y in δ_i contains no such pair of points. Then by Lemma 4.2 ρ consists of two successive subarcs of δ_i contained in the same edge of $Q_{G(i-1)}$. See Figure 4.4. Now, we can easily slide edges of $Q_{G(i-1)} - \tilde{\Gamma}_{i-1}$ and isotope D_i to shrink ρ into a point at the vertex v . ■




Figure 4.4

Then for $1 \leq i \leq k$ we perform the i th deformation of the spine Q_{i-1} along D_i as below. Note that ∂D_i cannot entirely be contained in ∂G since every connected component of $\tilde{\Gamma}_i$ is an edge in a copy of D such that it connects distinct copies of T' . Hence δ_i contains an edge e of $Q_{G(i-1)} - \tilde{\Gamma}_{i-1}$, we slide the edge e along the arcs $\delta_i - e$ and isotope along the disc D_i , to push e into ∂G . Then e is in the place of γ_i .

We consider the case where a copy of γ_k is of type (3) as in Figure 4.3. In the case of type (2), the proof is similar to that of this case and easier, and we omit it.

When a copy of γ_k is of type (3), in the graph $\tilde{\Gamma}_k$ a copy of the edge γ_k and a copy of some edge γ_m have an endpoint in the same vertex v of Q_{Gk} . We call these copies of edges γ_m and γ_k for simplicity.

The other endpoints $v_k \subset \partial\gamma_k$ and $v_m \subset \partial\gamma_m$ are in distinct vertices in the same copy t of T' . Let t' be the subarc of t between v_k and v_m . See Figure 4.5.

Lemma 4.4. *If γ_k is of type (3), then H is strongly cancellable.*

Proof. Let E be the disc bounded by $t' \cup \gamma_k \cup \gamma_m$ in R_k right after the k th deformation. Note that E is disjoint from the other edges of $\tilde{\Gamma}_k$ than γ_m and γ_k because of the definition of the number k .

First, suppose that t' contains no other vertices than v_k and v_m . Then t' is a ∂_+ -parallel arc component of T_2 and E gives a cancelling disc of t' in (C_2, T_2) . The vertex v_m does not coincide with the vertex p . Hence the ball neighbourhood $N(v_m)$ contains a cancelling disc E' of the ∂_+ -parallel arc of T_1 corresponding to v_m . We can take E' so that E' is perpendicular to the sphere ∂G inside G and so that E' is disjoint from the interior of the edges of Q_k . Then E and E' give a cancelling disc pair, and we see that H is strongly cancellable.

Secondly, suppose that t' contains other vertices than v_k and v_m . Let w_1, \dots, w_r be the vertices contained in t' other than v_k and v_m , appearing in this order in t' so that w_1 is the nearest to v_k . The arc t' is divided into $r + 1$ arcs when it is cut at the vertices w_1, \dots, w_r . Let t_i be the subarc of t' between w_{i-1} and w_i , where $v_k = w_0$ and $v_m = w_{r+1}$. The arc t_i is a ∂_+ -parallel arc component of T_2 . Let E_i be a cancelling disc of t_i in (C_2, T_2) . We can take E_1, \dots, E_{r+1} so that they are disjoint and they intersect the disc E only properly imbedded arcs connecting vertices of Q_k . Let $\mathcal{E} = \cup E_i$. See Figure 4.5.

If several arcs of $\mathcal{E} \cap E$ are incident to the vertex v , then we isotope \mathcal{E} along the edge γ_k to deform such arcs into arcs incident to the vertex v_k . See Figures 4.6 and 4.7.

Suppose there is an arc ρ of $\mathcal{E} \cap E$ such that ρ has two endpoints in the same vertex q . Then we can take ρ to be the outermost one among such arcs on E . That is, there is a disc E' cut off by ρ from E such that E' is disjoint from such arcs other than ρ . We perform a surgery on \mathcal{E}




Figure 4.5




Figure 4.6




Figure 4.7

along the disc E' and discard the component disjoint from T_2 , to obtain a union of cancelling discs which intersects E in fewer number of arcs. Hence we can assume that $\mathcal{E} \cap E$ does not contain such an arc.

We say that an arc of $\mathcal{E} \cap E$ is *good* if it connects adjacent vertices. We say a vertex w_i is *good* if all the arcs incident to w_i are good. There is such a good vertex w_i since every polygon has at least two vertices with no diagonal arc incident when divided into triangles by diagonal arcs. Then all the arcs incident to w_i have the other endpoint in the adjacent vertices w_{i-1} and w_{i+1} . Hence there is a cancelling disc E_i such that (a) E_i is a cancelling disc of t_i or t_{i+1} , (b) E_i intersects E transversely in arcs, and (c) only good arcs of $E_i \cap E$ are incident to w_i . Among such discs, we take E_i so that the number of intersection arcs $E_i \cap E$ is minimum.

We will show that no arc of $E_i \cap E$ is incident to w_i . Suppose there is an arc μ of $E_i \cap E$ either connecting w_{i-1} and w_i or connecting w_i and w_{i+1} . We consider the former case. The proof of latter case is similar and we omit it. We take μ to be the outermost one on E . The arc μ divides E_i into two discs, one of which, say E'_i intersects neither t_i nor t_{i+1} . Let E' be the subdisc of E cobounded by t_i and μ . Then $E' \cup E'_i$ is a cancelling disc of t_i which intersects in fewer number of arcs than E_i after an adequate small isotopy. This is a contradiction.

Thus there is no arc of $E_i \cap E$ which is incident to w_i . Let $N(w_i)$ be a ball neighbourhood of the vertex w_i . The disc $N(w_i) \cap E$ is a cancelling disc of the arc component of T_1 corresponding to the vertex w_i . Then the discs E_i and $N(w_i) \cap E$ show that H is strongly cancellable. ■

The proof of the next lemma is similar to that of Lemma 4.4. We omit it.

Lemma 4.5. *Suppose that γ_k is of type (2). Then the Heegaard splitting H is strongly cancellable.*

The next lemma completes the proof of the first sentence of Theorem 3.1.

Lemma 4.6. *Suppose that γ_k is of type (1). Then the Heegaard splitting H is cancellable. Moreover, if H is not strongly cancellable, then $T \cap Q = p$ and γ_k is an edge connecting the two copies of p in a copy of A .*

Proof. The arc T' and the edge γ_k divides the annulus A into two discs, one of which, say E , is disjoint from the toral boundary ∂V .

Then similar argument as in the proof of Lemma 4.4 will show that the Heegaard splitting H is cancellable.

When the arc $\partial E - \gamma_k$ does not contain a vertex of Q_{Gk} , then we can take a disc E' in $N(\partial\gamma_k)$ so that E and E' together form a cancelling pair of discs. In addition, if $\partial\gamma_k = p$, then we can take a cancelling disc of the arc $T_0 \cap N(p)$ in $N(p)$, and see that H is cancellable. Since we can also take a cancelling disc of the arc $\partial E \cap N(p)$ in $N(p)$, we see that H is strongly cancellable unless $T \cap Q = p$ and γ_k is an edge connecting the two copies of p in a copy of A .

When the arc $\partial E - \gamma_k$ contains a vertex of Q_{Gk} other than $\partial\gamma_k$, the proof is very similar as in that of Lemma 4.4. ■

The next lemma implies the second and third sentences of the statement of Theorem 3.1.

Lemma 4.7. *If H is not stabilized and not strongly cancellable, then $\text{genus}(H) = 1$. Moreover, such splitting surfaces are isotopic in (V, T) .*

Proof. Suppose that H is not stabilized and not strongly cancellable. Then by Lemmas 4.4, 4.5 and 4.6 the edge γ_k is contained in a copy of A and connects two copies of the vertex $p = T_0 \cap T' = T \cap Q_k$ as in the statement of Lemma 4.6. Then before the deformation Q_G is a connected tree intersecting ∂G in the two copies of the vertex p by Lemma 4.1. We can shrink Q_G to be an edge e connecting the two copies of p . Thus Q consists of a loop edge incident to p , and this implies that $\text{genus}(H) = 1$. Moreover, the edge e is not knotted in G since C_2 is homeomorphic to $(\text{a torus}) \times I$. ■

References

- [BO] F. Bonahon and J.-P. Otal, *Scindements de Heegaard des espaces lenticulaires*, Ann. Sci. Ec. Norm. Sup. **16(4)** (1983), 451-466.
- [CG] A.J. Casson and C.McA. Gordon, *Reducing Heegaard splittings*, Topology Appl. **27** (1987), 275-283.
- [D] H. Doll, *A generalized bridge number for links in 3-manifolds*, Math. Ann. **294** (1992), 701-717.
- [F] C. Frohman, *The topological uniqueness of triply periodic minimal surfaces in \mathbb{R}^3* , J. Differ. Geom. **31** (1990), 277-283.

- [G] H. Goda, *Bridge index for theta curves in the 3-sphere*, *Topology Appl.* **79** (1997), 177-196.
- [H] W. Haken, *Some results on surfaces in 3-manifolds*, *Studies in Modern Topology* (Math. Assoc. Amer., distributed by: Prentice-Hall) (1968), 34-98.
- [HS1] C. Hayashi and K. Shimokawa, *Thin position for 1-submanifold in 3-manifold*, *Pacific J. Math.* **197** (2001), 301-324.
- [HS2] C. Hayashi and K. Shimokawa, *Heegaard splittings of the trivial knot*, *J. Knot Ramifi.* **7** (1998), 1073-1085.
- [HS3] C. Hayashi and K. Shimokawa, *Heegaard splittings of trivial arcs in compression bodies*, *J. Knot Ramifi.* **10** (2001), 71-87.
- [J] W. Jaco, *Lectures on three-manifold topology*, (CBMS Reg. Conf. Ser., vol 43). Providence, RI: Am. Math. Soc., 1980.
- [O] M. Ochiai, *On Haken's Theorem and its extension*, *Osaka J. Math.* **20** (1983), 461-468.
- [ST] M. Scharlemann and A. Thompson, *Heegaard splitting of (surface) \times I are standard*, *Math. Ann.* **295** (1993), 549-564.
- [S] H. Schubert, *Ueber eine numerische Knoteninvariante*, *Math. Z.* **61** (1954), 245-288.
- [W] F. Waldhausen, *Heegaard-Zerlegungen der 3-Sphäre*, *Topology* **7** (1968), 195-203.

Department of Mathematics
 Faculty of Science
 Gakushuin University
 Graduate School of Information Sciences
 Tohoku University

1-5-1 Mejiro Toshima-ku Tokyo, 171, Japan
E-mail: Chuichiro.Hayashi@gakushuin.ac.jp

2-1-1 Katahira Aoba-ku Sendai, 980-8577, Japan
E-mail: koya@math.is.tohoku.ac.jp

Recibido: 20 de Marzo de 2000
 Revisado: 4 de Diciembre de 2000