

The starlikeness and convexity of multivalent functions involving certain inequalities

Hüseyin IRMAK⁽¹⁾ and R. K. RAINA⁽²⁾

(1) Department of Primary Education
Faculty of Education
Baskent University, Baglica Campus
Baglica-Etimesgut, Tr-06530, Ankara, Turkey
hisimya@baskent.edu.tr

(2) Department of Mathematics
College of Technology and Engineering
Maharana Pratap University of Agriculture and Technology
Udaipur 313 001, Rajasthan, India
rainark.7@hotmail.com

Recibido: 10 de Junio de 2002
Aceptado: 18 de Diciembre de 2002

ABSTRACT

In the present paper, a theorem for the starlikeness and convexity of multivalent functions involving certain inequalities is given. Some interesting consequences of the main result are also mentioned.

2000 Mathematics Subject Classification: Primary 30C45.

Key words: Open unit disc, analytic, multivalently starlike, multivalently convex, and Jack's Lemma

1. Introduction and definitions

Let $\mathcal{T}(p)$ denote the class of functions $f(z)$ of the form

$$f(z) = z^p + \sum_{k=p+1}^{\infty} a_k z^k, \quad (p \in \mathcal{N} = \{1, 2, 3, \dots\}), \quad (1.1)$$

which are *analytic* and *multivalent* in the open unit disc $\mathcal{U} = \{z : z \in \mathcal{C} \text{ and } |z| < 1\}$. A function $f(z)$ belonging to $\mathcal{T}(p)$ is said to be *multivalently starlike of order α* in \mathcal{U} if it satisfies the inequality:

$$\Re \left\{ \frac{z f'(z)}{f(z)} \right\} > \alpha, \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}), \quad (1.2)$$

and, a function $f(z) \in \mathcal{T}(p)$ is said to be *multivalently convex of order α* in \mathcal{U} if it satisfies the inequality:

$$\Re \left\{ \left(1 + \frac{zf''(z)}{f'(z)} \right) \right\} > \alpha, \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}). \quad (1.3)$$

For the aforementioned definitions, one may refer to [1] and [2] (see also [11]). Further, a function $f(z) \in \mathcal{T}(p)$ is said to be in the subclass $\mathcal{T}_\lambda(p; \alpha)$ if it satisfies the inequality

$$\Re \left\{ \frac{zf'(z) + \lambda z^2 f''(z)}{(1-\lambda)f(z) + \lambda z f'(z)} \right\} > \alpha, \quad (1.4)$$

$$(z \in \mathcal{U}; 0 \leq \lambda \leq 1; 0 \leq \alpha < p; p \in \mathcal{N}).$$

From the above definitions, the following subclasses of the classes $\mathcal{T}(p)$ and $\mathcal{T} \equiv \mathcal{T}(1)$ emerge from the families of functions $\mathcal{T}_\lambda(p; \alpha)$:

$$\mathcal{T}_\lambda(1; \alpha) = \mathcal{T}_\lambda(\alpha), \quad (0 \leq \lambda \leq 1; 0 \leq \alpha < 1), \quad (1.5)$$

$$\mathcal{T}_0(p; \alpha) = \mathcal{S}_p(\alpha), \quad (0 \leq \alpha < p; p \in \mathcal{N}), \quad (1.6)$$

$$\mathcal{T}_1(p; \alpha) = \mathcal{K}_p(\alpha), \quad (0 \leq \alpha < p; p \in \mathcal{N}), \quad (1.7)$$

$$\mathcal{S}_p(\alpha) \subseteq \mathcal{S}_p(0) = \mathcal{S}_p, \quad (0 \leq \alpha < p; p \in \mathcal{N}) \quad (1.8)$$

$$\mathcal{K}_p(\alpha) \subseteq \mathcal{K}_p(0) = \mathcal{K}_p, \quad (0 \leq \alpha < p; p \in \mathcal{N}), \quad (1.9)$$

$$\mathcal{T}_0(\alpha) = \mathcal{S}_1(\alpha) \subseteq \mathcal{S}(\alpha), \quad (0 \leq \alpha < 1), \quad (1.10)$$

$$\mathcal{T}_1(\alpha) = \mathcal{K}_1(\alpha) \subseteq \mathcal{K}(\alpha), \quad (0 \leq \alpha < 1), \quad (1.11)$$

$$\mathcal{T}_0(\alpha) = \mathcal{S}(\alpha) \subseteq \mathcal{S}(0) = \mathcal{S}, \quad (0 \leq \alpha < 1), \quad (1.12)$$

$$\mathcal{T}_1(\alpha) = \mathcal{K}(\alpha) \supseteq \mathcal{K}(0) = \mathcal{K}, \quad (0 \leq \alpha < 1). \quad (1.13)$$

The important subclasses in the Geometric function theory such as the multivalently starlike functions $\mathcal{S}_p(\alpha)$ of order α ($0 \leq \alpha < p; p \in \mathcal{N}$) in \mathcal{U} , the multivalently convex functions $\mathcal{K}_0(\alpha)$ of order α ($0 \leq \alpha < p; p \in \mathcal{N}$) in \mathcal{U} , the multivalently starlike functions \mathcal{S}_p in \mathcal{U} , the multivalently convex functions \mathcal{K}_p in \mathcal{U} , the starlike functions in $\mathcal{S}(\alpha)$ of order α ($0 \leq \alpha < 1$) in \mathcal{U} , the convex functions $\mathcal{K}(\alpha)$ of order α ($0 \leq \alpha < 1$) in \mathcal{U} , the starlike functions \mathcal{S} in \mathcal{U} , and the convex functions \mathcal{K} in \mathcal{U} , are seen to be easily identifiable with the aforementioned classes (*cf.*, *e.g.*, [1], [2], and [11]).

The purpose of considering inequality (1.4) is to obtain general results which combine certain types of inequalities concerning functions belonging to the classes $\mathcal{S}_p(\alpha)$, $\mathcal{K}_p(\alpha)$, \mathcal{S}_p , \mathcal{K}_p , $\mathcal{S}(\alpha)$, $\mathcal{K}(\alpha)$, \mathcal{S} , and \mathcal{K} . Some interesting corollaries are also deduced from our main results. Other interesting results involving certain inequalities and/or multivalent functions were also studied, for example, by Owa *et al.* ([8], [9], [10]), and Irmak *et al.* ([3], [4], [5]).

2. Main Result

Before stating and proving our main result, we require the following assertion (popularly known as Jacks's Lemma):

Lemma (cf., [6], [7]). *Let the function $x(z)$ be non-constant and regular in the unit disc \mathcal{U} , such that $w(0) = 0$. If $|w(z)|$ attains its maximum value on the circle $|z| = r < 1$ at a point z_0 , then*

$$z_0 w'(z_0) = c w(z_0), \quad (c \geq 1). \quad (2.1)$$

Theorem. *Let a function $f(z)$ belong to the class $\mathcal{T}(p)$. Define a function $F(z)$ by*

$$F(z) = (1 - \lambda)f(z) + \lambda z f'(z), \quad (0 \leq \lambda \leq 1), \quad (2.2)$$

and if $F(z)$ satisfies anyone of the following inequalities:

$$\left| \frac{1 + \frac{zF''(z)}{F'(z)} - p}{\frac{zF'(z)}{F(z)} - p} - 1 \right| < \frac{1}{2p - \alpha}, \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}), \quad (2.3)$$

$$\left| 1 + z \left(\frac{F''(z)}{F'(z)} - \frac{F'(z)}{F(z)} \right) \right| < \frac{p - \alpha}{2p - \alpha}, \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}), \quad (2.4)$$

$$\left| \frac{F(z)}{zF'(z)} \left(1 + \frac{zF''(z)}{F'(z)} \right) - 1 \right| < \frac{p - \alpha}{(2p - \alpha)^2}, \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}), \quad (2.5)$$

$$\left| \frac{zF'(z)}{F(z)} \left[1 + z \left(\frac{F''(z)}{F'(z)} - \frac{F'(z)}{F(z)} \right) \right] \right| < p - \alpha, \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}), \quad (2.6)$$

$$\Re \left\{ \frac{zF'(z)}{F(z)} \left(\frac{1 + \frac{zF''(z)}{F'(z)} - p}{\frac{zF'(z)}{F(z)} - p} - 1 \right) \right\} < 1, \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}), \quad (2.7)$$

then $f(z) \in \mathcal{T}_\lambda(p; \alpha)$.

Proof. Let $f(z) \in \mathcal{T}(p)$. then from (1.1) and (2.1), we find that

$$\begin{aligned} \frac{zF'(z)}{F(z)} &= \frac{z f'(z) + \lambda z^2 f''(z)}{(1 - \lambda)f(z) + \lambda z f'(z)} \\ &= \frac{p[1 + \lambda(p - 1)] + \sum_{k=p}^{\infty} k[1 + \lambda(k - 1)]a_k z^{k-p}}{1 + \lambda(p - 1) + \sum_{k=p}^{\infty} [1 + \lambda(k - 1)]a_k z^{k-p}}, \end{aligned} \quad (2.8)$$

$(z \in \mathcal{U}; 0 \leq \lambda \leq 1; p \in \mathcal{N}).$

Now, define a function $w(z)$ by

$$\frac{zF'(z)}{F(z)} - p = (p - \alpha)w(z), \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}), \quad (2.9)$$

then the function $w(z)$ is analytic in \mathcal{U} , and $w(0) = 0$. It follows from the above definition (2.9) that

$$1 + \frac{zF''(z)}{F'(z)} - p = (p - \alpha)w(z) \left(1 + \frac{zw'}{w(z)} \frac{1}{p + (p - \alpha)w(z)} \right), \tag{2.10}$$

$(z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N})$.

Hence, from (2.8) and (2.9), we have

$$F_1(z) = \frac{1 + \frac{zF''(z)}{zF'(z)} - p}{\frac{zF'(z)}{F(z)} - p} - 1 = \frac{zw'(z)}{w(z)} \frac{1}{p + (p - \alpha)w(z)}, \tag{2.11}$$

$$F_2(z) = 1 + z \left(\frac{F''(z)}{F'(z)} - \frac{F'(z)}{F(z)} \right) = \frac{(p - \alpha)zw'(z)}{p + (p - \alpha)w(z)}, \tag{2.12}$$

$$F_3(z) = \frac{F(z)}{zF'(z)} \left(1 + \frac{zF''(z)}{F'(z)} \right) - 1 = \frac{(p - \alpha)zw'(z)}{[p + (p - \alpha)w(z)]^2}, \tag{2.13}$$

$$F_4(z) = \frac{zF'(z)}{F(z)} \left[1 + z \left(\frac{F''(z)}{F'(z)} - \frac{F'(z)}{F(z)} \right) \right] = (p - \alpha)zw'(z), \tag{2.14}$$

$$F_5(z) = \frac{zF'(z)}{F(z)} \left(\frac{1 + \frac{zF''(z)}{F'(z)} - p}{\frac{zF'(z)}{F(z)} - p} - 1 \right) = \frac{zw'(z)}{w(z)}, \tag{2.15}$$

$(z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N})$.

We claim that $|w(z)| < 1$ in \mathcal{U} . For otherwise (by Jack's Lemma) there exists a point $z_0 \in \mathcal{U}$ such that

$$z_0w'(z_0) = cw(z_0), \tag{2.16}$$

where

$$|w(z_0)| = 1, (c \geq 1).$$

Therefore, the equations (2.11)-(2.15) in conjunction with (2.16) yield:

$$|F_1(z_0)| = \left| \frac{z_0w'(z_0)}{w(z_0)} \frac{1}{p + (p - \alpha)w(z_0)} \right| = \frac{c|w(z_0)|}{|p + (p - \alpha)w(z_0)|} \geq \frac{1}{2p - \alpha}, \tag{2.17}$$

$$|F_2(z_0)| = \left| \frac{(p - \alpha)z_0w'(z_0)}{p + (p - \alpha)w(z_0)} \right| = \frac{c(p - \alpha)|w(z_0)|}{|p + (p - \alpha)w(z_0)|} \geq \frac{p - \alpha}{2p - \alpha}, \tag{2.18}$$

$$|F_3(z_0)| = \left| \frac{(p - \alpha)z_0w'(z_0)}{p + (p - \alpha)w(z_0)} \right| = \frac{c(p - \alpha)|w(z_0)|}{|p + (p - \alpha)w(z_0)|^2} \geq \frac{p - \alpha}{(2p - \alpha)^2}, \tag{2.19}$$

$$|F_4(z_0)| = c(p - \alpha)|w(z_0)| \geq p - \alpha, \tag{2.20}$$

$$\Re\{F_5(z_0)\} = \Re\left\{ \frac{zw'(z_0)}{w(z_0)} \right\} = c \geq 1, \tag{2.21}$$

$$(0 \leq \alpha < p; p \in \mathcal{N}),$$

which contradict our assumptions (2.3)-(2.7), respectively. Therefore $|w(z)| < 1$ holds true for all $z \in \mathcal{U}$. From the definition (2.9) yields

$$\left| \frac{zF'(z)}{F(z)} - p \right| = (p - \alpha) |w(z)| < p - \alpha, \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}), \tag{2.22}$$

which implies that

$$\Re \left\{ \frac{zF'(z)}{F(z)} \right\} = \Re \left\{ \frac{zf'(z) + \lambda z^2 f''(z)}{(1 - \lambda)f(z) + \lambda z f'(z)} \right\} > \alpha, \tag{2.23}$$

$(z \in \mathcal{U}; 0 \leq \lambda \leq 1; 0 \leq \alpha < p; p \in \mathcal{N}),$

and hence $f(z) \in \mathcal{T}_\lambda(p; \alpha)$.

We mention now some interesting corollaries for the classes $\mathcal{T}_\lambda(\alpha)$, $\mathcal{S}_p(\alpha)$, $\mathcal{K}_p(\alpha)$, \mathcal{S}_p , \mathcal{K}_p , $\mathcal{S}(\alpha)$, $\mathcal{K}(\alpha)$, \mathcal{S} , and \mathcal{K} which are easily deducible from the main result.

Corollary 1. *Let a function $f(z)$ defined by (1.1) belong to the class $\mathcal{T}(p)$. If $f(z)$ satisfies anyone of the following inequalities:*

$$\left| \frac{1 + \frac{zf''(z)}{f'(z)} - p}{\frac{zf'(z)}{f(z)} - p} - 1 \right| < \frac{1}{2p - \alpha}, \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}) \tag{2.24}$$

$$\left| 1 + z \left(\frac{f''(z)}{f'(z)} - \frac{f'(z)}{f(z)} \right) \right| < \frac{p - \alpha}{2p - \alpha}, \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}), \tag{2.25}$$

$$\left| \frac{f(z)}{zf'(z)} \left(1 + \frac{zf''(z)}{f'(z)} \right) - 1 \right| < \frac{p - \alpha}{(2p - \alpha)^2}, \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}) \tag{2.26}$$

$$\left| \frac{zf'(z)}{f(z)} \left[1 + z \left(\frac{f''(z)}{f'(z)} - \frac{f'(z)}{f(z)} \right) \right] \right| < p - \alpha, \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}), \tag{2.27}$$

$$\Re \left\{ \frac{zf'(z)}{f(z)} \left(\frac{1 + \frac{zf''(z)}{f'(z)} - p}{\frac{zf'(z)}{f(z)} - p} - 1 \right) \right\} < 1, \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}), \tag{2.28}$$

then $f(z) \in \mathcal{S}_p(\alpha)$.

Corollary 2. *Let a function $f(z)$ defined by (1.1) belong to the class $\mathcal{T}(p)$. If $f(z)$ satisfies anyone of the following inequalities:*

$$\left| \frac{1 + \frac{z[2f''(z) + zf'''(z)]}{f'(z) + zf''(z)} - p}{1 + \frac{zf''(z)}{f'(z)} - p} - 1 \right| < \frac{1}{2p - \alpha}, \tag{2.29}$$

$$(z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}),$$

$$\left| z \left(\frac{2f''(z) + zf'''(z)}{f'(z) + zf''(z)} - \frac{f''(z)}{f'(z)} \right) \right| < \frac{p - \alpha}{2p - \alpha}, \quad (2.30)$$

$(z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}),$

$$\left| \frac{f'(z)}{f'(z) + zf''(z)} \left(1 + \frac{z[2f''(z) + zf'''(z)]}{f'(z) + zf''(z)} \right) - 1 \right| < \frac{p - \alpha}{(2p - \alpha)^2}, \quad (2.31)$$

$(z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}),$

$$\left| z \left(1 + \frac{zf''(z)}{f'(z)} \right) \left(\frac{2f''(z) + zf'''(z)}{f'(z) + zf''(z)} - \frac{f''(z)}{f'(z)} \right) \right| < p - \alpha, \quad (2.32)$$

$(z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}),$

$$\Re \left\{ \left(1 + \frac{zf''(z)}{f'(z)} \right) \left(\frac{1 + \frac{z[2f''(z) + zf'''(z)]}{f'(z) + zf''(z)} - p}{1 + \frac{zf''(z)}{f'(z)} - p} - 1 \right) \right\} < 1, \quad (2.33)$$

$(z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}),$

then $f(z) \in \mathcal{K}_p(\alpha)$.

Corollary 3. Let a function $f(z)$ defined by belong to the class \mathcal{T} . If $f(z)$ satisfies anyone of the following inequalities:

$$\left| \frac{\frac{zf''(z)}{zf'(z)}}{\frac{zf'(z)}{f(z)} - 1} - 1 \right| < \frac{1}{2 - \alpha}, \quad (z \in \mathcal{U}; 0 \leq \alpha < 1), \quad (2.34)$$

$$\left| 1 + z \left(\frac{f''(z)}{f'(z)} - \frac{f'(z)}{f(z)} \right) \right| < \frac{1 - \alpha}{2 - \alpha}, \quad (z \in \mathcal{U}; 0 \leq \alpha < 1), \quad (2.35)$$

$$\left| \frac{f(z)}{zf'(z)} \left(1 + \frac{zf''(z)}{f'(z)} \right) - 1 \right| < \frac{1 - \alpha}{(2 - \alpha)^2}, \quad (z \in \mathcal{U}; 0 \leq \alpha < 1), \quad (2.36)$$

$$\left| \frac{zf'(z)}{f(z)} \left[1 + z \left(\frac{f''(z)}{f'(z)} - \frac{f'(z)}{f(z)} \right) \right] \right| < 1 - \alpha, \quad (z \in \mathcal{U}; 0 \leq \alpha < 1), \quad (2.37)$$

$$\Re \left\{ \frac{zf'(z)}{f(z)} \left(\frac{\frac{zf''(z)}{f'(z)}}{\frac{zf'(z)}{f(z)} - 1} - 1 \right) \right\} < 1, \quad (z \in \mathcal{U}; 0 \leq \alpha < 1), \quad (2.38)$$

then $f(z) \in \mathcal{S}(\alpha)$.

Corollary 4. Let a function $f(z)$ defined belong to the class \mathcal{T} . If $f(z)$ satisfies anyone of the following inequalities:

$$\left| \frac{f'(z)}{zf''(z)} \left(\frac{2f''(z) + zf'''(z)}{f'(z) + zf''(z)} \right) - 1 \right| < \frac{1}{2 - \alpha}, \quad (2.39)$$

$(z \in \mathcal{U}; 0 \leq \alpha < 1),$

$$\left| z \left(\frac{2f''(z) + zf'''(z)}{f'(z) + zf''(z)} - \frac{f''(z)}{f'(z)} \right) \right| < \frac{1-\alpha}{2-\alpha}, \quad (2.40)$$

$$(z \in \mathcal{U}; 0 \leq \alpha < 1),$$

$$\left| \frac{f'(z)}{f'(z) + zf''(z)} \left(1 + \frac{z[2f''(z) + zf'''(z)]}{f'(z) + zf''(z)} \right) - 1 \right| < \frac{1-\alpha}{(2-\alpha)^2}, \quad (2.41)$$

$$(z \in \mathcal{U}; 0 \leq \alpha < 1),$$

$$\left| z \left(1 + \frac{zf''(z)}{f'(z)} \right) \left(\frac{2f''(z) + zf'''(z)}{f'(z) + zf''(z)} - \frac{f''(z)}{f'(z)} \right) \right| < 1 - \alpha, \quad (2.42)$$

$$(z \in \mathcal{U}; 0 \leq \alpha < 1),$$

$$\Re \left\{ \left(1 + \frac{zf''(z)}{f'(z)} \right) \left(\frac{f'(z)[2f''(z) + zf'''(z)]}{f''(z)[f'(z) + zf''(z)]} - 1 \right) \right\} < 1, \quad (2.43)$$

$$(z \in \mathcal{U}; 0 \leq \alpha < 1),$$

then $f(z) \in \mathcal{K}(\alpha)$.

ACKNOWLEDGEMENTS: The second-named author would like to express his thanks to the Council for Scientific and Industrial Research, Govt. of India, New Delhi for financial support.

References

- [1] P. L. Duren, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften 259, Springer-Verlag, New York, Berlin, Heidelberg and Tokyo, 1983.
- [2] A. W. Goodman, *Univalent Functions*, Vols. I and II, Polygonal Publishing House, Washington, New Jersey, 1983.
- [3] H. Irmak and Ö. F. Çetin, Some theorems involving inequalities on p -valent functions, *Turkish J. Math.* 23 (1999), 453-459.
- [4] H. Irmak and Ö. F. Çetin, Some inequalities on p -valently starlike and p -valently convex functions, *Bull. Natur. Sci. Engrg. Hacettepe Uni. Ser. B*, 28 (1999), 71-76.
- [5] H. Irmak and S. Owa, Certain inequalities for multivalent starlike and meromorphically multivalent starlike functions, *Bull. Ins. Math. Academia Sinica*, 31 (1) (2003), 11-21.
- [6] I. S. Jack, Functions starlike and convex of order α , *J. London Math. Soc.* 3 (1971), 469-474.
- [7] S. S. Miller and P. T. Mocanu, Second order differential inequalities in the complex plane, *J. Math. Anal. Appl.* 65 (1978), 289-305.
- [8] S. Owa, M. Nunokawa and H. Saitoh, Some inequalities involving multivalent functions, *Ann. Polon. Math.* 60 (2) (1994), 159-162.
- [9] S. Owa, M. Nunokawa and S. Fukui, A criteiron for p -valently starlike functions, *Internat. J. Math. and Math. Sci.* 17 (1) (1994), 205-207.

- [10] S. Owa, H. M. Srivastava, F.-Y. Ren and W.-Q. Yang, The starlikeness of a certain class of integral operators, *Complex Variables* 27 (1995), 185-1991.
- [11] H. M. Srivastava, S. Owa (Editors), *Current Topics in Analytic Function Theory*, World Scientific Publishing Company, Singapore, New Jersey, London and Hong Kong, 1992.