

## *A Criterion for the Minimal Closedness of the Lie Subalgebra Corresponding to a Connected Nonclosed Lie Subgroup*

JAN KUBARSKI

**ABSTRACT.** A Lie subalgebra  $\mathfrak{h}$  of a Lie algebra  $\mathfrak{g}$  is said to be *minimally closed* (after A. Malcev [11]) if the corresponding connected Lie subgroup is closed in the simply connected Lie group determined by  $\mathfrak{g}$ . The aim of this paper is to prove the following theorem:

*Let  $H \subset G$  be any connected (not necessarily closed) Lie subgroup of a Lie group  $G$ . Denote by  $\mathfrak{h}$ ,  $\bar{\mathfrak{h}}$  and  $\mathfrak{g}$  the Lie algebras of  $H$ , of its closure  $\bar{H}$  and of  $G$ , respectively. If there exists a Lie subalgebra  $\mathfrak{c} \subset \mathfrak{g}$  such that (a)  $\mathfrak{c} + \bar{\mathfrak{h}} = \mathfrak{g}$ , (b)  $\mathfrak{c} \cap \bar{\mathfrak{h}} = \mathfrak{h}$ , then  $\mathfrak{h}$  is minimally closed.*

As a corollary we obtain that if  $\pi_1(G)$  is finite, then no such a Lie subalgebra  $\mathfrak{c}$  exists provided that  $H$  is nonclosed.

The proof is carried out on the ground of the theory of Lie algebroids and by using some ideas from the theory of transversally complete foliations.

### 0. INTRODUCTION

**A)** Let  $G$  be any connected Lie group. Assume that  $H \subset G$  is any of its connected and nonclosed Lie subgroups. Denote by  $\mathfrak{h}$ ,  $\bar{\mathfrak{h}}$ ,  $\mathfrak{g}$  the Lie algebras of  $H$ , of its closure  $\bar{H}$  and of  $G$ , respectively.

**0.1. Problem.** *Does there exist a Lie subalgebra  $\mathfrak{c} \subset \mathfrak{g}$  such that (a)  $\mathfrak{c} + \bar{\mathfrak{h}} = \mathfrak{g}$ , (b)  $\mathfrak{c} \cap \bar{\mathfrak{h}} = \mathfrak{h}$ ?*

In work [7], some topological obstructions of the existence of such a Lie algebra  $\mathfrak{c}$  were found. Namely, the following theorem was proved.

**0.2. Theorem.** *If the following homomorphism of algebras*

$$V(\bar{\mathfrak{h}}/\mathfrak{h})^* \longrightarrow (V\bar{\mathfrak{h}}^*)_I \xrightarrow{h_p} H_{\text{dR}}(G/\bar{H}) \quad (1)$$

(where  $h_p$  is the Chern-Weil homomorphism of the  $\bar{H}$ -principal fibre bundle  $P = (G \rightarrow G/\bar{H})$ ) is nontrivial, then such a Lie subalgebra  $\mathfrak{c}$  does not exist. ■

Next, it was noticed that the case of a compact and semisimple Lie group is a case for which homomorphism (1) is always nontrivial. As a corollary we have.

**0.3. Theorem.** *If  $G$  is a compact and semisimple Lie group, then no Lie subalgebra  $\mathfrak{c}$  fulfilling (a) and (b) above exists. ■*

We add that (1) appears as the Chern-Weil homomorphism of the Lie algebroid of the TC-foliation  $\mathcal{F} = \{gH; g \in G\}$  of left cosets of  $G$  by  $H$ , determined by the author [7], [8].

**B)** In the present paper, a Lie algebroid of a connected (not necessarily closed) Lie subgroup  $H$  of a given Lie group  $G$  is constructed precisely. It can be noticed that it is the same as the one constructed in the theory of P. Molino [12] for the corresponding TC-foliation  $\mathcal{F}$  of left cosets. Next, we get to the core of the structure of this Lie algebroid and prove some strengthening of theorem 0.3 (by weakening the assumptions to the finiteness of  $\pi_1(G)$ ) without using any characteristic classes. This fact is obtained as a corollary from the theorem saying that:

**0.4. Theorem.** *The existence of a Lie subalgebra  $\mathfrak{c}$  fulfilling (a) and (b) above implies the minimal closedness of  $\mathfrak{h}$  (in the sense of Malcev [11]). ■*

## 1. PRELIMINARIES

We give a few elementary facts concerning the theory of Lie algebroids, needed in the sequel. We assume that in our paper all the manifolds considered are of  $C^\infty$ -class and Hausdorff. By  $\Omega^0(M)$  we denote the ring of  $C^\infty$  functions on a manifold  $M$ , by  $\mathfrak{X}(M)$  the Lie algebra of  $C^\infty$  vector fields on  $M$ , and by  $\text{Sec}A$  the  $\Omega^0(M)$ -module of all  $C^\infty$  global cross-sections of a given vector bundle  $A$  (over  $M$ ).

**1.1. Definition** [15], [16]. *By a transitive Lie algebroid on a manifold  $M$  we mean a system*

$$A = (A, [\![ \cdot, \cdot ]\!] , \gamma) \quad (2)$$

consisting of a vector bundle  $A$  (over  $M$ ) and mappings

$$[[\cdot, \cdot]]: SecA \times SecA \rightarrow SecA, \quad \gamma: A \rightarrow TM,$$

such that

- (i)  $(SecA, [[\cdot, \cdot]])$  is an  $\mathbf{R}$ -Lie algebra,
  - (ii)  $\gamma$ , called by K. Mackenzie [9] an *anchor*, is an epimorphism of vector bundles,
  - (iii)  $Sec\gamma: SecA \rightarrow \mathfrak{X}(M)$ ,  $\xi \rightarrow \gamma \circ \xi$ , is a homomorphism of Lie algebras,
  - (iv)  $[[\xi, f \cdot \eta]] = f \cdot [[\xi, \eta]] + (\gamma \circ \xi)(f) \cdot \eta$  for  $f \in \Omega^0(M)$ ,  $\xi, \eta \in SecA$ .
- $\mathfrak{g} := Ker\gamma$  is a vector bundle and the short exact sequence

$$0 \rightarrow \mathfrak{g} \hookrightarrow A \xrightarrow{\gamma} TM \rightarrow 0 \tag{3}$$

is called an *Atiyah sequence of (2)*; in each vector space  $\mathfrak{g}_{|x} = Ker\gamma_{|x}$ ,  $x \in M$ , some Lie algebra structure is defined by

$$[v, w]:= [[\xi, \eta]](x), \quad \xi, \eta \in SecA, \quad \xi(x) = v, \quad \eta(x) = w, \quad v, w \in \mathfrak{g}_{|x}.$$

$\mathfrak{g}_{|x}$  is called the *isotropy Lie algebra of (2) at  $x$* .  $\mathfrak{g}$  is a Lie algebra bundle [2], [5], [6], [9] called (after Mackenzie) the *adjoint of (2)*.

Let (2) and  $(A', [[\cdot, \cdot]]', \gamma')$  be two transitive Lie algebroids on the same manifold  $M$ . By a *strong homomorphism*

$$H: (A', [[\cdot, \cdot]]', \gamma') \rightarrow (A, [[\cdot, \cdot]], \gamma) \tag{4}$$

between them [4], [10, p. 273] we mean a strong homomorphism of vector bundles  $H: A' \rightarrow A$ , such that

- (i)  $\gamma \circ H = \gamma'$ ,
- (ii)  $SecH: SecA' \rightarrow SecA$ ,  $\xi \rightarrow H \circ \xi$ , is a homomorphism of Lie algebras.

If homomorphism (4) is a bijection, then  $H^{-1}$  is also a homomorphism of Lie algebroids; then  $H$  is called an *isomorphism of Lie algebroids*.

**1.2. Example.** By a *trivial Lie algebroid* [14] we mean any algebroid isomorphic to  $(TM \times \mathfrak{g}, \llbracket \cdot, \cdot \rrbracket, \rho_1)$  where  $\mathfrak{g}$  is a finitely dimensional Lie algebra and the bracket  $\llbracket \cdot, \cdot \rrbracket$  is defined by

$$\llbracket (X, \sigma), (Y, \eta) \rrbracket = ([X, Y], \mathcal{L}'_X \eta - \mathcal{L}'_Y \sigma + [\sigma, \eta]),$$

$X, Y \in \mathfrak{X}(M), \sigma, \eta: M \rightarrow \mathfrak{g}$  ( $[\sigma, \eta]$  is defined point by point:  $[\sigma, \eta](x) = [\sigma(x), \eta(x)], x \in M$ ).

**1.3. Example** (See [5], [6], [9]). By the *Lie algebroid  $A(P)$  of a principal fibre bundle  $P = (P, \pi, M, G, \cdot)$*  we mean a transitive Lie algebroid on  $M$   $(A(P), \llbracket \cdot, \cdot \rrbracket, \gamma)$  in which  $A(P) = TP/G, \gamma([v]) = \pi_*(v)$  where  $[v]$  denotes the equivalence class of  $v$ , and the bracket  $\llbracket \xi, \eta \rrbracket, \xi, \eta \in \text{Sec}A(P)$ , is constructed on the basis of the following observation: For each cross-section  $\eta \in \text{Sec}A(P)$ , there exists exactly one  $C^\infty$  right-invariant vector field  $\eta' \in \mathfrak{X}^R(P)$  such that  $[\eta'(z)] = \eta(\pi z)$ , and the mapping  $\text{Sec}A(P) \rightarrow \mathfrak{X}^R(P), \eta \rightarrow \eta'$ , is an isomorphism of  $\Omega^0(M)$ -modules. The bracket  $\llbracket \xi, \eta \rrbracket$  is a cross-section of  $A(P)$  such that  $\llbracket \xi, \eta \rrbracket' = [\xi', \eta']$ .

The Lie algebroid of a trivial principal fibre bundle  $P = M \times G$  is canonically isomorphic to the trivial Lie algebroid  $A = TM \times \mathfrak{g}$ ,  $\mathfrak{g}$  is the right Lie algebra of  $G$ , via

$$A(P) = T(M \times G)/G = TM \times (TG/G) \ni (v, [w]) \rightarrow (v, \Theta^R(w)) \in TM \times \mathfrak{g};$$

$\Theta^R$  denotes the canonical right-invariant 1-form on  $G$  [5], [6].

A transitive Lie algebroid strongly isomorphic to  $A(P)$  for some principal fibre bundle is called *integrable* [9]. There exist non-integrable Lie algebroids discovered by R. Almeida and P. Molino [1]. Lie algebroids of some TC-foliations are non-integrable, for example, the Lie algebroid of the foliation of left cosets of any connected and simply connected Lie group by a connected nonclosed Lie subgroup has this property.

**1.4. Definition.** By a *connection in transitive Lie algebroid* (2), see [5], [9], [15], we mean a homomorphism of vector bundles  $\lambda: TM \rightarrow A$  such that  $\gamma \circ \lambda = id_{TM}$ , i.e. a splitting of Atiyah sequence (3) of  $A$

$$0 \rightarrow \mathfrak{g} \hookrightarrow A \xrightarrow[\lambda]{\gamma} TM \rightarrow 0.$$

By a *curvature tensor of a connection  $\lambda$*  in (2) we shall mean a tensor  $\Omega_b \in \Omega^2(M; \mathfrak{g}) (= \text{Sec} \Lambda^2 T^* M \otimes \mathfrak{g})$  defined by

$$\Omega_b(X, Y) = \lambda[X, Y] - \llbracket \lambda X, \lambda Y \rrbracket, X, Y \in \mathfrak{X}(M).$$

$\lambda$  also determines a covariant derivative  $\nabla$  in  $\mathfrak{g}$  by

$$\nabla_X \sigma = [[\lambda X, \sigma]], \quad X \in \mathfrak{X}(M), \quad \sigma \in \text{Sec } \mathfrak{g},$$

See [5], [9].

It turns out that the Lie algebra structure in  $\text{Sec } A$  is uniquely determined by  $\mathfrak{g}$ ,  $\nabla$ ,  $\Omega_h$  and  $\lambda$ , namely, we have

**1.5. Theorem** [5], [9]. *The mapping  $\varphi: TM \oplus \mathfrak{g} \rightarrow A, (v, w) \rightarrow \lambda v + w$ , is an isomorphism of Lie algebroids provided that in  $TM \oplus \mathfrak{g}$  the following Lie algebroid structure is defined:*

(a) *the bracket:*

$$[[ (X, \sigma), (Y, \eta) ]] = ([X, Y], -\Omega_h(X, Y) + \nabla_X \eta - \nabla_Y \sigma + [\sigma, \eta]), \quad X, Y \in \mathfrak{X}(M), \\ \sigma, \eta \in \text{Sec } \mathfrak{g} \text{ (} [\sigma, \eta] \text{ is defined point by point: } [\sigma, \eta](x) = [\sigma(x), \eta(x)], x \in M).$$

(b) *the anchor:  $\gamma = pr_1: TM \oplus \mathfrak{g} \rightarrow TM$ . ■*

## 2. THE LIE ALGEBROID OF A CONNECTED (NOT NECESSARILY CLOSED) LIE SUBGROUP

Let  $G$  be any connected Lie group and  $H \subset G$  any connected (not necessarily closed) Lie subgroup of  $G$ .  $H$  determines the foliation  $\mathcal{F} = \{gH: g \in G\}$  of left cosets of  $G$  by  $H$ .  $\mathcal{F}$  is a transversally complete foliation [12], [13] because right-invariant vector fields are from the normalizer of  $\mathfrak{X}(\mathcal{F})$  and generate the entire tangent space  $T_g G$  for any  $g \in G$ .

Denote by  $E$  the tangent bundle to  $\mathcal{F}$  and  $Q = TG/E \xrightarrow{r} G$  the transversal bundle of  $\mathcal{F}$ . Let

$$\alpha: TG \rightarrow Q$$

be the canonical projection and let  $\bar{v}, v \in TG$ , denote the vector  $\alpha(v)$ .  $R_t: TG \rightarrow TG$  stands for the differential of the right translation by  $t \in G$ .

**2.1. Lemma.** (i)  $R_t, t \in \bar{H}$  ( $\bar{H}$  is the closure of  $H$ ), maps  $E$  into  $E$  inducing the isomorphism of vector bundles  $\bar{R}_t: Q \rightarrow Q, \bar{v} \mapsto \bar{R}_t(\bar{v})$ .

(ii) *The mapping  $\bar{R}: Q \times \bar{H} \rightarrow Q, (\bar{v}, t) \mapsto \bar{R}_t(\bar{v})$ , is a right strongly free action.*

**Proof.** Easy calculations. ■

As a corollary we obtain

**2.2.** The topological space  $A(G; H)$  of orbits of the action  $\bar{R}$ , i.e.

$$A(G; H) = Q/\approx \quad \text{where} \quad \bar{v} \approx \bar{w} \iff \exists t \in \bar{H} \quad (\bar{R}_t(\bar{v}) = \bar{w})$$

has a uniquely determined structure of a  $C^\infty$  manifold, such that the canonical projection  $\beta: Q \rightarrow A(G; H)$  is a submersion.

In the sequel, the vector  $\beta(\bar{v})$ ,  $\bar{v} \in Q$ , will be denoted by  $[\bar{v}]$  and  $\pi_b: G \rightarrow G/\bar{H}$  stands for the canonical projection. Of course,  $\bar{r}: A(G; H) \rightarrow G/\bar{H}$ ,  $[\bar{v}] \mapsto \pi_b(r\bar{v})$ , is a correctly defined projection. Its smoothness follows immediately from the commutativity of the diagram

$$\begin{array}{ccc} Q & \xrightarrow{\beta} & A(G; H) \\ \downarrow r & & \downarrow \bar{r} \\ G & \xrightarrow{\pi_b} & G/\bar{H} \end{array}$$

For the fibre  $A(G; H)_{|\bar{g}}$  of  $\bar{r}$  over  $\bar{g} \in G/\bar{H}$ , the mapping  $\beta_g: Q_{|g} \rightarrow A(G; H)_{|\bar{g}}$ ,  $g \in \pi_b^{-1}(\bar{g})$ , is a bijection. Via  $\beta_g$  we introduce in  $A(G; H)_{|\bar{g}}$  some structure of a real vector space and, clearly, it is independent of the choice of  $g$ . We wish to arrange the system  $(A(G; H), \bar{r}, G/\bar{H})$  to be a vector bundle. For the purpose, we find local trivializations of this system.

**2.3. Definition.** A  $C^\infty$  cross-section  $\zeta \in \text{Sec } Q$  is called a transversal field if, for any  $g \in G$  and  $t \in \bar{H}$ ,

$$\zeta(gt) = \bar{R}_t(\zeta(g))$$

(that is, if  $\zeta$  is  $\bar{H}$ -right-invariant).

**2.4. Example.** The  $C^\infty$  cross-section  $\bar{Y}_w := \alpha \circ Y_w$  where  $Y_w$  stands for the right-invariant vector field on  $G$  generated by  $w \in \mathfrak{g}$  ( $\mathfrak{g}$  is the Lie algebra of  $G$ ) is a transversal field. Therefore, transversal fields generate the entire space  $Q_{|g}$  for any  $g \in G$ .

**2.5. Remarks.** Denote by  $l(G; H)$  the space of all transversal fields.

(a)  $l(G; H)$  forms a module over the ring  $\Omega^0(G/\bar{H})$  under the multiplication  $\bar{f} \cdot \zeta := \bar{f} \circ \pi_b \cdot \zeta$ ,  $\bar{f} \in \Omega^0(G/\bar{H})$ ,  $\zeta \in l(G; H)$ .

(b) If transversal fields  $\zeta_1, \dots, \zeta_s$  are linearly independent at a point  $g \in G$ , then, immediately by the definition, they are linearly independent at each point  $gt, t \in \bar{H}$ , and, in consequence, at some open  $\mathcal{F}_b$ -saturated open subset where  $\mathcal{F}_b$  is the so-called *basic foliation*  $\mathcal{F}_b = \{g\bar{H}; g \in G\}$ .

(c) Let  $\zeta_i \in l(G; H), i \leq s$ . If  $\zeta_i$  are linearly independent on  $U = \pi_b^{-1}[\bar{U}]$  ( $\bar{U}$  open in  $G/\bar{H}$ ) and  $\zeta = \sum_{i=1}^s f^i \zeta_i$  for  $f^i \in \Omega^0(U)$ , then the functions  $f^i$  are of the form  $f^i = \tilde{f}^i \circ \pi_b|_U$  for some  $\tilde{f}^i \in \Omega^0(\bar{U})$ .

**2.6. Proposition.** Let  $q = \dim G - \dim H$ , i.e.  $q = \text{codim } \mathcal{F}$ . Suppose that  $\zeta_1, \dots, \zeta_q$  are transversal fields linearly independent at each point of a set  $U = \pi_b^{-1}[\bar{U}]$ ,  $\bar{U}$  open in  $G/\bar{H}$ . Then

$$\begin{aligned} \varphi: \bar{U} \times \mathbf{R}^q &\longrightarrow \bar{r}^{-1}[\bar{U}] \subset A(G; H) \\ (\bar{g}, \alpha) &\longrightarrow [\sum \alpha^i \zeta_i(g)], g \in \pi_b^{-1}(\bar{g}), \end{aligned}$$

is a local trivialization of  $\bar{r}: A(G; H) \longrightarrow G/\bar{H}$ .

**Proof.** Of course,  $\varphi_{\bar{g}}: \mathbf{R}^q \longrightarrow A(G; H)|_{\bar{g}}, \bar{g} \in \bar{U}$ , is an isomorphism of vector spaces. This proposition will be proved by showing that  $\varphi$  is a diffeomorphism. For the purpose, take the mapping  $\psi: U \times \mathbf{R}^q \longrightarrow r^{-1}[U] \subset Q, (g, \alpha) \longrightarrow \sum \alpha^i \zeta_i(g)$ , being a local trivialization of  $Q$ . Our assertion follows now from the commutativity of the diagram

$$\begin{array}{ccc} U \times \mathbf{R}^q & \xrightarrow{\psi} & r^{-1}[U] \subset Q & \xrightarrow{r} & G \\ \downarrow \pi_b \times id & & \downarrow \beta & & \downarrow \\ \bar{U} \times \mathbf{R}^q & \xrightarrow{\varphi} & \bar{r}^{-1}[\bar{U}] \subset A(G; H) & \xrightarrow{\bar{r}} & G/\bar{H} . \blacksquare \end{array}$$

**2.7. Remark.** The structure of a  $C^\infty$  manifold in  $A(G; H)$  can be obtained independently by demanding that  $\varphi$ 's be diffeomorphisms.

Now, we introduce a structure of a Lie algebroid into the vector bundle  $A(G; H)$ . Firstly, we define the anchor  $\gamma: A(G; H) \longrightarrow T(G/\bar{H})$  by  $[\bar{w}] \longrightarrow \pi_{b*}(w)$  (the correctness is easy to obtain). Secondly, we introduce in  $\text{Sec } A(G; H)$  a structure of a Lie algebra in the way described below.

Take a homomorphism of  $\Omega^0(G/\bar{H})$ -modules

$$c: l(G; H) \rightarrow \text{Sec } A(G; H), \quad \zeta \rightarrow c_\zeta, \quad (5)$$

where  $c_\zeta$  is a  $C^\infty$  cross-section of  $A(G; H)$  defined by  $c_\zeta(\bar{g}) = [\zeta(g)]$ ,  $g \in \pi_h^{-1}(\bar{g})$ .

**2.8. Lemma.** *c is an isomorphism of  $\Omega^0(G/\bar{H})$ -modules.*

**Proof.** We check at once that (5) is a monomorphism. To see that it is also an epimorphism, take an arbitrary  $C^\infty$  cross-section  $\xi \in \text{Sec } A(G; H)$  and define a cross-section  $\zeta$  of  $Q$  in such a way that the diagram

$$\begin{array}{ccc} & \beta & \\ & Q \rightarrow A(G; H) & \\ \zeta \uparrow & & \uparrow \xi \\ & G \xrightarrow{\pi_h} G/\bar{H} & \end{array}$$

commutes, i.e.  $c_\zeta = \xi$ . The smoothness of  $\zeta$  is the last thing to notice. In order to get this, take transversal fields  $\zeta_1, \dots, \zeta_q$  being a basis on  $U = \pi_h^{-1}(\bar{U})$  ( $\bar{U}$  is open in  $G/\bar{H}$  and contains an arbitrarily taken point of  $G/\bar{H}$ ). Then  $c_{\zeta_1}, \dots, c_{\zeta_q}$  forms a basis of  $A(G; H)$  on  $\bar{U}$ . Therefore,  $\xi = \sum \bar{f}^i c_{\zeta_i}$  on  $\bar{U}$  for some  $\bar{f}^i \in \Omega^0(\bar{U})$ . Of course,  $\zeta = \sum \bar{f}^i \circ \pi_h \cdot \zeta_i$  on  $U$ , which ends the proof. ■

**2.9.** The space  $l(G; H)$  has a natural structure of a real Lie algebra. Indeed, let  $\zeta, \nu \in l(G; H) \subset \text{Sec } Q$ . Take arbitrary vector fields  $X, Y \in \mathfrak{X}(G)$  such that  $\zeta = \bar{X}$  ( $:= \alpha \circ X$ ) and, analogously,  $\nu = \bar{Y}$ . Put

$$[\zeta, \nu] := \overline{[X, Y]}. \quad (6)$$

We need notice that

(a)  $\overline{[X, Y]} \in l(G; H)$ ,

(b) definition (6) is correct.

Let us first observe



**2.10. Lemma.** If  $\zeta \in l(G; H)$  is of the form  $\zeta = \bar{X}$  for a vector field  $X \in \mathfrak{X}(G)$ , then  $X$  belongs to the normalizer of  $\mathfrak{X}(\mathcal{F})$ , that is,

$$[X, Y] \in \mathfrak{X}(\mathcal{F}) \text{ for all } Y \in \mathfrak{X}(\mathcal{F}) \quad (7)$$

[i.e.  $X$  is the so-called foliate vector field for  $\mathcal{F}$ , see [13]].

**Proof.** Of course, it is sufficient to show relation (7) for left-invariant vector fields  $Y = X_h$ ,  $h \in \mathfrak{h}$ , only. To this end, take an arbitrary  $g \in G$  and express  $\zeta$  locally on a set  $U = \pi_b^{-1}[\bar{U}]$  containing  $g$  ( $\bar{U}$  open in  $G/\bar{H}$ ), in the form  $\zeta|_U = \sum \bar{f}^i \circ \pi_{h|U} \cdot Y_{w_i|U}$ ,  $\bar{f}^i \in \Omega(\bar{U})$ ,  $w_i \in \mathfrak{g}$  (for  $Y_w$ , see 2.4). Then

$$Z := \sum \bar{f}^i \circ \pi_{h|U} \cdot Y_{w_i|U} - X|_U \in \mathfrak{X}(\mathcal{F}_U)$$

and, furthermore, we have

$$\begin{aligned} [X, X_h]|_U &= [\sum \bar{f}^i \circ \pi_{h|U} \cdot Y_{w_i|U} - Z, X_h|_U] \\ &= -[Z, X_h]|_U \in \mathfrak{X}(\mathcal{F}_U), \end{aligned}$$

thus  $[X, X_h] \in \mathfrak{X}(\mathcal{F})$ . ■

**2.11. Remark.** It can be proved that condition (7) is equivalent to the fact that  $\zeta := \bar{X}$  is a transversal field; however, the sufficiency of this condition will not be used in the sequel.

Now, we are able to prove (a) and (b) from 2.9.

(a): To get the equality  $\bar{R}_t(\overline{[X, Y]}(g)) = \overline{[X, Y]}(gt)$ ,  $g \in G$ ,  $t \in \bar{H}$ , take the vector fields  $Z_1 = R_t X - X$  and  $Z_2 = R_t Y - Y$  tangent to  $\mathcal{F}$ . Applying 2.10, we deduce that

$$\begin{aligned} \bar{R}_t(\overline{[X, Y]}(g)) &= \overline{R_t(\overline{[X, Y]}(g))} = \overline{R_t(\overline{[X, Y]}(gt))} \\ &= \overline{[R_t X, R_t Y]}(gt) = \overline{[X + Z_1, Y + Z_2]}(gt) \\ &= \overline{[X, Y]}(gt). \end{aligned}$$

(b): Immediately from 2.10.

**2.12.** In  $\text{Sec}A(G; H)$  we introduce the bracket  $[[\cdot, \cdot]]$  (forming a Lie algebra) by demanding that (5) be an isomorphism of Lie algebras, i.e.  $[[c_\zeta, c_v]] := c_{[\zeta, v]}$ ,  $\zeta, v \in l(G; H)$ .

**2.13. Theorem.** *The system*

$$A(G; H) = (A(G; H), \llbracket \cdot, \cdot \rrbracket, \gamma) \quad (8)$$

*is a transitive Lie algebroid on  $G/\bar{H}$ .*

**Proof.** (1)  $\text{Sec}\gamma: \text{Sec}A(G; H) \rightarrow \mathfrak{H}(G/\bar{H})$  is a homomorphism of Lie algebras. To see this, take  $\xi, \eta \in \text{Sec}A(G; H)$ . Find vector fields  $X, Y \in \mathfrak{X}(G)$  such that  $\xi = c_{\bar{X}}, \eta = c_{\bar{Y}}$ . By the definition of  $\gamma$ ,

$$(\text{Sec}\gamma)(c_{\bar{X}})(\bar{g}) = \pi_{b*}g(X_g) \text{ for } \bar{g} = \pi_b(g), g \in G,$$

from which we obtain that  $X$  is  $\pi_b$ -related to  $\gamma \circ \xi$  and, analogously,  $Y$  to  $\gamma \circ \eta$ . Therefore  $[X, Y]$  is  $\pi_b$ -related to  $[\gamma \circ \xi, \gamma \circ \eta]$  and to  $\gamma \circ \llbracket \xi, \eta \rrbracket$  simultaneously, which confirms our assertion.

(2) The equality  $\llbracket \xi, \bar{f} \cdot \eta \rrbracket = \bar{f} \cdot \llbracket \xi, \eta \rrbracket + (\gamma \circ \xi)(\bar{f}) \cdot \eta$ ,  $\bar{f} \in \Omega^0(G/\bar{H})$ ,  $\xi, \eta \in \text{Sec}A(G; H)$ , follows easily from

$$[\bar{X}, \bar{f} \circ \pi_b \cdot \bar{Y}] = \bar{f} \circ \pi_b \cdot [\bar{X}, \bar{Y}] + (\gamma \circ c_{\bar{X}})(\bar{f}) \cdot \bar{Y}. \blacksquare$$

Lie algebroid (8) will be called the *Lie algebroid of a Lie subgroup  $H$  of  $G$* . It can be interesting only in the case of a nonclosed  $H$  because the closedness of  $H$  implies the triviality of  $A(G; H)$ :  $A(G; H) \cong T(G/\bar{H})$ .

**2.14. Remark.** One can prove [cf. [7]] that Lie algebroid (8) is equal to the one constructed by P. Molino [12], [13] for the TC-foliation  $\mathcal{F}$ .

### 3. STRUCTURE THEOREMS

Let (8) be the Lie algebroid of a connected Lie subgroup  $H$  of a connected Lie group  $G$  and

$$0 \rightarrow \mathfrak{g} \hookrightarrow A(G; H) \xrightarrow{\gamma} T(G/\bar{H}) \rightarrow 0$$

its Atiyah sequence. In this section we prove three fundamental facts concerning  $A(G; H)$ :

- *The adjoint Lie algebra bundle  $\mathfrak{g}$  of  $A(G; H)$  is a trivial bundle of abelian Lie algebras.*
- *If the Lie algebroid  $A(G; H)$  admits a flat connection (i.e. a connection with the zero curvature tensor), then it is trivial.*

• Let  $\mathfrak{h}, \bar{\mathfrak{h}}, \mathfrak{g}$  denote the Lie algebras of  $H, \bar{H}$  and  $G$ , respectively. Suppose that there exists a Lie subalgebra  $\mathfrak{c} \subset \mathfrak{g}$  such that (a)  $\mathfrak{c} + \bar{\mathfrak{h}} = \mathfrak{g}$ , (b)  $\mathfrak{c} \cap \bar{\mathfrak{h}} = \mathfrak{h}$ . Then  $A(G; H)$  admits a flat connection.

The crucial role in the proving of the first fact is played by the following Malcev theorem (for a short “foliated” proof of it, see [7]).

**3.1. The Malcev Theorem** [11], [17]. *If  $H$  is a dense connected Lie subgroup of a Lie group  $T$ , then  $H$  is a normal subgroup of  $T$  and  $T/H$  is abelian. ■*

By this, according to our notations,  $\mathfrak{h}$  is an ideal of  $\bar{\mathfrak{h}}$  and  $\bar{\mathfrak{h}}/\mathfrak{h}$  is an abelian Lie algebra.

**3.2. Theorem.** *For a vector  $w \in \bar{\mathfrak{h}}$ , the cross-section  $\bar{X}_w$  of the transversal bundle  $Q$ , induced by the left-invariant vector field  $X_w$ , is a transversal field, and the mapping*

$$\varphi: G/\bar{H} \times \bar{\mathfrak{h}}/\mathfrak{h} \rightarrow \mathfrak{g}, (\bar{g}, [w]) \rightarrow [\bar{X}_w(g)], g \in \pi_h^{-1}(\bar{g}), \quad (9)$$

is a global trivialization of the Lie algebra bundle  $\mathfrak{g}$ .

**Proof.** It is sufficient to show that  $\bar{X}_w, w \in \bar{\mathfrak{h}}$ , is a transversal field; the rest is easy. Clearly, for  $t \in \bar{H}$  and  $g \in G$   $\bar{R}_t(\bar{X}_w(g)) = \bar{L}_g(\bar{R}_t(w))$  and  $\bar{X}_w(gt) = \bar{L}_g(\bar{L}_t(w))$  where  $\bar{L}_g: Q \rightarrow Q$  is an automorphism of the vector bundle  $Q$ , determined by the differential  $L_g$  of the left-translation by  $g$ . Therefore, it remains to prove that  $R_t(w) - L_t(w) \in E_t$ , which means that the vector field  $X := Y_w - X_w$  is tangent to the foliation  $\mathcal{F}$  at each point of  $\bar{H}$ .

Firstly, we notice that  $X$  is foliate; to see this, we calculate: Let  $h \in \mathfrak{h}$ , then  $[X, X_h] = [Y_w - X_w, X_h] = X_{[h, w]} \in \mathfrak{X}(\mathcal{F})$  because  $[h, w] \in \mathfrak{h}$  according to the Malcev theorem.

Secondly, any foliate vector field  $X$  (for a foliation  $\mathcal{F}$ ) in any distinguished local coordinates  $x = (x^1, \dots, x^p, y^1, \dots, y^q)$  ( $p = \dim \mathcal{F}$ ,  $q = \text{codim } \mathcal{F}$ ) is of the form  $X(x, y) = \sum a^i(x, y) \frac{\partial}{\partial x^i} + \sum b^j(y) \frac{\partial}{\partial y^j}$  [13]; therefore, which is easy to see, if it is tangent to  $\mathcal{F}$  at a point  $z$ , then it is tangent to  $\mathcal{F}$  at each point of the closure of the leaf through  $z$ . In our situation,  $X(e) = [Y_w - X_w](e) = O \in E_{|e}$ , so - by the above - our theorem is proved. ■

Now, we proceed to the second problem.

**3.3. Theorem.** *If the Lie algebroid  $A(G; H)$  is flat, then it is trivial.*

**Proof.** Let  $\lambda: T(G/\bar{H}) \rightarrow A(G; H)$  be a flat connection in  $A(G; H)$ . Then, taking account of 1.4 and isomorphism (9) of Lie algebra bundles, we have an isomorphism of Lie algebroids

$$\rho: T(G/\bar{H}) \times \bar{\mathfrak{h}}/\mathfrak{h} \rightarrow A(G; H), (v, [w]) \rightarrow \lambda v + [\bar{X}_w(g)],$$

$v \in T_{\bar{g}}G/\bar{H}$ ,  $\bar{g} = \pi_b(g)$ ,  $g \in G$ , provided that in  $T(G/\bar{H}) \times \bar{\mathfrak{h}}/\mathfrak{h}$  the Lie algebroid structure is defined by the following formula

$$[[X, \sigma], (Y, \eta)] = ([X, Y], \nabla_X^0 \eta - \nabla_Y^0 \sigma + [\sigma, \eta]),$$

$X, Y \in \mathfrak{X}(G/\bar{H})$ ,  $\sigma, \eta: G/\bar{H} \rightarrow \bar{\mathfrak{h}}/\mathfrak{h}$ , where  $\nabla^0$  is a covariant derivative in the trivial vector bundle  $T(G/\bar{H}) \times \bar{\mathfrak{h}}/\mathfrak{h}$ , such that  $\varphi$  maps  $\nabla^0$  onto  $\nabla$ , i.e.

$$\nabla_X^0 \sigma = \varphi^{-1} \circ \nabla_X (\varphi \circ \sigma) = \varphi^{-1} \circ [[\lambda X, \varphi \circ \sigma]].$$

Looking at example 1.2, we see that to end the proof, it is sufficient to show the equality

$$\nabla_X^0 = \lambda X, \quad X \in \mathfrak{X}(G/\bar{H}),$$

which is equivalent to the fact that the covariant derivative  $\nabla_X^0$  of any constant function  $\bar{w}: G/\bar{H} \rightarrow \bar{\mathfrak{h}}/\mathfrak{h}$ ,  $\bar{g} \rightarrow [w]$ ,  $w \in \bar{\mathfrak{h}}$ , is zero, i.e. that  $[[\lambda X, c_{\bar{X}_w}]] = 0$ . The cross-section  $\lambda X$  is locally of the form  $\lambda X = \sum \bar{f}^i c_{\bar{Y}_{w_i}}$ ,  $\bar{f}^i \in \Omega^0(G/\bar{H})$ , thus

$$\begin{aligned} [[\lambda X, c_{\bar{X}_w}]] &= [[\sum \bar{f}^i c_{\bar{Y}_{w_i}}, c_{\bar{X}_w}]] \\ &= \sum \bar{f}^i [[c_{\bar{Y}_{w_i}}, c_{\bar{X}_w}]] - \gamma^0 c_{\bar{X}_w} (\bar{f}^i) \cdot c_{\bar{Y}_{w_i}} \\ &= 0 \end{aligned}$$

because  $\gamma^0 c_{\bar{X}_w} = 0$  and  $[[c_{\bar{Y}_{w_i}}, c_{\bar{X}_w}]] = c_{\{\bar{Y}_{w_i}, \bar{X}_w\}} = 0$ . ■

It remains to consider the third problem.

**3.4. Theorem.** *Suppose that there exists a Lie subalgebra  $\mathfrak{c} \subset \mathfrak{g}$  such that (a)  $\mathfrak{c} + \bar{\mathfrak{h}} = \mathfrak{g}$ , (b)  $\mathfrak{c} \cap \bar{\mathfrak{h}} = \mathfrak{h}$ . Then  $A(G; H)$  admits a flat connection.*

*Proof.* The construction of a flat connection in  $A(G; H)$  has four steps.

**Step 1.** Denote by  $\bar{C} \subset TG$  the left-invariant distribution generated by  $\mathfrak{c}$ , i.e. the vector bundle tangent to the foliation  $\{gF; g \in G\}$  where  $F$  is the connected Lie subgroup with the Lie algebra equalling  $\mathfrak{c}$ .  $\bar{C}$  fulfils the following conditions (in which  $E_h$  is the vector bundle tangent to the foliation  $\mathcal{F}_h = \{g\bar{H}; g \in G\}$ ):

- (1)  $\bar{C} + E_h = TG$ ,
- (2)  $\bar{C} \cap E_h = E$ ,
- (3)  $\bar{C}$  is  $\bar{H}$ -right-invariant [i.e.  $\bar{C}_{|g^t} = R_t[\bar{C}_{|g}]$ ,  $g \in G$ ,  $t \in \bar{H}$ ],
- (4)  $\bar{C}$  is involutive.

Clearly, (1), (2) and (4) hold. To see (3), take an arbitrary vector  $v \in \bar{C}_{|g}$ , we have  $v = L_x(w)$  for some  $w \in \mathfrak{c}$ . Since  $R_t(v) = L_x(R_t(w))$ , we need only to observe that  $R_t(w) \in \bar{C}_{|t}$  for  $t \in \bar{H}$ . Write  $t = \lim t_n$ ,  $t_n \in \bar{H}$ ; then, by the closedness of  $\bar{C}$  in  $TG$ , we obtain that  $R_t(w) = \lim R_{t_n}(w) \in \bar{C}$  because  $R_{t_n}[\bar{C}] = \bar{C}$ .

**Step 2.** Let  $\bar{C} \subset TG$  be a distribution realizing conditions (1) ÷ (4) above. Via the epimorphism  $\alpha: TG \rightarrow Q$  we define a subbundle  $C \subset Q$  by  $C_{|g} = \alpha_g[\bar{C}_{|g}]$ ,  $g \in G$ . [The fact that  $C$  is a subbundle is obtained from the relation  $E \subset \bar{C}$  which holds by (2)].  $C$  fulfils the following conditions:

- (1')  $Q' \oplus C = Q$  where  $Q' = E_h / E \subset Q$ ,
- (2')  $C$  is  $\bar{H}$ -right-invariant [i.e.  $C_{|g^t} = \bar{R}_t[C_{|g}]$ ,  $g \in G$ ,  $t \in \bar{H}$ ],
- (3')  $l_i(G; H) := \text{Sec } C \cap l(G; H)$  is a Lie subalgebra of  $l(G; H)$ .

(1') and (2') are obvious. To check (3'), take arbitrary  $\zeta, \nu \in l_i(G; H)$  and write  $\zeta = \bar{X}$ ,  $\nu = \bar{Y}$  for some vector fields  $X, Y \in \mathfrak{X}(\bar{C})$ . According to (4),  $[X, Y] \in \mathfrak{X}(\bar{C})$ , which gives the relation  $[\bar{X}, \bar{Y}] \in \text{Sec } C$ . On the other hand (see 2.9),  $[\zeta, \nu] = [\bar{X}, \bar{Y}] \in l(G; H)$ .

**Step 3.** Let  $C \subset Q$  be any vector subbundle realizing conditions (1') ÷ (3') above. Via the linear homomorphism  $\beta: Q \rightarrow A(G; H)$  we define a subbundle  $C \subset A(G; H)$  by  $C_{|\bar{g}} = \beta_g[C_{|g}]$ ,  $g \in \pi_h^{-1}(\bar{g})$ ,  $\bar{g} \in G/\bar{H}$ . Thanks to the equality  $\beta \circ \bar{R}_t = \beta$ ,  $t \in \bar{H}$ , the correctness of this definition is evident. To see that  $C$  is a  $C^\infty$  vector subbundle of  $A(G; H)$ , it is sufficient to notice that a local  $C^\infty$  cross-section of  $A(G; H)$  lying in  $C$  and passing through an arbitrarily taken vector from  $C$  exists. Let  $\bar{v} \in C_{|\bar{g}}$  and  $\bar{g} = \pi_h(g)$ . Take a local

$C^\infty$  cross-section  $\varphi: U \rightarrow G$  of the submersion  $\pi_h: G \rightarrow G/\bar{H}$ , such that  $\varphi(\bar{g}) = g$ , and consider the diagram

$$\begin{array}{ccccc}
 i^*C' & \xrightarrow{\bar{i}} & C \subset Q & \xrightarrow{\beta} & A(G; H) \\
 \downarrow & & \downarrow & & \downarrow \\
 \text{Im}\varphi & \xrightarrow{i} & G & \xrightarrow{\pi_h} & G/\bar{H} \\
 \uparrow & & \uparrow & & \uparrow \\
 & \xrightarrow{\varphi} & & & U
 \end{array}$$

Diminishing  $U$  if necessary, we may assume that the vector bundle  $i^*C' \rightarrow \text{Im}\varphi$  has a global  $C^\infty$  cross-section  $Z$  passing through  $\bar{v}$ . Put  $\xi = \beta \circ \bar{i} \circ Z \circ \varphi: U \rightarrow A(G; H)$ ;  $\xi$  is, of course, a  $C^\infty$  cross-section of  $A(G; H)$  over  $U$  such that  $\xi(\bar{g}) = [\bar{v}]$ . The vector bundle  $C$  fulfils the conditions

- (1)  $\mathfrak{g} \oplus C = A(G; H)$ ,
- (2)  $\text{Sec}C$  is a Lie subalgebra of  $\text{Sec}A(G; H)$ .

(1) is evident by the observation that  $\beta_g$  maps isomorphically  $Q|_g$  onto  $\mathfrak{g}|_{\bar{g}}$ . To see (2), take arbitrary  $\xi, \eta \in \text{Sec}C$ . According to 2.8, there exist transversal fields  $\zeta, v$  such that  $c_\zeta = \xi$  and  $c_v = \eta$ . Of course,  $\beta_g(\zeta_g) = \xi_{\bar{g}}$  and  $\beta_g(v_g) = \eta_{\bar{g}}$ ,  $g \in \pi_h^{-1}(\bar{g})$ . From the definition of  $C$  we obtain that  $\zeta$  and  $v$  belong to  $l_i(G; H)$ . By (3'),  $[\zeta, v] \in \text{Sec}C' \cap l(G; H)$ , therefore  $[[\xi, \eta]] = c_{[\zeta, v]} \in \text{Sec}C$ .

**Step 4.** Let  $C \subset A(G; H)$  be a vector subbundle realizing conditions (1) and (2) above. Then, of course, a splitting  $\lambda$  of the Atiyah sequence of  $A(G; H)$ , see the diagram

$$0 \rightarrow \mathfrak{g} \xrightarrow{\lambda} A(G; H) = \mathfrak{g} \oplus C \xrightarrow{\gamma} T(G/\bar{H}) \rightarrow 0$$

such that  $\text{Im}\lambda = C$ , is a flat connection in  $A(G; H)$ . ■

Combining the above theorems we get

**3.5. Corollary.** *The existence of a Lie subalgebra  $\mathfrak{c} \subset \mathfrak{g}$  fulfilling  $\mathfrak{c} + \bar{\mathfrak{h}} = \mathfrak{g}$  and  $\mathfrak{c} \cap \bar{\mathfrak{h}} = \mathfrak{h}$  implies the triviality of the Lie algebroid  $A(G; H)$ .*

#### 4. MAIN RESULTS

Let the symbols  $H, \bar{H}, G, \mathfrak{h}, \bar{\mathfrak{h}}, \mathfrak{g}$  have the same meaning as in the previous two sections.

**4.1. Theorem.** *If there is a Lie subalgebra  $\mathfrak{c} \subset \mathfrak{g}$  such that (a)  $\mathfrak{c} + \bar{\mathfrak{h}} = \mathfrak{g}$ , (b)  $\mathfrak{c} \cap \bar{\mathfrak{h}} = \mathfrak{h}$ , then the Lie algebra  $\mathfrak{h}$  is minimally closed.*

**Proof.** Corollary 3.5 states that the Lie algebroid  $A(G; H)$  of the Lie subgroup  $H \subset G$  is trivial, i.e. there exists a Lie algebroid isomorphism  $\Phi: A(G; H) \rightarrow A_o := T(G/\bar{H}) \times \bar{\mathfrak{h}}/\mathfrak{h}$ . Such a Lie algebroid is, of course, integrable:  $A_o$  is the Lie algebroid of the trivial principal fibre bundle  $P = G/\bar{H} \times F$  for an arbitrarily taken Lie group  $F$  with the abelian Lie algebra  $\bar{\mathfrak{h}}/\mathfrak{h}$ , see 1.3. The following reasoning is due to R. Almeida and P. Molino, see the proof of their theorem [13, p. 138]. Consider the Lie algebroid  $(TG \times \bar{\mathfrak{h}}/\mathfrak{h}, [\![\cdot, \cdot]\!] , pr_1)$  of the trivial principal fibre bundle  $G \times F$ . The linear homomorphism of vector bundles

$$\lambda: TG \rightarrow TG \times \bar{\mathfrak{h}}/\mathfrak{h}, \quad v \rightarrow (v, pr_2 \circ \Phi([\bar{v}]))$$

is a connection in this Lie algebroid.  $\lambda$  is flat. Indeed, it is sufficient to show the equality  $[\![\lambda X, \lambda Y]\!] = \lambda[X, Y]$  only for  $X, Y \in \mathfrak{X}(G)$  such that the corresponding cross-sections  $\bar{X}, \bar{Y}$  of  $Q$  are transversal fields. However, the equality is then easy to obtain by using the fact that  $\Phi$  is a homomorphism of Lie algebras, namely, writing  $\lambda X = (X, pr_2 \circ \Phi \circ c_{\bar{X}} \circ \pi_b)$  (and, analogously, for  $\lambda Y$ ), we have

$$\begin{aligned} [\![\lambda X, \lambda Y]\!] &= [\![ (X, pr_2 \circ \Phi \circ c_{\bar{X}} \circ \pi_b), (Y, pr_2 \circ \Phi \circ c_{\bar{Y}} \circ \pi_b) ]\!] \\ &= ([X, Y], \mathcal{L}_X(pr_2 \circ \Phi \circ c_{\bar{Y}} \circ \pi_b) - \mathcal{L}_Y(pr_2 \circ \Phi \circ c_{\bar{X}} \circ \pi_b) \\ &\quad + [pr_2 \circ \Phi \circ c_{\bar{X}} \circ \pi_b, pr_2 \circ \Phi \circ c_{\bar{Y}} \circ \pi_b]) \\ &= ([X, Y], \mathcal{L}_{\gamma \circ c_{\bar{X}}}(pr_2 \circ \Phi \circ c_{\bar{Y}}) \circ \pi_b - \mathcal{L}_{\gamma \circ c_{\bar{Y}}}(pr_2 \circ \Phi \circ c_{\bar{X}}) \circ \pi_b \\ &\quad + [pr_2 \circ \Phi \circ c_{\bar{X}}, pr_2 \circ \Phi \circ c_{\bar{Y}}] \circ \pi_b) \\ &= ([X, Y], pr_2 \circ [\!(\gamma \circ c_{\bar{X}}, pr_2 \circ \Phi \circ c_{\bar{Y}}), (\gamma \circ c_{\bar{Y}}, pr_2 \circ \Phi \circ c_{\bar{X}})\!] \circ \pi_b) \\ &= ([X, Y], pr_2 \circ [\![\Phi \circ c_{\bar{X}}, \Phi \circ c_{\bar{Y}}]\!] \circ \pi_b) \\ &= ([X, Y], pr_2 \circ \Phi \circ [\![c_{\bar{X}}, c_{\bar{Y}}]\!] \circ \pi_b) \\ &= ([X, Y], pr_2 \circ \Phi \circ c_{[\bar{X}, \bar{Y}]} \circ \pi_b) \\ &= \lambda[X, Y]. \end{aligned}$$

Let  $D$  be the connection in  $G \times F$  determined by  $\lambda$ , i.e. the right-invariant distribution  $D \subset T(G \times F)$  for which

$$D_{|(g, e)} = \{(v, pr_2 \circ \Phi([\bar{v}]); v \in TG\}, \quad g \in G,$$

where  $e$  denotes the neutral element of  $F$ . The flatness of  $\lambda$  implies the involutivity of  $D$ . Consider the diagram

$$\begin{array}{ccc} G \times F & \xrightarrow{p_2 = \pi_{\bar{h}} \times id} & G/\bar{H} \times F \\ \downarrow p_1 & & \downarrow \\ G & \xrightarrow{\pi_{\bar{h}}} & G/\bar{H}. \end{array}$$

Let  $\tilde{G} \subset G \times F$  be any leaf of the distribution  $D$ . Of course,  $\tilde{p}_1 = p_1|_{\tilde{G}}: \tilde{G} \rightarrow G$  is a covering and, which is easy to obtain,  $\tilde{p}_2 = p_2|_{\tilde{G}}: \tilde{G} \rightarrow G/\bar{H} \times F$  is a submersion. Denote by  $\tilde{\mathcal{F}}$  the lifting (by  $\tilde{p}_1$ ) of the foliation  $\mathcal{F}$  in  $\tilde{G}$ . Let  $(g, a) \in \tilde{G}$ . For  $v \in T_g G$ , the following conditions are equivalent:

- (1)  $v$  is tangent to  $\mathcal{F}$ ,
- (2)  $\tilde{p}_{1*(g, a)}^{-1}(v)$  is tangent to  $\tilde{\mathcal{F}}$ ,
- (3)  $\tilde{p}_{2*(g, a)}(\tilde{p}_{1*(g, a)}^{-1}(v)) = 0$ .

From this we obtain that  $\tilde{\mathcal{F}}$  is defined by the submersion  $\tilde{p}_2: \tilde{G} \rightarrow G/\bar{H} \times F$ , in particular, the leaves of  $\tilde{\mathcal{F}}$  are closed. Introducing in  $\tilde{G}$  a structure of a group in the standard way we obtain:  $\tilde{G}$  is a Lie group and  $\tilde{p}_1$  is a local isomorphism of Lie groups. It is a standard calculation to obtain that  $\tilde{\mathcal{F}}$  is then the foliation of left cosets of  $\tilde{G}$  by  $\tilde{F}$  where  $\tilde{F}$  is a connected Lie subgroup of  $\tilde{G}$  with the Lie algebra equalling  $\tilde{\mathfrak{h}} = \tilde{p}_{1*\tilde{e}}^{-1}[\mathfrak{h}]$  ( $\tilde{e}$  being the neutral element of  $\tilde{G}$ ). Therefore  $\tilde{F}$  is a closed Lie subgroup. Of course,  $F$ , being closed after the lifting to some covering, is also closed after lifting it to the universal one, which means that  $\mathfrak{h}$  is minimally closed. ■

**4.2. Theorem.** *If  $\pi_1(G)$  is finite and  $H \neq \bar{H}$ , then there exists no Lie subalgebra  $\mathfrak{c} \subset \mathfrak{g}$  fulfilling the conditions  $\mathfrak{c} + \bar{\mathfrak{h}} = \mathfrak{g}$  and  $\mathfrak{c} \cap \bar{\mathfrak{h}} = \mathfrak{h}$ .*

**Proof.** Let  $\pi_1(G)$  be finite. Then, the universal covering is finite, which implies the nonclosedness of the lifting  $\tilde{H}$  of  $H$ . Our assertion follows now trivially from the previous theorem. ■

To finish with, we can ask



Can Theorem 4.1 be inverted ?

It turns out that the answer is no.

**4.3. Example.** Let  $G = U(2)$ . Suppose that  $\bar{H} = T$  is a maximal torus in  $G$ . It is well known that  $\dim T = 2$  and the lifting of  $T$  to the universal covering  $\mathbf{R} \times SU(2) \rightarrow U(2)$  is isomorphic to the cylinder  $\mathbf{R} \times S^1$ . Therefore, any Lie subalgebra  $\mathfrak{h}$  of  $\bar{\mathfrak{h}}$  ( $\bar{\mathfrak{h}}$  — the Lie algebra of  $\bar{H}$ ) is minimally closed. We prove, using theorem 0.2, that there exists some 1-dimensional Lie subalgebra  $\mathfrak{h}$  of  $\bar{\mathfrak{h}}$  for which.

- (i) no Lie subalgebra  $\mathfrak{c} \subset \mathfrak{g}$  fulfilling (a) and (b) from 0.1 exists.
- (ii) the corresponding connected Lie subgroup of  $T$  is dense in  $T$ .

Let  $h_p: V(\bar{\mathfrak{h}}^*) \rightarrow H_{dR}^2(G/T)$  be the Chern-Weil homomorphism of the  $T$ -principal fibre bundle  $P = (G \rightarrow G/T)$ .  $G$  and  $T$  have the same rank, therefore, according to [3; Th.VII, p. 467], we have that

$$h_p^{(2)}: \bar{\mathfrak{h}}^* \rightarrow H_{dR}^2(G/T)$$

is surjective. Moreover,  $\dim \bar{\mathfrak{h}}^* = 2$  and  $\dim H_{dR}^2(G/T) = 1$ , thus  $\dim \text{Ker } h_p^{(2)} = 1$ . Then it is obvious that there exists a covector  $O \neq \beta \in \bar{\mathfrak{h}}^*$  such that (1)  $h_p^{(2)}(\beta) \neq 0$ , (2)  $\mathfrak{h} := \text{Ker } \beta \subset \bar{\mathfrak{h}}$  is a subspace such that the corresponding Lie subgroup  $H \subset T$  is dense in  $T$ . Of course, the superposition

$$(\bar{\mathfrak{h}}/\mathfrak{h})^* \xrightarrow{j} \bar{\mathfrak{h}}^* \xrightarrow{h_p^{(2)}} H_{dR}^2(G/T)$$

is nontrivial:  $h_p^{(2)} \circ j(\bar{\beta}) \neq 0$  where  $\bar{\beta} \in (\bar{\mathfrak{h}}/\mathfrak{h})^*$  is a linear homomorphism determined by  $\beta$ . Theorem 0.2 implies the nonexistence of a Lie subalgebra  $\mathfrak{c} \subset \mathfrak{g}$  fulfilling (a) and (b) above.

### References

- [1] R. ALMEIDA, P. MOLINO: Suites d'Atiyah et feuilletages transversalement complets, *C. R. Acad. Sci. Paris Ser. I Math.*, 300 (1985), 13-15.
- [2] — Suites d'Atiyah, feuilletages et quantification geometrique, *Estrait du Seminaire du Geometrie differentielle*, Montpellier, 1984-85.
- [3] W. GREUB, S. HALPERIN, R. VANSTONE: *Connections, Curvature, and Cohomology, Vol. III*, Academic Press, New York and London, 1976.
- [4] J. KUBARSKI: Exponential mapping for Lie groupoids, *Colloq. Math.* XLVII (1982), 267-282.

- [5] — Lie algebroid of a principal fibre bundle, *Publications du Department de Mathematiques de L'Universite de LYON I*, in printing.
- [6] — Lie algebroid of a principal fibre bundle - three equivalent definitions, *Prace Naukowe Politechniki Szczecinskiej*, 11 (1988), 123-145.
- [7] — The Chern-Weil homomorphism of regular Lie algebroids and TC-foliations, *Preprint No 9*, Institute of Mathematics, Technical University of Łódź, November 1988.
- [8] — Pontryagin algebra of a transitive Lie algebroid, *Supplemento ai Rendiconti del Circolo Matematico di Palermo, Proceedings of the Winter School on Geometry and Physics*, SRN1, 9-16 January, 1988 (in printing).
- [9] K. MACKENZIE: *Lie groupoids and Lie algebroids in differential Geometry*, Cambridge, 1987.
- [10] — A note on Lie algebroids which arise from groupoid actions, *Cahiers de Topologie et Geometrie Differentielle Categoriques*, XXVIII-4 (1987), 283-302.
- [11] A. MAL'CEV: Subgroups of Lie groups in the large, *C. R. (Doklady) Acad. Sci. URSS (N. S.)* 36 (1942), 5-7 (Kraus Reprint LTD. VADUZ 1963).
- [12] P. MOI'INO: Etude des feuilletages transversalement complets et applications, *Ann. Sci. École Norm. Sup.*, 10(3) (1977), 289-307.
- [13] — Riemannian Foliations, *Progress in Mathematics Vol. 73*, Birkhäuser Boston Basel, 1988.
- [14] NGO-VAN-QUE: Sur l'espace de prolongement differentiable, *J. Differential Geom.* 2.1. (1968), 33-40.
- [15] J. PRADINES: Theorie de Lie pour les groupoides differentiables dans la categorie des groupoides, Calcul differential dans la categorie des groupoides infinitesimaux, *C. R. Acad. Sci. Ser. A-B*, Paris, 264 (1967), 265-248.
- [16] — : Theorie de Lie pour les groupoides differentiables, *Atti Conv Intern Geom 7 Diff.* Bologna, 1967, Bologna-Amsterdam.
- [17] E. WINBERG, A. ONISHCHIK: *A seminar on Lie groups and algebraic groups* (in Russian), Moscow, 1988.

Institute of Mathematics,  
 Technical University of Lodz,  
 Poland

Recibido: 28 de marzo de 1990