

## *Monotonicity in Time and Stationary Solutions for a Quasilinear Heat Equation with Source*

VICTOR A. GALAKTIONOV and SERGEY A. POSASHKOV

**ABSTRACT.** We consider the Cauchy problem for the quasilinear parabolic heat equation with source  $u_t = \Delta u^{\sigma+1} + u^\beta$  in  $R^N \times (0, T)$ ,  $\sigma > 0, \beta > 1$  are fixed constants, with nonnegative bounded symmetric initial function. Two properties of monotone behaviours of the solution  $u(|x|, t)$  for  $x=0$  are investigated. 1. Monotonicity of large solutions: there exists a constant  $M_k > 0$  such that if  $u(0, t_0) \geq M_k$  for some  $t_0 \in [0, T)$ , then  $u_t(0, t) \geq 0$  for all  $t \in [t_0, T)$ . 2.  $u(0, t)$  does not decrease in  $(0, T)$ . It is shown that sufficient conditions for these properties are quite different for five cases: i)  $1 < \beta < \sigma + 1$ , ii)  $\beta = \sigma + 1$ , iii)  $\sigma + 1 < \beta < \beta_*$ , iv)  $\beta = \beta_*$ , v)  $\beta > \beta_*$ , where  $\beta_* = (\sigma + 1)(N + 2)/(N - 2)$  for  $N > 2$  ( $\beta_* = \infty$  for  $N = 1, 2$ ) is the critical Sobolev exponent.

### 1. INTRODUCTION. MAIN RESULTS

In this paper we consider the Cauchy problem for the quasilinear parabolic heat equation with source

$$u_t = \Delta u^{\sigma+1} + u^\beta \quad \text{in } R^N \times (0, T) \quad (1)$$

$$u(x, 0) = u_0(x) \geq 0 \quad \text{in } R^N, \quad (2)$$

where  $\sigma > 0$  and  $\beta > 1$  are fixed constants. Equation (1) is well-known equation arising in the theory and in different applications of the processes of the heat conduction and combustion in a medium, where the heat conductivity coefficient  $k(u) \equiv (\sigma + 1)u^\sigma$  and the power of the energy emission  $Q(u) \equiv u^\beta$  depend upon the temperature of the medium  $u = u(x, t) \geq 0$ .

We shall assume the initial function (2) satisfies the following hypotheses:

$$\begin{aligned} u_0 = u_0(r) \geq 0 \quad \text{in } R^N, r = |x|; M_1 = \sup u_0 < \infty; \\ u_0(r) \text{ is the continuous function in } R_+^1 = (0, \infty), \\ (u_0^\sigma u_0)(0) = 0. \end{aligned} \quad (3)$$

Under these hypotheses, there exists the unique weak local (in time) solution  $u = u(r, t)$  of the problem (1), (2), which is nonnegative continuous function in  $R^N \times (0, T)$ , where  $T \in (0, \infty]$  is a finite or an infinite *existence time*. See a full list of references given in [11]. Notice that  $u(r, t)$  is the classical solution at any point  $(r, t)$  where  $u(r, t)$  is strictly positive.

The main results of the paper are devoted to the analysis of the behaviour with time of the temperature at the single point  $x=0$ . This yields the conditions on the initial temperature for the *ignition* of the combustion process at the single point  $x=0$ .

We shall consider two types of the monotone behaviours of  $u(0, t)$ .

**Property (ML) (Monotonicity of the Large solution):** there exists a constant  $M_k > 0$ , depending on the initial function, such that if  $u(0, t_0) \geq M_k$  for some  $t_0 \in [0, T)$ , then

$$u_t(0, t) \geq 0 \quad \text{for all } t \in [t_0, T). \quad (4)$$

**Property (M) (Monotonicity for arbitrary  $t$ ):**

$$u(0, t) \text{ doesn't decrease in } (0, T). \quad (5)$$

We show that above properties (ML) and (M) depend on the initial function  $u_0$ , the dimension of the space  $N \geq 1$  and exponents  $\sigma, \beta$  of equation (1). These properties are quite different for five cases: i)  $1 < \beta < \sigma + 1$ , ii)  $\beta = \sigma + 1$ , iii)  $\sigma + 1 < \beta < \beta_*$ , iv)  $\beta = \beta_*$ , v)  $\beta > \beta_*$ , where

$$\beta_* = (\sigma + 1)(N + 2)/(N - 2) \quad \text{for } N > 2 \quad (\beta_* = \infty \quad \text{for } N = 1, 2) \quad (6)$$

is the critical Sobolev exponent for the nonlinear elliptic operator in the right-hand side of (1).

Let  $B_\varepsilon = \{r < \varepsilon\}$  be the ball in  $R^N$  of a radius  $\varepsilon > 0$  with the boundary  $S_\varepsilon = \{r = \varepsilon\}$ .  $D_\varepsilon = R^N \setminus B_\varepsilon$  and denote

$$h^*(r) = c_* r^{-(N-2) \cdot 2(\sigma+1)} \quad \text{for } r > 0.$$

$$\begin{aligned}
 c_* &= \left[ \frac{N(N-2)}{4} \right]^{(N-2)/4(\sigma+1)} \quad (N > 2); \\
 h(r) &= c r^{-2/[\beta-(\sigma+1)]} \quad \text{for } r > 0, \\
 c &= \left[ \frac{2N(\sigma+1)}{\beta-(\sigma+1)} \right]^{1/[\beta-(\sigma+1)]} \quad (\beta > \sigma+1); \\
 h_\infty(r) &= c_\infty r^{-2/[\beta-(\sigma+1)]} \quad \text{for } r > 0, \\
 c_\infty &= \left\{ \frac{2(\sigma+1)[\beta(N-2) - N(\sigma+1)]}{[\beta-(\sigma+1)]^2} \right\}^{-1/[\beta-(\sigma+1)]}, \\
 & \quad (\beta > (\sigma+1)N/(N-2) \quad \text{for } N > 2).
 \end{aligned}
 \tag{7}$$

We now state the main results of the paper.

**Theorem 1** ( $1 < \beta < \sigma + 1$ ). *Let  $1 < \beta < \sigma + 1$  and let (3) holds. Then*

(i) *If  $u_0(r)$  satisfies*

$$\begin{aligned}
 & \text{there exists a large constant } R > 0 \text{ such that } u_0^{\sigma+1} \text{ is} \\
 & \text{uniformly Lipschitz continuous in } D_R,
 \end{aligned}
 \tag{8}$$

*then (ML) holds.*

(ii) *If  $u_0^{\sigma+1}(r) \in C^1$  and*

$$\left| [u_0^{\sigma+1}(r)]' \right| = o(r^{(\beta+\sigma+1)/(\sigma+1-\beta)}) \quad \text{as } r \rightarrow +\infty,
 \tag{9}$$

*then (ML) holds.*

(iii) *If  $u_0^{\sigma+1}(r) \in C^1$  and*

$$(u_0^{\sigma+1}(r))' + u_0^\beta(r) r/N > 0 \quad \text{in } \{r > 0\} \cap \{u_0 > 0\},
 \tag{10}$$

*then (M) holds.*

**Theorem 2** ( $\beta = \sigma + 1$ ). *Let  $\beta = \sigma + 1$  and let (3) holds. Then*

(i) *If  $u_0(r)$  satisfies*

there exists small enough  $\varepsilon > 0$  such that

$$u_0^{\sigma+1} \text{ is Lipschitz continuous in } \{z_N - \varepsilon < r < z_N\}, \quad (11)$$

where  $z_N > 0$  is the first zero of Bessel function  $J_{(N-2)/2}$ , then (ML) holds.

(ii) If  $u_0^{\sigma+1}(r) \in C^1$  and

$$(u_0^{\sigma+1}(r))' + u_0^{\sigma+1}(r) r/N > 0 \text{ in } \{0 < r < z_N\} \cap \{u_0 > 0\}, \quad (12)$$

then (M) holds.

(iii) If  $u_0(r)$  is the nondecreasing function in  $\{0 < r < z_N\}$ , then (M) holds.

**Theorem 3** ( $\sigma + 1 < \beta < \beta_*$ ). Let  $\sigma + 1 < \beta < \beta_*$  and let (3) holds. Then

(i) If for some small  $\varepsilon > 0$

$$u_0^{\sigma+1}(r) \text{ is Lipschitz continuous in } B_\varepsilon, \quad (13)$$

then (ML) holds.

(ii) If  $u_0^{\sigma+1}(r) \in C^1$  and

$$(u_0^{\sigma+1}(r))' + u_0^\beta(r) r/N > 0 \text{ in } \{r > 0\} \cap \{u_0 > 0\}, \quad (14)$$

then (M) holds.

(iii) If  $u_0(r)$  is the nondecreasing function in  $\{r < l_0\}$ , where

$$l_0 = c_l (u_0(0))^{-1/\beta - (\sigma+1)/2} \quad (15)$$

and  $c_l = c_l(\sigma, \beta, N)$  is some positive constant, then (M) holds.

(iv) If  $u_0(r)$  is the nondecreasing function in  $\{r > 0 \mid u_0(r) < h(r)\}$ , then (M) holds.

(v) Let  $l_* = \sup \{a > 0 \mid u_0(r) \text{ is the nondecreasing function for } r \in (0, a)\} > 0$ . Then (ML) holds for

$$M_{k_N} = (c_l/l_*)^{2/(\beta - (\sigma+1))}.$$

**Theorem 4** ( $\beta = \beta_*$ ). Let  $\beta = \beta_*$  for  $N \geq 3$  and let (3) holds. Then

(i) If  $u_0(r)$  satisfies (13) for some  $\varepsilon > 0$  and

$$u_0(r) > 0 \text{ for all } r > 0, \tag{16}$$

there exists constant  $m_0 > 0$  such that

$$u_0(r) > m_0 \cdot r^{-(N-2)/(\alpha+1)} \text{ for any large } r > 0, \tag{16'}$$

then (ML) holds.

(ii) Assertion (ii) of Theorem 3 is valid.

(iii) If  $u_0(r)$  is the nondecreasing function in  $\{r > 0 \mid u_0(r) < h^*(r)\}$ , then (M) holds.

**Theorem 5** ( $\beta > \beta_*$ ). Let  $\beta > \beta_*$  for  $N \geq 3$  and let (3) holds. Then

(i) Let  $u_0(r) = h_\infty(r)$  for a unique point  $r = r_* > 0$ ,  $u_0 \in C^1$  in a small neighbourhood of  $r = r_*$  and  $u'_0(r_*) > h'_\infty(r_*)$ . Let for any sufficiently small  $\varepsilon > 0$  the set  $\{r > 0 \mid |u_0(r)/h_\infty(r) - 1| < \varepsilon\}$  be a finite connected interval containing the point  $r = r_*$ . Then (ML) holds.

(ii) Assertion (ii) of Theorem 3 is valid.

(iii) If  $u_0(r)$  is the nondecreasing function in  $\{r > 0 \mid u_0(r) < h(r)\}$ , then (M) holds.

Proofs of Theorems 1-5 are based on intersection comparison of the solution  $u(r, t)$  with the set of the stationary solutions  $\{U\}$ . Therefore, we begin with the analysis of the set  $\{U\}$ .

## 2. SOME PROPERTIES OF THE SET OF THE STATIONARY SOLUTIONS

We now describe some well-known properties of the stationary solutions  $U(r; \lambda)$  satisfying

$$\begin{aligned} \Delta U^{\alpha+1} + U^\beta &\equiv r^{1-N} (r^{N-1} (U^{\alpha+1})_r)_r + \dot{U}^\beta = 0 \text{ for } r > 0, \\ U(0; \lambda) &= \lambda > 0, \\ U_r(0; \lambda) &= 0, \end{aligned} \tag{17}$$

where  $\lambda > 0$  is an arbitrary fixed constant.

One can see that equation (17) is invariant under the self-similar transformation and hence

$$U(r; \lambda) = \lambda U(r\lambda^m; 1), \quad m = [\beta - (\sigma + 1)]/2, \quad \text{for any fixed } \lambda > 0. \quad (18)$$

It is well-known (see e.g. [10]) that for  $\beta \geq \beta_*$  any solution of (17) is strictly positive for all  $r > 0$  and  $U(\infty; \lambda) = 0$ . For  $1 < \beta < \beta_*$  the function  $U(r; \lambda)$  vanishes at some point  $r = l(\lambda)$ , where

$$l(\lambda) = c_1 \cdot \lambda^{-m}, \quad (19)$$

$c_1(\sigma, \beta, N) \equiv l(1) > 0$  is given in (15). In this case we let  $U(r; \lambda) \equiv 0$  for  $r > l(\lambda)$ . For  $\beta \geq \beta_*$  we formally let  $l(\lambda) = +\infty$ .

Thus, for  $\beta > 1$  we have introduced one parametric set of the stationary solutions  $\{U(r; \lambda); \lambda \geq 0\}$  ( $U(r; 0) \equiv 0$ ). In all cases for arbitrary fixed  $\lambda > 0$   $U(r; \lambda)$  is the monotone decreasing function in the domain of positivity. The function  $U(r; \lambda)$  is the continuous function with respect to  $r$  and  $\lambda$ . From well-known properties of solution of the Cauchy problem for the ordinary differential equation (17) we get that there exists the continuous dependence on  $\lambda$  of the solution  $U(r; \lambda)$  in any compact from  $R_+^1$  and the derivative  $(U^{\sigma+1})_r$  in any compact from the domain  $\omega(\lambda) = [0, l(\lambda))$  of positivity of the stationary solution.

*The case  $\beta \in (1, \sigma + 1)$ .* For  $\beta \in (1, \sigma + 1)$  from (19) it follows that

$$l(\lambda) \rightarrow +\infty \text{ monotone as } \lambda \rightarrow +\infty. \quad (20)$$

Identity (18) yields the following conditions

$$U(r; \lambda) \rightarrow +\infty \text{ uniformly in } [0, l/2] \text{ as } \lambda \rightarrow +\infty, \quad (21)$$

$$(U^{\sigma+1})_r(r; \lambda) \rightarrow -\infty \text{ uniformly in } [l/2, l] \text{ as } \lambda \rightarrow +\infty. \quad (22)$$

Introduce for a fixed constant  $M > 0$  the set  $g(\lambda, M) = \{r > 0 \mid 0 < U(r; \lambda) < M\}$ . From (18), (19) one can obtain the following estimates: if  $r \in g(\lambda, M)$  then

$$r = l(\lambda) (1 + o(1)) \rightarrow \infty \text{ as } \lambda \rightarrow \infty, \quad (22')$$

$$|(U^{\sigma+1})_r(r; \lambda)| = k_1 r^{-(\beta + \sigma + 1)2m} (1 + o(1)) \rightarrow \infty \text{ as } \lambda \rightarrow \infty,$$

where  $k_1 = k_1(\sigma, \beta, N) > 0$  is some constant.

*The case  $\beta = \sigma + 1$ .* In this case equation (17) has the explicit solution

$$U(r; \lambda) \equiv U_S(r; \lambda) = \lambda \{\Gamma(N/2) 2^{\nu} r^{-\nu} J_{\nu}(r)\}, \quad r \in [0, z_N), \quad (23)$$

where  $\nu = (N-2)/2$ ,  $\Gamma(N/2)$  is Euler's Gamma function,  $z_N$  is the first zero of Bessel function  $J_\nu(r)$  ( $U(z_N; \lambda) = 0$ ).

For any fixed arbitrary small  $\varepsilon > 0$

$$U_S \rightarrow +\infty \text{ uniformly in } r \in [0, z_N - \varepsilon] \text{ as } \lambda \rightarrow +\infty, \tag{24}$$

$$(U_S^{\sigma+1})_r \rightarrow -\infty \text{ uniformly in } r \in [z_N - \varepsilon, z_N] \text{ as } \lambda \rightarrow +\infty. \tag{25}$$

The case  $\beta \in (\sigma + 1, \beta_*)$ . Then it follows from (19) that

$$I(\lambda) \rightarrow 0 \text{ monotone as } \lambda \rightarrow +\infty. \tag{26}$$

and properties (21), (22) are also valid.

The case  $\beta = \beta_*$  ( $N \geq 3$ ). It is well-known [12] that for  $\beta = \beta_*$ ,  $N \geq 3$ , equation (17) admits the explicit strictly positive solution

$$U(r; \lambda) = \lambda \cdot \left[ \frac{N(N-2)}{N(N-2) + \lambda^{4(\sigma+1)} (N-2) r^2} \right]^{(N-2) 2(\sigma+1)} \tag{27}$$

for  $r > 0$ . It has the following properties:

$$U(r; \lambda) = c_2 \lambda^{-1} r^{-(N-2)(\sigma+1)} (1 + o(1)) \text{ as } r \rightarrow \infty, \tag{28}$$

where  $c_2 = [N(N-2)]^{(N-2) 2(\sigma+1)}$ ;

$U(r; \lambda) \rightarrow 0$  uniformly in any set  $[\delta, +\infty)$  as  $\lambda \rightarrow \infty$ ,

$$(U^{\sigma+1})_r(r; \lambda) \rightarrow -\infty \text{ uniformly in any set} \tag{29}$$

$$\{r > 0 \mid m_1 \leq U(r; \lambda) \leq m_2\} \text{ as } \lambda \rightarrow \infty,$$

where  $\delta, m_1 < m_2$  are arbitrary fixed positive constants.

The case  $\beta > \beta_*$ ,  $N \geq 3$ . Then  $U(r; \lambda) > 0$  for any  $r > 0$ ,  $U(+\infty; \lambda) = 0$  and there hold [10]

$$U(r; \lambda) = h_\infty(r) (1 + o(1)), \tag{30}$$

$$U_r(r; \lambda) = h'_\infty(r) (1 + o(1)) \text{ as } r \rightarrow +\infty,$$

where the function  $h_\infty(r)$  is given by (7). From (18), (30) one can easily verify that for any fixed  $\delta > 0$

$$U(r; \lambda) / h_\infty(r) \rightarrow 1. \tag{31}$$

$U_r(r; \lambda)/h'_\infty(r) \rightarrow 1$  uniformly in any set  $[\delta, +\infty)$  as  $\lambda \rightarrow \infty$ .

*The estimates of  $U(r; \lambda)$ .* The first estimate holds for arbitrary  $\beta > 1$ . By using monotonicity of the function  $U(r; \lambda)$  for  $r > 0$  we obtain the following inequality

$$r^{1-N}(r^{N-1}(U^{\sigma+1})_r)_r = -U^\beta \geq -\lambda^\beta \text{ for } r > 0.$$

Integrating twice this inequality yields the lower estimate

$$U(r; \lambda) \geq U_-(r; \lambda) \equiv \lambda(1 - r^2/r_0^2)_+^{1,(\sigma+1)} \text{ for } r > 0, \quad (32)$$

where  $r_0 = r_0(\lambda) \equiv (2N)^{1/2} \cdot \lambda^{-[\beta - (\sigma+1)]/2}$ .

By integrating equation (17) and by using the monotonicity of the function  $U(r; \lambda)$  for  $r > 0$  we get the following inequality:

$$(U^{\sigma+1})_r + U^\beta \frac{r}{N} \leq 0 \text{ for } r > 0. \quad (33)$$

The second estimate for  $\beta \in (\sigma+1, +\infty)$  we shall derive by integrating (33). Then we have for  $r > 0$

$$U(r; \lambda) \leq \left[ \lambda^{-[\beta - (\sigma+1)]} + \frac{r^2[\beta - (\sigma+1)]}{2N(\sigma+1)} \right]^{-1/[\beta - (\sigma+1)]},$$

and hence

$$U(r; \lambda) \leq h(r), r > 0, \text{ for any fixed } \lambda > 0, \quad (34)$$

where  $h(r)$  is given in (7).

*The tangent curve.* Upper estimate (34) implies that for any  $\beta > \sigma+1$  there exists the tangent curve  $L = L(r)$  of the set  $\{U(r; \lambda)\}$  and the following upper estimate

$$L(r) \equiv \sup_{\lambda > 0} U(r; \lambda) \leq h(r) \text{ for } r > 0 \quad (35)$$

holds. Moreover, from (32) for  $\beta > \sigma+1$  we obtain the lower estimate of  $L(r)$ :

$$L(r) \geq L_-(r) \equiv \sup_{\lambda > 0} U_-(r; \lambda) = c_3 r^{-2[\beta - (\sigma+1)]} \text{ for } r > 0, \quad (36)$$

where

$$c_3 \equiv \left[ \frac{\beta - (\sigma+1)}{\beta} \right]^{(\sigma+1)} \left[ \frac{2N(\sigma+1)}{\beta} \right]^{[\beta - (\sigma+1)]}$$



For  $\beta = \beta_*$ , the tangent curve can be calculated explicitly:

$$L(r) = h^*(r) \text{ for } r > 0,$$

where the function  $h^*(r)$  is determined by (7).

### 3. INTERSECTION COMPARISON

1. Let  $\omega(\lambda) = [0, l(\lambda))$  be the domain of positivity of the function  $U(r; \lambda)$  for fixed  $\lambda > 0$  ( $\omega(\lambda) \equiv [0, +\infty)$  for  $\beta \geq \beta_*$ ). Let  $N(t; \lambda)$  for fixed  $t \in [0, T)$  be the number of intersections of the functions  $u(r, t)$  and  $U(r; \lambda)$  in  $\omega(\lambda)$  (see e.g. [6]–[8], [15]) or, which is the same, the number of sign changes of the difference  $u(r, t) - U(r; \lambda)$  [1], [14], [16]. This implies that the functions  $u(r, t)$  and  $U(r; \lambda)$  are positive in a small neighbourhood of any intersection. Since the solutions  $u(r, t)$  and  $U(r; \lambda)$  are classical there, each intersection is an isolated point for any fixed  $t > 0$  [1], [13], [16]. Without loss of generality we shall assume that for  $t = 0$  and for all  $\lambda > 0$  there exist only points of intersection, and  $N(0; \lambda) < +\infty$ .

The following Lemma 1 is well-known for classical solutions, see [1], [6], [7], [13]–[16]; for weak solutions of degenerate equations see results in [8], [15].

**Lemma 1.** Fix arbitrary  $\lambda > 0$ . Let (3) holds and  $N(0; \lambda) < \infty$ . Then  $N(t; \lambda) \leq N(0; \lambda) + 1$ . If  $N(0; \lambda) \leq 1$  for some fixed  $\lambda > u_0(0)$ , then

$$N(t; \lambda) \leq 1 \text{ for any } t \in (0, T). \tag{37}$$

**Proof.** Below we use the standard technique of construction of the weak solution of the Cauchy problem (1), (2) [11]. Fix an arbitrary small  $\varepsilon > 0$  and denote

$$u_{0\varepsilon}(x) \equiv \max \{u_0(r), \varepsilon\} > 0 \text{ for } r \geq 0. \tag{38}$$

Clearly,  $u_{0\varepsilon}(x) \rightarrow u_0(r)$  as  $\varepsilon \rightarrow 0$  uniformly in  $R^N$ . The Cauchy problem for (1) with initial function (38) has the unique classical strictly positive solution  $u_\varepsilon(r, t) \geq \varepsilon$  in  $R^N \times (0, T)$  [3]. Moreover (see [11] and references therein)

$$u_\varepsilon(r, t) \rightarrow u(r, t) \text{ as } \varepsilon \rightarrow 0 \tag{39}$$

uniformly in any compact set from  $R^N \times (0, T)$ .

Let  $N_\varepsilon(t; \lambda)$  for fixed  $t \in [0, T)$  be the number of the points of intersection of the solutions  $u_\varepsilon(r, t)$  and  $U(r; \lambda)$  in  $\omega(\lambda)$ . One can see that  $N_\varepsilon(0; \lambda) \leq N(0; \lambda) + 1$  (one additional point of intersection «given» in the right-hand side of the last inequality can arise on the boundary  $r = l(\lambda)$  of  $\omega(\lambda)$ ). Consider for arbitrary fixed  $t \in (0, T)$  the domain  $\Omega(t) = \omega(\lambda) \times (0, t)$ . It is well-known that  $N_\varepsilon(t; \lambda)$  is not greater than the number of the sign changes of the difference  $w_\varepsilon(r, t; \lambda) \equiv u_\varepsilon(r, t) - U(r; \lambda)$  on the parabolic boundary of  $\Omega(t)$  [1], [13], [14], [16]. Since  $u_\varepsilon(l(\lambda), t) > 0$  for all  $t \in (0, T)$  we get the inequality

$$N_\varepsilon(t; \lambda) \leq N_\varepsilon(0; \lambda) \quad \text{for } t \in (0, T).$$

From (39) by using the continuity of  $u(r, t)$  we can conclude that

$$N(t; \lambda) \leq N_\varepsilon(t; \lambda) \quad \text{for any small } \varepsilon > 0,$$

and hence  $N(t; \lambda) \leq N(0; \lambda) + 1$ .

If  $\lambda > u_0(0)$  and  $N(0; \lambda) = 0$  or  $1$ , then it is easily seen that  $N_\varepsilon(0; \lambda) \leq 1$  for any small  $\varepsilon > 0$ . Then in both cases we have  $N(t; \lambda) \leq 1$  for all  $t \in (0, T)$ , which completes the proof.

**2. Comparison with the set of the stationary solutions.** The following two Lemmas 2, 3 are based on the analysis of the number of intersection of the initial function  $u_0(r)$  with the set  $\{U(r; \lambda)\}$  of the stationary solutions.

These Lemmas yield the sufficient conditions for the properties (ML) and (M).

**Lemma 2** (property (ML)). *Let (3) holds and there exists  $\lambda_* > M_1$  such that*

$$N(0; \lambda) \leq 1 \quad \text{for all } \lambda > \lambda_*. \quad (40)$$

*Then (ML) holds with  $M_k = \lambda_*$ .*

**Lemma 3** (property (M)). *Let (3) holds and*

$$\begin{aligned} N(0; \lambda) &\leq 1 \quad \text{for all } \lambda > \lambda_0 = u_0(0), \\ u_0(r) &\geq U(r; \lambda_0) \quad \text{in } R_+^1. \end{aligned} \quad (41)$$

*Then (M) holds.*

**Proof of Lemma 2.** Let  $u_0 \notin \{U(r; \lambda)\}$  and (40) is valid for some  $\lambda_*$ . Fix arbitrary  $\lambda > \lambda_*$ . Let  $u(0, t_\lambda) = \lambda$  for some  $t_\lambda \in (0, T)$  such that  $u(0, t) < \lambda$  for all  $t \in (0, t_\lambda)$ . We shall show that (see also [4], [15, Chapter IV])

$$u(r, t_\lambda) \geq U(r; \lambda) \text{ for all } r \geq 0. \tag{42}$$

Suppose (42) is not valid. Then from (40) and Lemma 1 it follows that  $N(t_\lambda; \lambda) \leq 1$ .

Consider the first case when  $N(t_\lambda; \lambda) = 0$ . Then  $u(r, t_\lambda) \leq U(r; \lambda)$  in  $\omega(\lambda)$ . Hence, without loss of generality we may assume  $\text{supp } u(r, t_\lambda) \subsetneq \omega(\lambda)$ . Since by well-known properties of the interface [11]  $\text{supp } u_0 \subsetneq \text{supp } u(r, t_\lambda)$  and hence by Lemma 1 we have  $N(0; \lambda) = 0$ . Therefore,  $u_0(r) \leq U(r; \lambda)$  in  $\omega(\lambda)$  and  $u(l(\lambda), t) = 0$  for  $t \in [0, t_\lambda]$ . Then by the strong maximum principle [3, Chapter II], applied for  $u(r, t)$  in the domain of positivity near the origin, we get  $u(0, t_\lambda) < U(0; \lambda)$ , whence the contradiction.

Consider the second case when  $N(t_\lambda; \lambda) = 1$ , i.e. there exists one sign change of the difference  $w(r, t_\lambda; \lambda) \equiv u(r, t_\lambda) - U(r; \lambda)$  in  $\omega(\lambda)$ . Then by using the continuity of  $U(r; \lambda)$  with respect to  $\lambda$  we get that there exists some sufficiently small  $|\varepsilon| > 0$ ,  $\lambda + \varepsilon > \lambda_*$ , such that the difference  $w(r, t_\lambda; \lambda)$  has at least two sign changes in  $\omega(\lambda + \varepsilon)$  and  $N(t_\lambda; \lambda + \varepsilon) \geq 2$ . See the similar analysis in [6-8], [15, p. 384]. This contradicts (37) and (40).

Thus, (42) is valid. Then by the comparison theorem  $u(r, t) \geq U(r; \lambda)$  in  $\omega(\lambda)$  for all  $t \in (t_\lambda, T)$ . Since  $u(0, t_\lambda) = \lambda = U(0; \lambda) > 0$  and  $u(0, t) \geq \lambda$  for any  $t \in (t_\lambda, T)$  we obtain the inequality  $u_t(0, t_\lambda) \geq 0$  and hence (ML) is valid because  $\lambda > \lambda_*$  is arbitrary.

**Proof of Lemma 3.** Since  $u_0(r) \geq U(r; \lambda_0)$  in  $R_+^1$  by the comparison theorem  $u(r, t) \geq U(r; \lambda_0)$  in  $R_+^1 \times (0, T)$ . This implies that  $u(r, t)$  doesn't decrease with time for  $t = 0$  at any point  $r = r_*$ , where  $u_0(r_*) = U(r_*, \lambda_0) > 0$ , in particular, at the point  $r = 0$ . The end of the proof is quite similar to the proof of Lemma 2 with  $\lambda_* = \lambda_0$ .

#### 4. PROOFS OF THEOREMS 1-5

Proofs of Theorems 1-5 are based on the properties of the set of the stationary solution and on Lemmas 2 and 3.

**Proof of Theorem 1.** (i) From (20)–(22) and (8) it follows that there exists some sufficiently large  $\lambda_* = \lambda_*(R) > 0$  such that  $N(0; \lambda) \leq 1$  for all  $\lambda > \lambda_*$ . Then by Lemma 2 (i) holds.

Notice that, roughly speaking, for  $\beta \in (1, \sigma + 1)$  the property (ML) doesn't depend on the behaviour of the initial function  $u_0(r)$  in a neighbourhood of origin.

(ii) Estimates (22') with  $M = M_1$  (see (3)) and the condition (9) imply that for any sufficiently large  $\lambda > 0$  the inequality  $N(0; \lambda) \leq 1$  holds. Hence by Lemma 2 (ii) is valid.

(iii) From estimate (33) and condition (10) we get the ordinary differential inequality of the first order for the difference  $z(r) = u_0(r) - U(r; \lambda_0)$ ,  $\lambda_0 = u_0(0) > 0$ :

$$(a(r)z)_r + b(r)z > 0 \text{ in } \omega(\lambda) \cap \{u_0 > 0\}, \quad z(0) = 0,$$

where  $a(r)$  and  $b(r)$  are some smooth nonnegative functions. Then by the comparison theorem  $z \geq 0$  in  $\omega(\lambda) \cap \{u_0 > 0\}$  and hence  $u_0(r) \geq U(r; \lambda_0)$  in  $R_+^1$ . Moreover, if for  $\lambda > \lambda_0$  there exists some point of intersection  $r = r_*$  of the functions  $u_0(r)$  and  $U(r; \lambda)$  in  $\omega(\lambda)$ , then  $(u_0)_r(r_*) > U_r(r_*; \lambda)$ . This implies that  $u_0(r) \geq U(r; \lambda)$  for all  $r > r_*$  and hence  $N(0; \lambda) \leq 1$ . Then by Lemma 3 (iii) is valid.

From (i) we have

**Corollary 1.** *Let  $\beta \in (1, \sigma + 1)$ , (3) holds and let  $u_0(r)$  be a compactly supported function. Then (ML) holds.*

**Proof of Theorem 2.** (i) This is similar to the proof of (i) of Theorem 1. Explicit solution (23) and properties given in (24), (25) are used.

Note that for  $\beta = \sigma + 1$  the property (ML) at the origin depends on the behaviour of  $u_0$  in a small left neighbourhood of the point  $r = z_N$ .

(ii) See the proof of (iii) of Theorem 1.

(iii) Denote  $\lambda_0 = u_0(0)$ . Then  $u_0(r) > U(r; \lambda_0) \equiv U_S(r; \lambda_0)$  for all  $r \in (0, z_N)$  since  $u_0(r)$  is nondecreasing function in  $(0, z_N)$ . Since  $U_r(r; \lambda) < 0$ ,  $r \in (0, z_N)$ , for any  $\lambda > \lambda_0$  and  $U(r; \lambda) \equiv 0$  for  $r \geq z_N$ , we get  $N(0; \lambda) = 1$  and Lemma 3 can be used.

The assertion (i) yields

**Corollary 2.** *Let  $\beta = \sigma + 1$ , (3) holds and let  $u_0(r)$  be a compactly supported function such that  $\sup \{r > 0 \mid u_0(r) > 0\} < z_N$ . Then (ML) is valid.*

**Proof of Theorem 3.** (i) Properties (21), (22), (26) and the condition (13) guarantee the inequality  $N(0; \lambda) \leq 1$  for all sufficiently large  $\lambda > 0$ . Hence Lemma 2 can be applied. Therefore, the property (ML) depends on the behaviour of the initial function near  $r = 0$ .

(ii) See the proof of (iii) of Theorem 1.

(iii) Since  $U_r(r; \lambda) < 0$  in  $\omega(\lambda)$ , which is the domain of positivity of the function  $U(r; \lambda)$ , from (15), (18), (19) we get that  $u_0(0) = U(0; u_0(0))$  and  $u_0(r) > U(r; u_0(0))$  for  $r \in (0, l_0)$ . From (26) we obtain the inequality  $N(0; \lambda) = 1$  for all  $\lambda > u_0(0)$ . Now one can use Lemma 3.

(iv) Since  $h(r) \geq \sup_{\lambda > 0} U(r; \lambda)$  (see (35)) and  $u_0(r)$  is nondecreasing function in  $\{r > 0 \mid u_0(r) < h(r)\}$  we get that  $u_0(r) \geq U(r; u_0(0))$  for all  $r > 0$  and  $N(0; \lambda) = 1$  for all  $\lambda > u_0(0)$ . Then by Lemma 3 (iv) holds.

(v) See (19), (26) and the proof of (iii) of this Theorem.

**Proof of Theorem 4.** (i) Choose  $\delta = \varepsilon$  (see (13)). Then from (13) and (29) for all sufficiently large  $\lambda > 0$  there exists unique intersection of the functions  $u_0(r)$  and  $U(r; \lambda)$  in  $(0, \delta)$ . By using (16), (16'), (28), (29) we get that there are no points of intersection in  $[\delta, \infty)$ , i.e.  $N(0; \lambda) = 1$  for large  $\lambda > 0$ . Hence, (ML) holds.

(ii) See the proof of assertion (iii) of Theorem 1. Notice that for  $\beta \geq \beta_*$  under hypothesis (14) the initial function should be strictly positive.

(iii) See the proof of (iv) of Theorem 3.

**Proof of Theorem 5.** (i) Fix small  $\delta > 0$  such that  $h_\infty(\delta) > 2M_1$ . By using (31) for any large  $\lambda > 0$  we can choose sufficiently small  $\varepsilon > 0$  such that

$$|U(r; \lambda)/h_\infty(r) - 1| < \varepsilon, \quad |U_r(r; \lambda)/h'_\infty(r) - 1| < \varepsilon \quad (43)$$

in  $[\delta, \infty)$ . Since  $u_0 \in C^1$  in the neighbourhood of unique point  $r = r_*$ , where  $u_0(r_*) = h_\infty(r_*)$  and  $u'_0(r_*) > h'_\infty(r_*)$ , from (43) for small  $\varepsilon > 0$  and for all large  $\lambda > 0$  it follows that there exists a unique intersection of the functions  $u_0(r)$  and  $U(r; \lambda)$  in the connected interval  $\{r > 0 \mid |u_0(r)/h_\infty(r) - 1| < \varepsilon\}$ . Then by Lemma 2 (ML) is valid.

(ii) See the proof of (iii) of Theorem 1.

(iii) See the proof of (iv) of Theorem 3.

## References

- [1] S. ANGENENT: *The zero set of a solution of a parabolic equation*, J. reine angew. Math., **390**, 78-96 (1988).
- [2] A. FRIEDMAN: *On the regularity of the solutions of nonlinear elliptic and parabolic systems of partial differential equations*, J. Math. Mech., **7**, 43-59 (1958).
- [3] A. FRIEDMAN: *Partial Differential Equations of Parabolic Type*, Prentice-Hall, Englewood Cliffs, New Jersey (1964).
- [4] V. A. GALAKTIONOV: *On conditions of localization of unbounded solutions of quasilinear parabolic equations*, Doklady AN SSSR, **264**, 1035-1040 (1982) (in Russian).
- [5] V. A. GALAKTIONOV, S. P. KURDYUMOV and A. A. SAMARSKII: *On structure of limiting distributions for blow-up solutions of quasilinear parabolic equations*, Doklady AN SSSR, **301**, 781-785 (1988) (in Russian).
- [6] V. A. GALAKTIONOV and S. A. POSASHKOV: *Applications of new comparison theorems for unbounded solutions of nonlinear parabolic equations*, Differentsial'nye Uravnenia, **22**, 1165-1173 (1986) (in Russian).
- [7] V. A. GALAKTIONOV and S. A. POSASHKOV: *Any large solution of nonlinear heat conduction equation with source becomes monotone in time*, Preprint Inst. Appl. Math. Acad. Sci. USSR, n.<sup>o</sup> 109 (1989) (in Russian).
- [8] V. A. GALAKTIONOV and S. A. POSASHKOV: *On the property of monotonicity for quasilinear degenerate parabolic equation*, Differentsial'nye Uravnenia, **26**, 1127-1136 (1990) (in Russian).
- [9] Y. GIGA and R. V. KOHN: *Characterizing blow-up using similarity variables*, Indiana Univ. Math. J., **36**, 1-40 (1987).
- [10] D. D. JOSEPH and T. S. LUNDGREN: *Quasilinear Dirichlet problems driven by positive sources*, Arch. Rational Mech. Anal., **60**, 171-183 (1976).
- [11] A. S. KALASHNIKOV: *Some questions of qualitative theory of nonlinear degenerate parabolic equations of the second order*, Uspekhi Matem. Nauk, **42**, 135-176 (1987) (in Russian).
- [12] C. LOEWNER and L. NIRENBERG: *Partial differential equations invariant under conformal and projective transformations*. In: Contributions to Analysis, Academic Press, New York, 245-272 (1974).
- [13] H. MATANO: *Nonincrease of the lap number of a solution for a one-dimensional semi-linear parabolic equation*, J. Fac. Sci. Univ. Tokyo, Sec. 1A, **29**, 401-441 (1982).
- [14] K. NICKEL: *Gestaltaussagen über Lösungen Parabolischer Differenzialgleichungen*, J. reine angew. Math., **211**, 78-94 (1962).
- [15] A. A. SAMARSKII, V. A. GALAKTIONOV, S. P. KURDYUMOV and A. P. MIKHAILOV: *Blow-up in Problems for Quasilinear Parabolic Equations*, Nauka, Moscow (1987) (in Russian).
- [16] D. H. SATTINGER: *On the total variation of solutions of parabolic equations*, Math. Ann., **183**, 78-92 (1969).