

On Slice Knots in the Complex Projective Plane

AKIRA YASUHARA

ABSTRACT. We investigate the knots in the boundary of the punctured complex projective plane. Our result gives an affirmative answer to a question raised by Suzuki. As an application, we answer to a question by Mathieu.

1. INTRODUCTION

Throughout this paper, we work in the smooth category, all manifolds are oriented and all the homology groups are with integral coefficients.

Let M be a closed 4-manifold, B^4 an embedded 4-ball in M , and K a knot in $\partial(M - \text{Int } B^4)$. If K bounds a properly embedded 2-disk in $M - \text{Int } B^4$ then we call the knot K a *slice knot in M* . Let $\text{Slice}(M)$ be the set of slice knots in M . It is well-known that $\text{Slice}(S^4)$ is proper subset of the set of knots (Fox and Milnor [3]) and $\text{Slice}(S^4)$ is a subset of $\text{Slice}(M)$. In [17], Suzuki proved that $\text{Slice}(S^2 \times S^2)$ is equal to the set of knots, and asked the following question.

Question 1. *Is there a 4-manifold M such that $\text{Slice}(S^4)$ is a proper subset of $\text{Slice}(M)$ and $\text{Slice}(M)$ is a proper subset of the set of knots?*

In [20], the author has proved that $\text{Slice}(CP^2)$ does not contain a $(-2, 15)$ -torus knot. This assertion gives an affirmative answer to Question 1 since $\text{Slice}(S^4)$ is a proper subset of $\text{Slice}(CP^2)$ (Kervaire and Milnor [6]). In [20], the author could not find a knot that belongs to neither $\text{Slice}(CP^2)$ nor $\text{Slice}(\overline{CP^2})$. In Section 2, we show that there exist the knots that belongs to neither $\text{Slice}(CP^2)$ nor $\text{Slice}(\overline{CP^2})$.

Let K be a knot in $\partial(n_1 CP^2 \# n_2 \overline{CP^2} - \text{Int } B^4)$. The knot K is an *evenly slice knot* in $n_1 CP^2 \# n_2 \overline{CP^2}$ if K bounds a properly embedded 2-disk in $n_1 CP^2 \# n_2 \overline{CP^2} - \text{Int } B^4$ that represents an element $z(\varepsilon_1 \gamma_1 + \dots + \varepsilon_{n_1} \gamma_{n_1} + \bar{\varepsilon}_1 \bar{\gamma}_1 + \dots + \bar{\varepsilon}_{n_2} \bar{\gamma}_{n_2})$ in $H_2(n_1 CP^2 \# n_2 \overline{CP^2} - \text{Int } B^4, \partial)$, where $\gamma_1, \dots, \gamma_{n_1}, \bar{\gamma}_1, \dots, \bar{\gamma}_{n_2}$ are standard generators of $H_2(n_1 CP^2 \# n_2 \overline{CP^2} - \text{Int } B^4, \partial)$, $\varepsilon_i = \pm 1$, $\bar{\varepsilon}_j = \pm 1$ and z is an integer. Let $e\text{-Slice}(n_1 CP^2 \# n_2 \overline{CP^2})$ be the set of evenly slice knots in $n_1 CP^2 \# n_2 \overline{CP^2}$. (Note that $e\text{-Slice}(CP^2) = \text{Slice}(CP^2)$ and $e\text{-Slice}(\overline{CP^2}) = \text{Slice}(\overline{CP^2})$.) In Section 3, we deal with in the case $n_1 = n_2 = 1$ or $n_1 = 0$.

Let K_0 be a knot and D^2 a 2-disk intersecting transversely K_0 with the linking number $lk(\partial D^2, K_0) = l$. Let p be a positive integer and $\varepsilon = \pm 1$. By performing $\frac{\varepsilon}{p}$ -Dehn surgery along ∂D^2 , we have a new knot. The new knot is said to be the knot obtained from K_0 by an $(\varepsilon p, l)$ -twisting. Let \mathcal{H}_p be the set of knots obtained from a trivial knot by an $(\varepsilon p, l)$ -twisting for some integer l and $\varepsilon = \pm 1$. Section 4 is devoted to two applications. Our first application is to find infinitely many knots that give a negative answer to the following question given by Mathieu [12].

Question 2. For any knot K , is there a positive integer p such that $K \in \mathcal{H}_p$?

Our second one is to find infinitely many counterexamples to the following conjecture made by Akbulut and Kirby.

Conjecture. If K is a knot with Arf invariant zero, then K is obtained from a slice knot by a $(\pm 1, \pm 1)$ -twisting. (Problem 1.46 (B) of [9].)

It is shown that a $(2, 7)$ -torus knot cannot be obtained from a ribbon knot by a $(|, |)$ -twisting by using Donaldson's outstanding theorem [1, Theorem 1] (see [10]). Since then Donaldson improved this result to drop "simply connectedness assumption" [2, Theorem 1], a $(2, 7)$ -torus knot cannot be obtained from a slice knot by a $(|, |)$ -twisting. Here we give infinitely many counterexamples in different knot cobordism classes.

Similar results for Question 2 were obtained independently by Katura Miyazaki [13].

1. PRELIMINARIES

In this section we introduce some useful lemmas to us. In particular, Lemmas 1.8 and 1.11 are key lemmas in this paper.

Let α, β be the standard generators of $H_2(S^2 \times S^2)$ with $\alpha^2 = \beta^2 = 0$, $\alpha \cdot \beta = 1$ and let γ or γ_i (resp. $\bar{\gamma}$ or $\bar{\gamma}_i$) be the standard generator of $H_2(CP^2)$ (resp. $H_2(\overline{CP^2})$) with $\gamma^2 = \gamma_i^2 = 1$ (resp. $\bar{\gamma}^2 = \bar{\gamma}_i^2 = -1$). From now on a homology class in $H_2(M - \text{Int } B^4, \partial)$ is identified with its image by the homomorphism

$$H_2(M - \text{Int } B^4, \partial) \cong H_2(M - \text{Int } B^4) \rightarrow H_2(M).$$

Let l and m be nonnegative integers and $\varepsilon = \pm 1$. An $(\varepsilon l, m)$ -torus link is the link that wraps around the standardly embedded solid torus in S^3 in the longitudinal direction l times and in the meridional direction m times, where the intersection number of the meridian and longitude is ε . When l and m are relatively prime, it is a knot and called an $(\varepsilon l, m)$ -torus knot. An $(\varepsilon l, m)$ -torus knot is denoted by $T(\varepsilon l, m)$.

Let L be a μ -component link in S^3 . Let $f_i: I \times I \rightarrow S^3, i = 1, \dots, m-1$ ($m \leq \mu$) be mutually disjoint embeddings such that

- (i) $f_i(I \times I) \cap L = f_i(I \times \partial I)$ for each i ($i = 1, \dots, m-1$) and
- (ii) the link $L' = Cl(L \cup \cup f_i(\partial I \times I) - \cup f_i(I \times \partial I))$ has the orientation compatible with that of $L - \cup f_i(I \times \partial I)$ and $\cup f_i(\partial I \times I)$.

The link L' is said to be the link obtained from L by m -fusion if the number of the components of L' is $\mu - m$. In particular if the number of the components of L' is one, then L' is said to be the knot obtained from L by complete fusion. We call the images $f_1(I \times I), \dots, f_m(I \times I)$ the strips connecting L . Let $\mathcal{T}_{\varepsilon x}$ ($\varepsilon = \pm 1, x \geq 0$) be the set of knots obtained from a $(2\varepsilon, 4x)$ -torus link by 1-fusion. Note that a knot K belongs to \mathcal{T}_x if and only if the reflected inverse $-K^!$ belongs to \mathcal{T}_{-x} .

1.1. Lemma. For any knot $K \in \mathcal{T}_{\varepsilon x}$, there exists an embedded 2-disk Δ in $S^2 \times S^2 - \text{Int } B^4$ such that Δ represents an element $2\alpha + 2\varepsilon x\beta$ in $H_2(S^2 \times S^2 - \text{Int } B^4, \partial)$ and $\partial\Delta \subset \partial(S^2 \times S^2 - \text{Int } B^4)$ is $-K^!$.

Proof. We first deal with the case that $K \in \mathcal{T}_x$. It is easily seen that there exist mutually disjoint $2x+2$ properly embedded 2-disks $\Delta_1, \dots, \Delta_{2x+2}$ in $S^2 \times S^2 - \text{Int } B^4$ such that $\cup \Delta_i$ represents an element $2\alpha + 2x\beta$ and $\partial(\cup \Delta_i) \subset \partial(S^2 \times S^2 - \text{Int } B^4)$ is a Figure 1. Since a $(-2, 4x)$ -torus link is obtained from $\partial(\cup \Delta_i)$ by $2x$ -fusion, there exist $2x+1$ strips b_1, \dots, b_{2x+1} connecting the link $\partial(\cup \Delta_i)$ such that $\Delta = \Delta_1 \cup \dots \cup \Delta_{2x+2} \cup b_1 \cup \dots \cup b_{2x+1}$ is an embedded 2-disk in $S^2 \times S^2 - \text{Int } B^4$ and $\partial\Delta \subset \partial(S^2 \times S^2 - \text{Int } B^4)$ is $-K^!$.

The above argument remains valid in case $K \in \mathcal{T}_{-x}$ \square




Figure 1

1.2. Lemma. For any knot $K \in \mathcal{T}_{\varepsilon x}$, there exists an embedded 2-disk Δ in $CP^2 \# \overline{CP^2} - \text{Int } B^4$ such that Δ represents an element $(2x + \varepsilon)\gamma + (2x - \varepsilon)\tilde{\gamma}$ in $H_2(CP^2 \# \overline{CP^2} - \text{Int } B^4, \partial)$ and $\partial\Delta \subset \partial(CP^2 \# \overline{CP^2} - \text{Int } B^4)$ is $-K^1$.

Proof. We first deal with the case that $K \in \mathcal{T}_x$. Let $O_1 \cup O_{-1}$ be a 2-component trivial link in ∂B^4 such that O_j is framed by j ($j = \pm 1$). By considering the ‘‘Kirby’s calculus’’[8] as Figure 2, we note that there exist mutually disjoint $2x + 1$ properly embedded 2-disks $\Delta_1, \dots, \Delta_{2x+1}$ in $CP^2 \# \overline{CP^2} - \text{Int } B^4$ such that $\cup \Delta_i$ represents an element $(2x + 1)\gamma + (2x - 1)\tilde{\gamma}$ in $H_2(CP^2 \# \overline{CP^2} - \text{Int } B^4, \partial)$ and $\partial(\cup \Delta_i) \subset \partial(CP^2 \# \overline{CP^2} - \text{Int } B^4)$ is as Figure 3. Since a $(-2, 4x)$ -torus link is obtained from $\partial(\cup \Delta_i)$ by $(2x - 1)$ -fusion, there exist $2x$ strips b_1, \dots, b_{2x} connecting the link $\partial(\cup \Delta_i)$ such that $\Delta = \Delta_1 \cup \dots \cup \Delta_{2x+1} \cup b_1 \cup \dots \cup b_{2x}$ is an embedded 2-disk in $CP^2 \# \overline{CP^2} - \text{Int } B^4$ and $\partial\Delta \subset \partial(CP^2 \# \overline{CP^2} - \text{Int } B^4)$ is $-K^1$.

By considering the Kirby’s calculus as in Figure 4, the above argument remains valid in case $K \in \mathcal{T}_{-x}$ \square

1.3. Lemma. (Rohlin [16]) Let M be a connected, simply connected, closed 4-manifold. If $\xi \in H_2(M)$ is represented by an embedded 2-sphere in M , then




Figure 2

- (a) $\left| \frac{\xi^2}{2} - \sigma(M) \right| \leq \text{rank } H_2(M)$ if ξ is divisible by 2,
- (b) $\left| \frac{\xi^2(q^2 - 1)}{2q^2} - \sigma(M) \right| \leq \text{rank } H_2(M)$ if ξ is divisible by an odd prime integer q , where $\sigma(M)$ is the signature of M .

1.4. Lemma. (Weintraub [18], Yamamoto [19]) *Let K be a knot. If the unknotting number of K is less than or equal to u then there exists embedded 2-disk Δ in $u(CP^2 \# \overline{CP^2}) - \text{Int } B^4$ such that Δ represents the zero element in $H_2(u(CP^2 \# \overline{CP^2}) - \text{Int } B^4, \partial)$ and $\partial\Delta \subset \partial(u(CP^2 \# \overline{CP^2}) - \text{Int } B^4)$ is $-K$.*

1.5. Lemma. (Lawson [11]) *Let $\xi \in H_2(CP^2 \# 2\overline{CP^2})$ be a characteristic element. The element ξ is represented by a 2-sphere in $CP^2 \# 2\overline{CP^2}$ if and only if $\xi^2 = -1$.*

1.6. Lemma. (Lawson [11]) *Let $\xi \in H_2(CP^2 \# n\overline{CP^2})$ ($n \geq 3$) be a characteristic element. If ξ is represented by a 2-sphere in $CP^2 \# n\overline{CP^2}$ then $\xi^2 \leq -2$.*




Figure 3




Figure 4

1.7. Lemma. (Kikuchi [7]) *Let $\xi \in H_2(\overline{CP^2 \# 3CP^2})$ be a characteristic element. The element ξ is represented by a 2-sphere in $CP^2 \# 3CP^2$ if and only if $\xi^2 = -2$.*

1.8. Lemma. *Let p be a positive integer and x a nonnegative integer. Let $K \in \mathcal{F}_x$ be a knot such that the unknotting number of K is less than or equal to u . If $K \in e\text{-Slice}(p\overline{CP^2})$ then there exists an integer z such that z satisfies a condition*

- (a) $\frac{8x-4}{p} \leq z^2 \leq \frac{4u}{p} + 4$ and z is even, or
- (b) $\begin{cases} z^2 = 8x+1 \text{ if } p=1, \\ z^2 = 4x+1 \text{ if } p=2, \\ \frac{8x+2}{p} \leq z^2 \leq \frac{9}{2} \left(\frac{u}{p} + 1 \right) \text{ and } z \text{ is odd if } p \geq 3. \end{cases}$

Proof. Suppose that $K \in \mathcal{F}_x \cap e\text{-Slice}(p\overline{CP^2})$ and the unknotting number of K is less than or equal to u . Since $K \in \mathcal{F}_x \cap e\text{-Slice}(p\overline{CP^2})$, there exists an integer z such that

- (1) $2\alpha + 2x\beta + z(\bar{\epsilon}_1 \bar{\gamma}_1 + \dots + \bar{\epsilon}_p \bar{\gamma}_p) \in H_2(S^2 \times S^2 \# p\overline{CP^2})$ is represented by a 2-sphere in $S^2 \times S^2 \# p\overline{CP^2}$ and
- (2) $(2x+1)\gamma + (2x-1)\bar{\gamma} + z(\bar{\epsilon}_1 \bar{\gamma}_1 + \dots + \bar{\epsilon}_p \bar{\gamma}_p) \in H_2(CP^2 \# (p+1)\overline{CP^2})$ is represented by a 2-sphere in $CP^2 \# (p+1)\overline{CP^2}$,

by Lemmas 1.1, 1.2 and the definition of evenly slice knots. Since the unknotting number of K is less than or equal to u , by Lemma 1.4,

- (3) $z(\bar{\epsilon}_1 \bar{\gamma}_1 + \dots + \bar{\epsilon}_p \bar{\gamma}_p)$ is represented by a 2-sphere in $p\overline{CP^2} \# u(CP^2 \# \overline{CP^2})$.

In case that z is even. By Lemma 1.3, (1) and (3),

$$\left| \frac{8x - pz^2}{2} + p \right| \leq p + 2,$$

$$\left| \frac{-pz^2}{2} + p \right| \leq p + 2u.$$

It follows that

$$\frac{8x-4}{p} \leq z^2 \leq \frac{4u}{p} + 4.$$

In case that z is odd and $|z| \geq 3$. By Lemma 1.3 and (3), there exists an odd prime integer q such that

$$\left| \frac{-pz^2(q^2-1)}{2q^2} + p \right| \leq p + 2u.$$

This implies

$$(1-1) \quad z^2 \leq \frac{9}{2} \left(\frac{u}{p} + 1 \right).$$

We note that

$$(1-2) \quad 1 < \frac{9}{2} \left(\frac{u}{p} + 1 \right).$$

The inequations (1-1) and (1-2) imply that any odd integer z satisfies

$$(1-3) \quad 1 \leq z^2 \leq \frac{9}{2} \left(\frac{u}{p} + 1 \right).$$

Moreover if z is odd then $(2x+1)\gamma + (2x-1)\bar{\gamma} + z(\bar{\epsilon}\bar{\gamma}_1 + \dots + \bar{\epsilon}_p\bar{\gamma}_p)$ is a characteristic element in $H_2(CP^2 \# (p+1)CP^2)$. By Lemmas 1.5, 1.6, 1.7 and (2),

$$(1-4) \quad 8x - z^2 = -1 \text{ if } p=1,$$

$$(1-5) \quad 8x - 2z^2 = -2 \text{ if } p=2,$$

$$(1-6) \quad 8x - pz^2 \leq -2 \text{ if } p \geq 3.$$

By (1-3), (1-4), (1-5) and (1-6), we have

$$z^2 = 8x + 1 \text{ if } p=1,$$

$$z^2 = 4x + 1 \text{ if } p=2,$$

$$\frac{8x+2}{p} \leq z^2 \leq \frac{9}{2} \left(\frac{u}{p} + 1 \right) \text{ if } p \geq 3.$$

This completes the proof. \square

Suppose that knots K_+ and K_- have representatives in S^3 that are identical outside a 3-ball within which they are as in Figure 5. Then we say that K_- is obtained from K_+ by *changing a positive crossing* and that K_+ is obtained from K_- by *changing a negative crossing*. We define the *positive unknotting number* (resp. *negative unknotting number*) of a knot K , to be the minimum, over all sequences transforming K to be a trivial knot, of the number of positive (resp. negative) crossings which are changed. If K cannot be a trivial knot by changing only positive (resp. negative) crossings, then we define the positive unknotting number (resp. negative unknotting number) of K is *infinite*.




Figure 5

1.9. Lemma. (Weintraub [18]) *Let K be a knot. If the positive unknotting number (resp. negative unknotting number) of K is less than or equal to u , then there exists an embedded 2-disk Δ in $u\mathbb{C}P^2 - \text{Int } B^4$ (resp. $u\mathbb{C}P^2 - \text{Int } B^4$) such that Δ represents the zero element in $H_2(u\mathbb{C}P^2 - \text{Int } B^4, \partial)$ (resp. $H_2(u\mathbb{C}P^2 - \text{Int } B^4, \partial)$) and $\partial\Delta \subset \partial(u\mathbb{C}P^2 - \text{Int } B^4)$ (resp. $\partial\Delta \subset \partial(u\mathbb{C}P^2 - \text{Int } B^4)$) is $-K$.*

1.10. Lemma. (Kervaire and Milnor [6]) *Let M be a connected, simply connected, closed 4-manifold. Let $\xi \in H_2(M)$ be a characteristic element. If ξ is represented by an embedded 2-sphere in M , then $\xi^2 \equiv \sigma(M) \pmod{16}$.*

1.11. Lemma. *Let p be a positive integer and x a nonnegative integer. Let $K \in \mathcal{T}_{-x}$ be a knot such that the negative unknotting number of K is finite. If $K \in e\text{-Slice}(p\overline{CP^2})$ then there exists an integer z such that z satisfies a condition*

- (a) $z^2 \leq 4 + \frac{4-8x}{p}$ and z is even, or
- (b) $\begin{cases} z^2 = 1 \text{ only if } x=0 \text{ and } p=1, 2, \\ z^2 = 1 \text{ only if } x \equiv 0 \pmod{2} \text{ and } p \geq 3. \end{cases}$

Proof. Suppose $K \in \mathcal{T}_{-x} \cap e\text{-Slice}(p\overline{CP^2})$ and the negative unknotting number of K is u . Since $K \in \mathcal{T}_{-x} \cap e\text{-Slice}(p\overline{CP^2})$, there exists an integer z such that

$$(4) \quad 2\alpha - 2x\beta + z(\bar{\epsilon}_1 \bar{\gamma}_1 + \dots + \bar{\epsilon}_p \bar{\gamma}_p) \in H_2(S^2 \times S^2 \# p\overline{CP^2}) \text{ is represented by a 2-sphere in } S^2 \times S^2 \# p\overline{CP^2} \text{ and}$$

$$(5) \quad (2x-1)\gamma + (2x+1)\bar{\gamma} + z(\bar{\epsilon}_1 \bar{\gamma}_1 + \dots + \bar{\epsilon}_p \bar{\gamma}_p) \in H_2(CP^2 \# (p+1)\overline{CP^2}) \text{ is represented by a 2-sphere in } CP^2 \# (p+1)\overline{CP^2},$$

by Lemmas 1.1, 1.2 and the definition of evenly slice knots. Since the negative unknotting number of K is u , by Lemma 1.9,

$$(6) \quad z(\bar{\epsilon}_1 \bar{\gamma}_1 + \dots + \bar{\epsilon}_p \bar{\gamma}_p) \text{ is represented by a 2-sphere in } p\overline{CP^2} \# u\overline{CP^2}.$$

In case that z is even. By Lemma 1.3 and (4),

$$\left| \frac{-8x - pz^2}{2} + p \right| \leq p + 2.$$

This implies

$$z^2 \leq 4 + \frac{4-8x}{p}.$$

In case that z is odd. If $|z| \geq 3$, then by Lemma 1.3 and (6), there exists an odd prime integer q such that

$$\left| \frac{-pz^2(q^2-1)}{2q^2} + p - u \right| \leq p + u.$$

It follows that

$$z^2 \leq \frac{9}{2}.$$

This is a contradiction. Thus $|z| = 1$. Moreover, by Lemmas 1.5, 1.7, 1.10 and (5), we have

$$-8x - pz^2 = -p \text{ if } p = 1, 2,$$

$$-8x - pz^2 \equiv -p \pmod{16}.$$

Since $|z| = 1$,

$$-8x = 0 \text{ if } p = 1, 2,$$

$$-8x \equiv 0 \pmod{16}.$$

This implies

$$x = 0 \text{ if } p = 1, 2,$$

$$x \equiv 0 \pmod{2}.$$

This completes the proof. \square

2. SLICE KNOTS IN CP^2 or $\overline{CP^2}$

In this section we shall prove the following two theorems.

2.1. Theorem. *Let x be a positive integer.*

(a) *If Slice($\overline{CP^2}$) contains $T(2, 4x-1)$, then $2x-1$, $2x$ or $8x+1$ is a square number.*

(b) *If Slice($\overline{CP^2}$) contains $T(2, 4x+1)$, then $2x$, $2x+1$ or $8x+1$ is a square number.*

2.2. Theorem. *Let t be a nonnegative integer. The set Slice($\overline{CP^2}$) does not contain $T(-2, 2t+1)$ if and only if $t \geq 2$.*

2.3. Remark. Since Slice(CP^2) contains a knot K if and only if Slice($\overline{CP^2}$) contains $-K^!$, Slice(CP^2) contains $T(l, m)$ if and only if

$\text{Slice}(\overline{CP^2})$ contains $T(-l, m)$. It follows that Theorems 2.1 and 2.2 imply that there exist infinitely many integer $x_i (i=1, 2, \dots)$ such that $T(2, 2x_i+1)$ belongs to neither $\text{Slice}(CP^2)$ nor $\text{Slice}(\overline{CP^2})$ for any x_i .

2.4. Lemma. For any $T(2\varepsilon, 4x+1)$ ($\varepsilon=\pm 1, x\geq 0$), there exists an embedded 2-disk Δ in $CP^2\#\overline{CP^2}-\text{Int } B^4$ such that Δ represents an element $(2x+1+\varepsilon)\gamma + (2x+1-\varepsilon)\bar{\gamma}$ in $H_2(CP^2\#\overline{CP^2}-\text{Int } B^4, \partial)$ and $\partial\Delta \subset \partial(CP^2\#\overline{CP^2}-\text{Int } B^4)$ is $T(-2\varepsilon, 4x+1)$.

Proof. By considering the Kirby's calculus as in Figure 2, we note that there exist mutually disjoint $2x+2$ properly embedded 2-disk $\Delta_1, \dots, \Delta_{2x+2}$ in $CP^2\#\overline{CP^2}-\text{Int } B^4$ such that $\cup \Delta_i$ represents an element $(2x+2)\gamma + 2x\bar{\gamma}$ in $H_2(CP^2\#\overline{CP^2}-\text{Int } B^4, \partial)$ and $\partial(\cup \Delta_i) \subset \partial(CP^2\#\overline{CP^2}-\text{Int } B^4)$ is as Figure 6. Since a $(-2, 4x+2)$ -torus link is obtained from $\partial(\cup \Delta_i)$ by $2x$ -fusion, there exist $2x+1$ strips b_1, \dots, b_{2x+1} connecting the link $\partial(\cup \Delta_i)$ such that $\Delta = \Delta_1 \cup \dots \cup \Delta_{2x+2} \cup b_1 \cup \dots \cup b_{2x+1}$ is an embedded 2-disk in $CP^2\#\overline{CP^2}-\text{Int } B^4$ and $\partial\Delta \subset \partial(CP^2\#\overline{CP^2}-\text{Int } B^4)$ is $T(-2, 4x+1)$.

By considering the Kirby's calculus as in Figure 4, the above argument remains valid for $T(-2, 4x+1)$. \square




Figure 6

Proof of Theorem 2.1. Suppose $T(2, 4x - 1) \in \text{Slice}(\overline{CP^2})$. Since the unknotting number of $T(2, 4x - 1)$ is $2x - 1$, $T(2, 4x - 1) \in \mathcal{T}_x$ and $e\text{-Slice}(\overline{CP^2}) = \text{Slice}(\overline{CP^2})$, by Lemma 1.8, there exists an integer z such that z satisfies a condition

$$(2-7) \quad 8x - 4 \leq z^2 \leq 8x \text{ and } z \text{ is even, or}$$

$$(2-8) \quad z^2 = 8x + 1.$$

We set $z = 2k$ in (2-7), then we have

$$2x - 1 \leq k^2 \leq 2x.$$

It follows that

$$(2-9) \quad k^2 = 2x - 1, 2x.$$

By (2-8) and (2-9), we obtain Theorem 2.1 (a).

Suppose $T(2, 4x + 1) \in \text{Slice}(\overline{CP^2})$. Since the unknotting number of $T(2, 4x + 1)$ is $2x$ and $T(2, 4x + 1) \in \mathcal{T}_x$, by Lemma 1.8, there exists an integer z such that z satisfies a condition

$$(2-10) \quad 8x - 4 \leq z^2 \leq 8x + 4 \text{ and } z \text{ is even, or}$$

$$(2-11) \quad z^2 = 8x + 1.$$

The fact that $T(2, 4x + 1)$ belongs to $\text{Slice}(\overline{CP^2})$ and Lemma 2.4 imply that $(2x + 2)\gamma + 2x\bar{\gamma} + z\bar{\gamma}_1 \in H_2(CP^2 \# 2CP^2)$ is represented by a 2-sphere in $CP^2 \# 2CP^2$. If z is even, then by Lemma 1.3, we have

$$\left| \frac{8x + 4 - z^2}{2} + 1 \right| \leq 3.$$

This implies

$$(2-12) \quad 8x \leq z^2 \leq 8x + 12.$$

By (2-10) and (2-12), we have

$$(2-13) \quad 8x \leq z^2 \leq 8x + 4 \text{ and } z \text{ is even.}$$

We set $z = 2k$ in (2-13) then

$$2x \leq k^2 \leq 2x + 1.$$

It follows that

$$(2-14) \quad k^2 = 2x, 2x + 1.$$

By (2-11) and (2-14), we obtain Theorem 2.1 (b). \square

2.5. Proposition. *If $t \geq 3$ then $\text{Slice}(\overline{CP^2})$ does not contain $T(-2, 2t+1)$.*

Proof. Note that \mathcal{F}_{-x} contains both $T(-2, 4x-1)$ and $T(-2, 4x+1)$ and that the negative unknotting number of $T(-2, 4x-1)$ and that the negative unknotting number of $T(-2, 4x+1)$ are finite. If $\text{Slice}(\overline{CP^2})$ contains $T(-2, 4x-1)$ or $T(-2, 4x+1)$, then by Lemma 1.11, there exists an integer z such that z satisfies a condition

$$(2-15) \quad z^2 = 8 - 8x \text{ and } z \text{ is even, or}$$

$$(2-16) \quad z^2 = 1 \text{ and } x = 0.$$

The conditions (2-15) and (2-16) imply

$$x = 0, 1.$$

This completes the proof. \square

2.5.1. Remark. By the proofs of Lemma 1.11 and Proposition 2.5, we note that if $\text{Slice}(\overline{CP^2})$ contains $T(-2, 5)$ then there exists a properly embedded 2-disk Δ in $\overline{CP^2} - \text{Int } B^4$ such that Δ represents the zero element in $H_2(\overline{CP^2} - \text{Int } B^4, \partial)$ and $\partial\Delta \subset \partial(\overline{CP^2} - \text{Int } B^4)$ is $T(-2, 5)$.

2.6. Proposition. *The set $\text{Slice}(\overline{CP^2})$ does not contain $T(-2, 5)$.*

Proof. Suppose $\text{Slice}(\overline{CP^2})$ contains $T(-2, 5)$. Remark 2.5.1 and Lemma 2.4 imply that $2\gamma + 4\bar{\gamma} \in H_2(CP^2 \# \overline{CP^2})$ is represented by a 2-sphere in $CP^2 \# \overline{2CP^2}$. By Lemma 1.3, we have

$$\left| \frac{4-16}{2} + 1 \right| \leq 3.$$

This is a contradiction. \square

Proof of Theorem 2.2. By Propositions 2.5 and 2.6, if $t \geq 2$ then $Slice(\overline{CP^2})$ does not contain $T(-2, 2t+1)$. If $t=0$ or 1 then $Slice(\overline{CP^2})$ contains $T(-2, 2t+1)$, see Proposition 3.7. \square

3. EVENLY SLICE KNOTS IN $n_1 CP^2 \# n_2 \overline{CP^2}$

In [15], Norman proved that $Slice(CP^2 \# \overline{CP^2})$ is equal to the set of knots, but the following theorem implies that there exist infinitely many knots that do not belong to $e\text{-Slice}(CP^2 \# \overline{CP^2})$, i.e., $e\text{-Slice}(CP^2 \# \overline{CP^2})$ is a proper subset of $Slice(CP^2 \# \overline{CP^2})$.

3.1. Theorem. Let t be a nonnegative integer and $\varepsilon = \pm 1$. The set $e\text{-Slice}(CP^2 \# \overline{CP^2})$ contains $T(2\varepsilon, 2t+1)$ if and only if $t=0$ or 1.

3.2. Lemma. (Hirai [4]) Let $\xi \in H_2(2(CP^2 \# \overline{CP^2}))$ be a characteristic element. The element ξ represented by a 2-sphere in $2(CP^2 \# \overline{CP^2})$ if and only if $\xi^2=0$.

3.3. Proposition. For $\varepsilon = \pm 1$, if $t \geq 3$ then $e\text{-Slice}(CP^2 \# \overline{CP^2})$ does not contain $T(2\varepsilon, 2t+1)$.

Proof. Let x be a nonnegative integer. If either $T(2\varepsilon, 4x-1)$ or $T(2\varepsilon, 4x+1)$ belongs to $e\text{-Slice}(CP^2 \# \overline{CP^2})$ then there exists an integer z such that

$$(7) \quad 2\alpha + 2\varepsilon x\beta + z(\varepsilon_1 \gamma_1 + \bar{\varepsilon}_1 \bar{\gamma}_1) \in H_2(S^2 \times S^2 \# CP^2 \# \overline{CP^2}) \text{ is represented by a 2-sphere in } S^2 \times S^2 \# CP^2 \# \overline{CP^2} \text{ and}$$

$$(8) \quad (2x + \varepsilon)\gamma + (2x - \varepsilon)\bar{\gamma} + z(\varepsilon_1 \gamma_1 + \bar{\varepsilon}_1 \bar{\gamma}_1) \in H_2(2(CP^2 \# \overline{CP^2})) \text{ is represented by a 2-sphere in } 2(CP^2 \# \overline{CP^2}),$$

by Lemmas 1.1, 1.2 and the definition of evenly slice knots. If z is even, then by Lemma 1.3 and (7),

$$\left| \frac{8\varepsilon x}{2} \right| \leq 4.$$

This implies

$$x = 0, 1.$$

If z is odd, then by Lemma 3.2 and (8),

$$8\epsilon x = 0.$$

It follows that if $x \geq 2$, then neither $T(2\epsilon, 4x-1)$ nor $T(2\epsilon, 4x+1)$ belongs to $e\text{-Slice}(CP^2\#\overline{CP^2})$. This completes the proof. \square

3.4. Proposition. *The set $e\text{-Slice}(CP^2\#\overline{CP^2})$ does not contain $T(2\epsilon, 5)$ for $\epsilon = \pm 1$.*

Proof. Suppose $e\text{-Slice}(CP^2\#\overline{CP^2})$ contains $T(2\epsilon, 5)$. Proof of Proposition 3.3 and Lemma 2.4 implies that there exists an even integer z such that $(3+\epsilon)\gamma + (3-\epsilon)\bar{\gamma} + z(\epsilon_1\gamma_1 + \bar{\epsilon}_1\bar{\gamma}_1) \in H_2(2(CP^2\#\overline{CP^2}))$ is represented by a 2-sphere in $2(CP^2\#\overline{CP^2})$. By Lemma 1.3, we have

$$\left| \frac{12\epsilon}{2} \right| \leq 4.$$

This is a contradiction. \square

Proof of Theorem 3.1. By Propositions 3.3 and 3.4, if $t \geq 2$ then $e\text{-Slice}(CP^2\#\overline{CP^2})$ does not contain $T(2\epsilon, 2t+1)$. If $t=0$ or 1 then $e\text{-Slice}(CP^2\#\overline{CP^2})$ contains $T(2\epsilon, 2t+1)$, see Proposition 3.7. \square

The same arguments as proof of Theorem 2.1 and Proposition 2.5 lead to the following Theorem 3.5 and Proposition 3.6, respectively.

3.5. Theorem. *Let x be a positive integer.*

- (a) *If $e\text{-Slice}(2\overline{CP^2})$ contains $T(2, 4x-1)$ then x or $4x+1$ is a square number.*
- (b) *If $e\text{-Slice}(2\overline{CP^2})$ contains $T(2, 4x+1)$ then x , $x+1$ or $4x+1$ is a square number.*

3.6. Proposition. *If $t \geq 3$ then $e\text{-Slice}(2\overline{CP^2})$ does not contain $T(-2, 2t+1)$.*

3.7. Proposition. *Let K be a knot. If the positive unknotting number or the negative unknotting number of K is less than or equal to p , then both $e\text{-Slice}(pCP^2)$ and $e\text{-Slice}(p\overline{CP^2})$ contain K .*

Proof. Suppose K is a knot and the positive or negative unknotting number of K is less than or equal to p . Let L_ε be the Hopf link in $\partial(CP^2 - \text{Int } B^4)$ with linking number $\varepsilon (\varepsilon = \pm 1)$. It is easily seen that L_ε bounds a properly embedded 2-disk in $CP^2 - \text{Int } B^4$ that represents an element $(1-\varepsilon)\gamma$ in $H_2(CP^2 - \text{Int } B^4, \partial)$. Since the positive or negative unknotting number of K is less than or equal to p , K is obtained from the p copies of L_ε by complete fusion. It follows that K bounds a properly embedded 2-disk in $pCP^2 - \text{Int } B^4$ that represents an element $(1-\varepsilon)(\varepsilon_1\gamma_1 + \dots + \varepsilon_p\gamma_p)$ in $H_2(pCP^2 - \text{Int } B^4, \partial)$. This implies that K belongs to $e\text{-Slice}(pCP^2)$.

The above argument remains valid to show that K belongs to $e\text{-Slice}(p\overline{CP^2})$. This completes the proof. \square

By Propositions 3.6 and 3.7, we have the following theorem.

3.8. Theorem. *Let t be a nonnegative integer. The set $e\text{-Slice}(2\overline{CP^2})$ does not contain $T(-2, 2t+1)$ if and only if $t \geq 3$.*

3.9. Theorem. *For any integer $p \geq 3$, $e\text{-Slice}(p\overline{CP^2})$ contains neither $T(2, 8p+3)$ nor $T(-2, 8p+3)$.*

Proof. Suppose that $e\text{-Slice}(p\overline{CP^2})$ contains $T(2, 8p+3)$. Since $T(2, 8p+3)$ belongs to \mathcal{T}_{2p+1} and the unknotting number of $T(2, 8p+3)$ is $4p+1$, by Lemma 1.8, there exists an integer z such that z satisfies a condition

$$(3-17) \quad \frac{16p+4}{-p} \leq z^2 \leq \frac{16p+4}{p} + 4 \text{ and } z \text{ is even, or}$$

$$(3-18) \quad \frac{16p+10}{p} \leq z^2 \leq \frac{9}{2} \left(\frac{4p+1}{p} + 1 \right) \text{ and } z \text{ is odd.}$$

Since $p \geq 3$, (3-17) and (3-18) imply

$$16 < z^2 < 25 \text{ and } z \text{ is even,}$$

$$16 < z^2 < 25 \text{ and } z \text{ is odd.}$$

This is a contradiction.

Suppose that $e\text{-Slice}(p\overline{CP^2})$ contains $T(-2, 8p+3)$. Since $T(-2, 8p+3)$ belongs to \mathcal{T}_{-2p-1} and the negative unknotting number of $T(-2, 8p+3)$ is

finite, by Lemma 1.11, there exists an integer z such that z satisfies the following condition

$$z^2 \leq 4 + \frac{-16p-4}{p-1} < 0.$$

This is a contradiction. \square

3.10. Claim. Let K be a knot. Neither $e\text{-Slice}(pCP^2)$ nor $e\text{-Slice}(\overline{pCP^2})$ contains K if and only if $e\text{-Slice}(\overline{pCP^2})$ contains neither K nor $-K$.

3.11. Remark. By Theorem 3.9 and Claim 3.10, we have that $T(2, 8p+3)$ belongs to neither $e\text{-Slice}(pCP^2)$ nor $e\text{-Slice}(\overline{pCP^2})$ for any $p \geq 3$.

4. APPLICATIONS

4.1. Proposition. If $K \in \mathcal{S}_p$ then K belongs to either $e\text{-Slice}(pCP^2)$ or $e\text{-Slice}(\overline{pCP^2})$.

Proof. If $K \in \mathcal{S}_p$ then there exists a 2-disk D^2 and a trivial knot K_0 in S^3 such that K is obtained from K_0 by $\frac{\varepsilon}{p}$ -Dehn surgery along ∂D^2 . We take the parallel copies D_1^2, \dots, D_p^2 of D^2 as in Figure 7. It is easily seen that K is obtained from K_0 by Dehn surgery along $\partial(\cup D_i^2)$ in which the surgery coefficients are all ε . Suppose that K_0 and $\cup D_i^2$ are in the boundary of a 4-ball B_0^4 , then K_0 bounds a properly embedded 2-disk Δ in B_0^4 . Let $\{h_i^2\}$ ($1 \leq i \leq p$) be 2-handles on B_0^4 whose attaching sphere are $\{\partial D_i^2\}$ and all framings are ε . We note that $K_0 \subset \partial(B_0^4 \cup \cup h_i^2)$ is K , K bounds the 2-disk Δ in $B_0^4 \cup \cup h_i^2$ and $B_0^4 \cup \cup h_i^2$ is diffeomorphic to either punctured pCP^2 or punctured $\overline{pCP^2}$. Let the punctured pCP^2 and punctured $\overline{pCP^2}$ be denoted by $pCP^2 - \text{Int } B^4$ and $\overline{pCP^2} - \text{Int } B^4$, respectively. Suppose the linking number $lk(K_0, \partial D^2) = z$ then $lk(K_0, \partial D_i^2)$ ($1 \leq i \leq p$) are the same number as z . It is not hard to see that Δ represents either an element $z(\varepsilon_1 \gamma_1 + \dots + \varepsilon_p \gamma_p)$ in $H_2(pCP^2 - \text{Int } B^4, \partial)$ or an element $z(\bar{\varepsilon}_1 \bar{\gamma}_1 + \dots + \bar{\varepsilon}_p \bar{\gamma}_p)$ in $H_2(\overline{pCP^2} - \text{Int } B^4, \partial)$. This implies that K belongs to either $e\text{-Slice}(pCP^2)$ or $e\text{-Slice}(\overline{pCP^2})$. \square

By Remark 3.11, Proposition 4.1 and the definition of evenly slice knots, we have the following theorem.

4.2. Theorem. For any integer $p \geq 3$, \mathcal{S}_p does not contain any knot that is cobordant to $T(2, 8p+3)$.




Figure 7

By Lemmas 1.8 and 1.11, we have the following proposition.

4.3. Proposition. *For any p ($1 \leq p \leq 5$), $e\text{-Slice}(p\overline{CP^2})$ contains neither $T(2, 75)$ nor $T(-2, 75)$.*

By Claim 3.10, Propositions 4.1, 4.3 and the definition of evenly slice knots, we have the following proposition.

4.4. Proposition. *For any p ($1 \leq p \leq 5$), \mathcal{K}_p does not contain any knot that is cobordant to $T(2, 75)$.*

4.5. Lemma. (Moteji [14]) *If $p \geq 6$ then \mathcal{K}_p does not contain any composite knot.*

Let K be a nontrivial slice knot. Proposition 4.4 and Lemma 4.5 imply that \mathcal{K}_p does not contain $T(2, 75)\#K$ for any $p \geq 1$. Hence we have the following theorem that gives a negative answer to Question 2.

4.6. Theorem. *There exist infinitely many knots that do not belong to any \mathcal{K}_p ($p \geq 1$).*

Let K be a knot in $\partial(CP^2\#\overline{CP^2} - \text{Int } B^4)$. If K is obtained from a slice knot by a $(\pm 1, \pm 1)$ -twisting, then by proof of Proposition 4.1, K bounds a properly embedded 2-disk in $CP^2\#\overline{CP^2} - \text{Int } B^4$ that represents an element $\pm\gamma_1$ or $\pm\tilde{\gamma}_1$ in $H_2(CP^2\#\overline{CP^2} - \text{Int } B^4, \partial)$. It follows that K bounds a properly embedded 2-disk in $CP^2\#\overline{CP^2} - \text{Int } B^4$ that represent an element $\pm\gamma_1 + \tilde{\gamma}_1$ or $\gamma_1 \pm \tilde{\gamma}_1$ in $H_2(CP^2\#\overline{CP^2} - \text{Int } B^4, \partial)$. We have the following proposition.

4.7. Proposition. *If K is obtained from a slice knot by $(\pm 1, \pm 1)$ -twisting, then K belongs to $e\text{-Slice}(CP^2\#\overline{CP^2})$.*

Since a $(\pm 1, \pm 1)$ -twisting does not change the Arf invariant of a knot, thus $T(2\varepsilon, 3)$ cannot be obtained from a slice knot by a $(\pm 1, \pm 1)$ -twisting. By Theorem 3.1, Proposition 4.7 and the definition of evenly slice knots, we have the following theorem.

4.8. Theorem. *Let t be a nonnegative integer and $\varepsilon = \pm 1$. A knot cobordant to $T(2\varepsilon, 2t+1)$ is obtained from a slice knot by a $(\pm 1, \pm 1)$ -twisting if and only if $t = 0$.*

If $2t+1 \equiv \pm 1 \pmod{8}$, then the Arf invariant of $T(2\varepsilon, 2t+1)$ is zero (for example, see p266 in [5]). Thus Theorem 4.8 gives infinitely many counterexamples to Conjecture.

Acknowledgement. The author would like to thank Professor Shin'ichi Suzuki for his encouragement.

References

- [1] S. K. DONALDSON: *An application of gauge theory to four-dimensional topology*, J. Diff. Geometry, 18(1983), 279-315.
- [2] S. K. DONALDSON: *The orientation of Yang-Mills moduli spaces and 4-manifold topology*, J. Diff. Geometry, 26(1987), 397-428.
- [3] R. H. FOX and J. W. MILNOR: *Singularities of 2-spheres in 4-space and cobordism of knots*, Osaka J. Math., 3(1966), 257-267.
- [4] Y. HIRAI: *Representing homology classes of connected sums of 2-sphere bundles over 2-spheres*, Kobe J. Math., 6(1989), 233-240.
- [5] L. H. KUAFFMAN: *On knots*, Ann. of Math. Studies, 115, Princeton Univ. Press, Princeton, N. J., (1987).
- [6] M. A. KERVAIRE and J. W. MILNOR: *On 2-sphere in 4-manifolds*, Proc. Nat. Acad. Sci. U.S.A., 47(1961), 1651-1657.
- [7] K. KIKUCHI: *Representing positive homology classes of $CP^2 \# 2\overline{CP^2}$ and $CP^2 \# 3CP^2$* , to appear in Proc. Amer. Math. Soc.
- [8] R. KIRBY: *A calculus for framed links in S^3* , Invent. Math. 45(1978), 35-56.
- [9] R. KIRBY: *Problems in low-dimensional manifold theory*, Proc. Symp. Pure Math. 32, American Mathematical Society, Providence, Rhode Island (1978), 273-312.
- [10] R. KIRBY: *4-manifold problems*, Contemp. Math., 35(1984), 513-528.
- [11] T. LAWSON: *Representing homology classes of almost definite 4-manifolds*, Michigan Math. J., 34(1987), 85-91.
- [12] Y. MATHIEU: *Sur des noeuds qui ne sont pas déterminés par leur complément et problèmes de chirurgie dans les variétés de dimension 3*, These, L'Université de Provence, 1990.
- [13] K. MIYAZAKI: private communication.
- [14] K. MOTEGI: *Primeness of twisted knots*, preprint.
- [15] R. A. NORMAN: *Dehn's Lemma for certain 4-manifolds*, Invent. Math., 7(1969), 143-147.
- [16] V. A. ROHLIN: *Two-dimensional submanifolds of four-dimensional manifolds*, Functional Anal. Appl., 5(1974), 39-48.
- [17] S. SUZUKI: *Local knots of 2-spheres in 4-manifolds*, Proc. Japan Acad., 45(1969), 34-38.
- [18] S. H. WEINTRAUB: *Inefficiently embedded surfaces in 4-manifolds*, Algebraic Topology Aarhus 1978 (J. L. Dupont and I. H. Madsen, eds.), Lecture Notes in Math., 763, Springer-Verlag, Berlin and New York, 1979.
- [19] M. YAMAMOTO: *Lower bounds for the unknotting numbers of certain torus knots*, Proc. Amer. Math. Soc., 86(1982), 519-524.

- [20] A. YASUHARA: *(2, 15)-torus knot is not slice in CP^2* , Proc. Japan Acad. ser. A. Math. Sci., 67(1991), 353-355.

Department of Mathematics
School of Science and Engineering
Waseda University
Tokyo 169-50
JAPAN

Recibido: 18 de marzo de 1992.