

A Generalized Mixed Topology on Orlicz Spaces

MARIAN NOWAK

ABSTRACT. Let L^φ be an Orlicz space defined by an arbitrary Orlicz function φ over a positive measure space (Ω, Σ, μ) and provided with its usual F -norm $\|\cdot\|_\varphi$. In L^φ a natural convergence can be defined as follows: a sequence (x_n) in L^φ is said to be γ_φ -convergent to $x \in L^\varphi$ whenever $x_n \rightarrow x$ ($\mu - \Omega$) and $\sup \|x_n\|_\varphi < \infty$. In this paper we examine some kind of generalized inductive-limit topology (in the sense of Turpin) \mathcal{J}_I^φ in L^φ that generates our γ_φ -convergence in L^φ . The main aim of the paper is to obtain a description of the topology \mathcal{J}_I^φ in terms of some family of F -norms defined by other Orlicz functions. As an application we obtain a topological characterization of the γ_φ -convergence in L^φ .

1. INTRODUCTION AND PRELIMINARIES

Every Orlicz space L^φ defined by an Orlicz function φ (not necessarily convex) over a measure space (Ω, Σ, μ) can be equipped with two

F -norms: $\|\cdot\|_\varphi$ - the usual F -norm on L^φ and $\|\cdot\|_\mu$ - the F -norm of convergence in measure (on Ω) restricted to L^φ . Thus a natural sequential convergence in L^φ can be defined as follows: a sequence (x_n) in L^φ is said to be γ_φ -convergent to $x \in L^\varphi$, in symbols $x_n \xrightarrow{\gamma_\varphi} x$, whenever

$$x_n \rightarrow x(\mu - \Omega) \text{ (i.e., } \|x_n - x\|_\mu \rightarrow 0 \text{) and } \sup_n \|x_n\|_\varphi < \infty.$$

When we replace in the above definition the condition: $\sup \|x_n\|_\varphi < \infty$ with the boundedness of the set $\{x_n : n \geq 0\}$ for the topology $\mathcal{J}_{\|\cdot\|_\varphi}$, then this new convergence comes under the definition of the so-called two-norm convergence or γ -convergence in the sense of Alexiewicz ([1,1954]). The general theory of two-norm convergence has been extensively developed by A.Alexiewicz [1], W. Orlicz [19], A. Alexiewicz and Z. Semadeni [2], A. Wiweger [23], [24], [25].

It is well known that the theory of two-norm convergence is closely related to the Wiweger's theory of mixed topologies [23], [24]. Indeed, in case when $\|\cdot\|$ is a homogenous norm and $\|\cdot\|^*$ is an F -norm on a linear space X and $\|x_n - x\|^* \rightarrow 0$ implies $\liminf \|x_n\| \geq \|x\|$, then the sequential γ -convergence in X is generated by the so-called mixed topology $\gamma[\mathcal{J}_{\|\cdot\|}, \mathcal{J}_{\|\cdot\|^*}]$.

The notion of the mixed topology was a starting point for the theory of generalized inductive-limit topologies. There are many kinds of such topologies introduced for different reasons by A. Persson [21], D.J.H. Garling [7], J.B. Cooper [3], P. Turpin [22] and others.

The question arises whether our γ_φ -convergence in L^φ is topologized by some linear topology. It turns out that there is a positive answer to this question when we take into account an appropriate generalized inductive-limit topology in the sense of Turpin. This topology will be called here a generalized mixed topology and denoted by \mathcal{J}_I^φ . This term is justified by the fact that \mathcal{J}_I^φ coincides with the usual mixed topology $\gamma[\mathcal{J}_{\|\cdot\|_\varphi}, \mathcal{J}_{\mu|_{L^\varphi}}]$ (in the sense of Wiweger) when the space $(L^\varphi, \mathcal{J}_{\|\cdot\|_\varphi})$ is locally bounded.

In this paper we investigate the generalized mixed topology \mathcal{J}_I^φ . Our main aim is to obtain a description of \mathcal{J}_I^φ in terms of some family of F -norms defined by other Orlicz functions. As application we obtain

a topological characterization of our γ_φ -convergence in L^φ . Moreover, for φ being a convex Orlicz function we establish the general form of \mathcal{J}_I^φ -continuous linear functionals on L^φ .

In some special cases the topology \mathcal{J}_I^φ was examined by P. Turpin [22] and the author [14], [15], [16].

Given a linear topological space (X, ξ) by $Bd(X, \xi)$ we will denote the collection of all ξ -bounded subsets of X . As usual \mathcal{N} stands for the set of all natural numbers. We assume that $0 \cdot \infty = 0$.

Now we recall some notation and terminology concerning Orlicz spaces (see [9], [11], [12], [22] for more details).

By an Orlicz function we mean a function $\varphi : [0, \infty) \rightarrow [0, \infty]$ which is non-decreasing, left continuous, continuous at 0 with $\varphi(0) = 0$, and not identically equal to 0.

An Orlicz function φ is called convex whenever $\varphi(\alpha u + \beta v) \leq \alpha\varphi(u) + \beta\varphi(v)$ for $\alpha, \beta \geq 0$, $\alpha + \beta = 1$ and $u, v \geq 0$. A convex Orlicz function is usually called a Young function.

For a Young function φ we denote by φ^* the function complementary to φ in the sense of Young, i.e.,

$$\varphi^*(v) = \sup\{uv - \varphi(u) : u \geq 0\} \text{ for } v \geq 0.$$

For a set Ψ of Young functions we will write: $\Psi^* = \{\psi^* : \psi \in \Psi\}$.

Let φ and ψ be a pair of Orlicz functions vanishing only at zero (resp. taking only finite values). We say that φ increases essentially more rapidly than ψ for small u (resp. for large u) in symbols $\psi \overset{s}{\ll} \varphi$ (resp. $\psi \overset{l}{\ll} \varphi$) whenever for any $c > 0$, $\psi(cu)/\varphi(u) \rightarrow 0$ as $u \rightarrow 0$ (resp. $u \rightarrow \infty$).

We will write $\psi \overset{a}{\ll} \varphi$ when $\psi \overset{s}{\ll} \varphi$ and $\psi \overset{l}{\ll} \varphi$ hold.

For φ and ψ being Young functions the condition $\psi \overset{s}{\ll} \varphi$ (resp. $\psi \overset{l}{\ll} \varphi$) implies $\varphi^* \overset{s}{\ll} \psi^*$ (resp. $\varphi^* \overset{l}{\ll} \psi^*$) (see [9, Lemma 13.1]).

Let (Ω, Σ, μ) be a positive measure space, and let L^0 denote the set of equivalence classes of all real valued measurable functions defined

and finite a.e. on Ω . For a subset A of Ω and $x \in L^0$ we will write $x_A = x \cdot \chi_A$, where χ_A stands for the characteristic function of A .

An Orlicz function φ determines a functional $m_\varphi : L^0 \rightarrow [0, \infty]$ by

$$m_\varphi(x) = \int_{\Omega} \varphi(|x(t)|) d\mu.$$

The Orlicz space generated by φ is the ideal of L^0 defined by

$$L^\varphi = \{x \in L^0 : m_\varphi(\lambda x) < \infty \text{ for some } \lambda > 0\}.$$

The functional m_φ restricted to L^φ is an orthogonally additive semi-modular.

L^φ can be equipped with the complete metrizable topology \mathcal{J}_φ of the F -norm

$$\|x\|_\varphi = \inf\{\lambda > 0 : m_\varphi(x/\lambda) \leq \lambda\}.$$

Moreover, if φ is a Young function, then the topology \mathcal{J}_φ can be generated by the Luxemburg norm

$$\| \|x\| \|_\varphi = \inf\{\lambda > 0 : m_\varphi(x/\lambda) \leq 1\}.$$

For $r > 0$ let

$$B_\varphi(r) = \{x \in L^\varphi : \|x\|_\varphi \leq r\}$$

and let

$$B_{(\varphi)}(r) = \{x \in L^\varphi : \| \|x\| \|_\varphi \leq r\}$$

whenever φ is a Young function.

We shall need the following lemma.

Lemma 1.1. *Let φ_1, φ_2 be Orlicz functions, and let $\varphi(u) = \varphi_1(u) \vee \varphi_2(u)$ for $u \geq 0$. Then φ is an Orlicz function and the following statements hold:*

(i) $L^\varphi = L^{\varphi_1} \cap L^{\varphi_2}$.

(ii) $\|x\|_{\varphi_1} \vee \|x\|_{\varphi_2} \leq \|x\|_\varphi \leq \|x\|_{\varphi_1} + \|x\|_{\varphi_2}$ for $x \in L^\varphi$.

(iii) $\mathcal{J}_\varphi = \mathcal{J}_{\varphi_1|_{L^\varphi}} \vee \mathcal{J}_{\varphi_2|_{L^\varphi}}$

and $Bd(L^\varphi, \mathcal{J}_\varphi) = Bd(L^\varphi, \mathcal{J}_{\varphi_1|_{L^\varphi}}) \cap Bd(L^\varphi, \mathcal{J}_{\varphi_2|_{L^\varphi}})$.

Proof. (i) See [8, Theorem 1].

(ii) It follows from the definition of $\|\cdot\|_\varphi$.

(iii) It follows from (ii).

Let

$$E^\varphi = \{x \in L^0 : m_\varphi(\lambda x) < \infty \text{ for all } \lambda > 0\}.$$

It is known that $L^\varphi = E^\varphi$ whenever φ satisfies the Δ_2 -condition, i.e., $\limsup \varphi(2u)/\varphi(u) < \infty$ as $u \rightarrow 0$ and $u \rightarrow \infty$.

Let

$$\varphi_0(u) = \begin{cases} 0 & \text{for } 0 \leq u \leq 1, \\ 1 & \text{for } u > 1. \end{cases}$$

It is known that L^{φ_0} is the largest Orlicz space and consists of all those $x \in L^0$ that are bounded outside of some set of finite measure, and

$$\|x\|_{\varphi_0} = \inf\{\lambda > 0 : \mu(\{t \in \Omega : |x(t)| > \lambda\}) \leq \lambda\}.$$

It is seen that $\|x_n - x\|_{\varphi_0} \rightarrow 0$ in L^{φ_0} iff $x_n \rightarrow x$ in measure on Ω (in symbols $x_n \rightarrow x$ (μ - Ω)). Therefore we will write $\|\cdot\|_\mu$ instead of $\|\cdot\|_{\varphi_0}$, and by \mathcal{J}_μ we will denote the topology of the F -norm $\|\cdot\|_{\varphi_0}$.

For $\varepsilon > 0$ let

$$B_\mu(\varepsilon) = \{x \in L^{\varphi_0} : \|x\|_\mu \leq \varepsilon\}.$$

We shall need the following lemma.

Lemma 1.2. *Let φ be an Orlicz function such that $\varphi(u) \rightarrow \infty$ as $u \rightarrow \infty$. Then for $r > 0$, $B_\varphi(r) \in Bd(L^\varphi, \mathcal{J}_{\mu|_{L^\varphi}})$.*

Proof. Let $x_n \in B_\varphi(r)$ ($n = 1, 2, \dots$) and let $\lambda_n \rightarrow 0$. For $\varepsilon > 0$ let $\Omega_n(\varepsilon) = \{t \in \Omega : |\lambda_n x_n(t)| > \varepsilon\}$. Then

$$\begin{aligned} \mu(\Omega_n(\varepsilon))\varphi\left(\frac{\varepsilon}{r|\lambda_n|}\right) &\leq \int_{\Omega_n(\varepsilon)} \varphi\left(\frac{|x_n(t)|}{r}\right) d\mu \\ &\leq m_\varphi\left(\frac{x_n}{r}\right) \leq r. \end{aligned}$$

Since $\varphi(u) \rightarrow \infty$ as $u \rightarrow \infty$ we get $\mu(\Omega_n(\varepsilon)) \rightarrow 0$, and this means that $\|\lambda_n x_n\|_\mu \rightarrow 0$.

2. A GENERALIZED MIXED TOPOLOGY ON L^φ - GENERAL PROPERTIES

In this section we consider some kind of generalized inductive limit topology on L^φ .

Let φ be an arbitrary Orlicz function, and let

$$F_n = B_\varphi(2^n) \text{ and } \mathcal{J}_n = \mathcal{J}_{\mu|_{F_n}} \text{ for } n \geq 0.$$

Then the family $\mathcal{B}_\varphi = \{F_n : n \geq 0\}$ forms a base of metric bounded sets in $(L^\varphi, \|\cdot\|_\varphi)$.

Moreover, the sequence (F_n, \mathcal{J}_n) ($n \geq 0$) of balanced topological spaces satisfies the following conditions:

- (i) $L^\varphi = \bigcup_{n \geq 0} F_n$.
- (ii) $F_n + F_n \subset F_{n+1}$, and the function

$$F_n \times F_n \ni (x, y) \rightarrow x + y \in F_{n+1}$$

is continuous.

- (iii) The function $[-1, 1] \times F_n \ni (\lambda, x) \mapsto \lambda \cdot x \in F_n$ is continuous.
- (iv) $\mathcal{J}_{n+1}|_{F_n} = \mathcal{J}_n$ for $n \geq 0$.

Thus the space L^φ with the system $\{(F_n, \mathcal{J}_n) : n \geq 0\}$ comes under the conditions of the strict inductive limit of balanced topological spaces in the sense of P. Turpin [22, Ch. I].

Definition 2.1. *The family of all sets of the form*

$$\bigcup_{N=0}^{\infty} \left(\sum_{n=0}^N (B_\varphi(2^n) \cap B_\mu(\varepsilon_n)) \right) \quad (2.1)$$

where $(\varepsilon_n : n \geq 0)$ is a sequence of positive numbers, forms a base of neighbourhoods of zero for a linear topology on L^φ (in the sense of Turpin) which will be denoted by \mathcal{J}_I^φ .

According to [22, Theorem 1.1.6] \mathcal{J}_I^φ is the finest of all linear topologies ξ on L^φ which satisfy the conditions:

$$\xi|_{F_n} \subset \mathcal{J}_\mu|_{F_n} \text{ for } n \geq 0. \quad (2.2)$$

Moreover, in view of [22, Theorem 1.1.8] we have

$$\mathcal{J}_I^\varphi|_{F_n} = \mathcal{J}_\mu|_{F_n} \text{ for } n \geq 0. \quad (2.3)$$

Since $\mathcal{J}_\mu|_{L^\varphi} \subset \mathcal{J}_\varphi$ we have $\mathcal{J}_I^\varphi \subset \mathcal{J}_\varphi$; hence $\mathcal{J}_\mu|_{L^\varphi} \subset \mathcal{J}_I^\varphi \subset \mathcal{J}_\varphi$.

Henceforth in this section we assume that $\varphi(u) \rightarrow \infty$ as $u \rightarrow \infty$.

The basic properties of the topology \mathcal{J}_I^φ are included in the following theorems.

Theorem 2.1. *The space $(L^\varphi, \mathcal{J}_I^\varphi)$ is complete.*

Proof. It is known that the balls $B_\varphi(2^n)$ are closed subsets of $(L^{\varphi_0}, \mathcal{J}_\mu)$ (see [22, 0.3.6]), so the spaces $(B_\varphi(2^n), \mathcal{J}_\mu|_{B_\varphi(2^n)})$ ($n \geq 0$) are complete. Hence, by [22, Theorem 1.1.10] the space $(L^\varphi, \mathcal{J}_I^\varphi)$ is complete.

Theorem 2.2. *For a subset $Z \subset L^\varphi$ the following statements are equivalent:*

- (i) $\sup\{\|x\|_\varphi : x \in Z\} < \infty$.
(ii) Z is bounded for \mathcal{J}_I^φ .

Proof. By Lemma 1.2 the balls $B_\varphi(2^n)$ are bounded subsets of $(L^\varphi, \mathcal{J}_{\mu|_{L^\varphi}})$. Moreover, the balls $B_\varphi(2^n)$ are also closed in $(L^\varphi, \mathcal{J}_{\mu|_{L^\varphi}})$ (see [22, 0.3.6]). In view of (2.2) and (2.3) \mathcal{J}_I^φ is the finest of all linear topologies ξ on L^φ such that $\xi|_{F_n} = \mathcal{J}_{\mu|_{F_n}}$. Hence by [22, Corollary 1.1.12] the equivalence (i) \Leftrightarrow (ii) holds.

Theorem 2.3. *For a subset $Z \subset L^\varphi$ the following statements are equivalent:*

- (i) Z is relatively compact for \mathcal{J}_I^φ .
(ii) Z is relatively compact for $\mathcal{J}_{\mu|_{L^\varphi}}$ and

$$\sup\{\|x\|_\varphi : x \in Z\} < \infty.$$

Proof. It follows from Theorem 2.2 and (2.3).

Let us recall that a sequence (x_n) in L^φ is said to be γ_φ -convergent to $x \in L^\varphi$, in symbols $x_n \xrightarrow{\gamma_\varphi} x$, whenever

$$x_n \rightarrow x \ (\mu - \Omega) \quad \text{and} \quad \sup_n \|x_n\|_\varphi < \infty.$$

Theorem 2.4. *For a sequence (x_n) in L^φ the following statements are equivalent:*

- (i) $x_n \rightarrow 0$ for \mathcal{J}_I^φ .
(ii) $x_n \xrightarrow{\gamma_\varphi} 0$.

Moreover, \mathcal{J}_I^φ is the finest of all linear topologies ξ on L^φ which satisfy the condition:

$$x_n \xrightarrow{\gamma_\varphi} 0 \quad \text{implies} \quad x_n \rightarrow 0 \text{ for } \xi. \quad (+)$$

Proof. The equivalence (i) \Leftrightarrow (ii) follows from Theorem 2.2 and (2.3).

Now let ξ be a linear topology on L^φ for which the condition (+) holds. Then $\xi|_{B_\varphi(r)} \subset \mathcal{J}_\mu|_{B_\varphi(r)}$ for $r > 0$, because \mathcal{J}_μ is a metrizable linear topology. Hence by (2.2) we get that $\xi \subset \mathcal{J}_I^\varphi$.

Definition 2.2. Let (X, η) be a linear topological space. A linear mapping $A : L^\varphi \rightarrow X$ is said to be γ_φ -linear, if

$$x_n \xrightarrow{\gamma_\varphi} 0 \quad \text{implies} \quad A(x_n) \rightarrow 0 \text{ for } \eta.$$

The next theorem gives a characterization of γ_φ -linear functionals on L^φ .

Theorem 2.5. For a linear topological space (X, η) and a linear mapping $A : L^\varphi \rightarrow X$ the following statements are equivalent:

- (i) A is $(\mathcal{J}_I^\varphi, \eta)$ -continuous.
- (ii) A is γ_φ -linear.
- (iii) For every $r > 0$, the restriction $A|_{B_\varphi(r)}$ is $(\mathcal{J}_\mu|_{B_\varphi(r)}, \eta)$ -continuous.

Proof. (i) \Rightarrow (ii) It follows from Theorem 2.4.

(ii) \Rightarrow (iii) it is obvious.

(iii) \Rightarrow (i) Let W be a neighbourhood of zero in X for η . Then there exists a sequence $(W_n : n \geq 0)$ of neighbourhoods of zero for η such that $\sum_{n=0}^N W_n \subset W$ for every $N \geq 0$. Thus by our assumption there exists a sequence $(\varepsilon_n : n \geq 0)$ of positive numbers such that $A(B_\varphi(2^n) \cap B_\mu(\varepsilon_n)) \subset W_n$. Thus for $N \geq 0$

$$A\left(\sum_{n=0}^N (B_\varphi(2^n) \cap B_\mu(\varepsilon_n))\right) \subset \sum_{n=0}^N W_n \subset W,$$

so

$$\begin{aligned} & A\left(\bigcup_{N=0}^{\infty} \left(\sum_{n=0}^N (B_{\varphi}(2^n) \cap B_{\mu}(\varepsilon_n))\right)\right) \subset \\ & \subset \bigcup_{N=0}^{\infty} A\left(\sum_{n=0}^N (B_{\varphi}(2^n) \cap B_{\mu}(\varepsilon_n))\right) \subset W. \end{aligned}$$

This means that A is $(\mathcal{J}_I^{\varphi}, \eta)$ -continuous.

Now we are going to compare the topology \mathcal{J}_I^{φ} with the mixed topology $\gamma[\mathcal{J}_{\varphi}, \mathcal{J}_{\mu|L^{\varphi}}]$ in the sense of Wiweger (see [24]). For this purpose we shall need the following

Theorem 2.6. *Assume that (Ω, Σ, μ) is an atomless measure space or that μ is the counting measure on \mathcal{N} . If $(L^{\varphi}, \mathcal{J}_{\varphi})$ is a locally bounded space then for a subset Z of L^{φ} the following statements are equivalent:*

- (i) Z is bounded for \mathcal{J}_I^{φ} .
- (ii) $\sup \{\|x\|_{\varphi} : x \in Z\} < \infty$.
- (iii) Z is bounded for \mathcal{J}_{φ} .

Proof. (i) \Leftrightarrow (ii) See Theorem 2.2.

(ii) \Rightarrow (iii) In view of [22, 0.3.10.2] $\sup\{\|x\|_{\varphi} : x \in Z\} < \infty$ if and only if Z is additively bounded (see [22, 0.3.10.1]), so arguing as in the proof of [15, Lemma 2.5] we obtain that Z is bounded for \mathcal{J}_{φ} .

(iii) \Rightarrow (i) Obvious.

Theorem 2.7. *Assume that (Ω, Σ, μ) is an atomless measure space or that μ is the counting measure on \mathcal{N} . If $(L^{\varphi}, \mathcal{J}_{\varphi})$ is a locally bounded space, then the generalized mixed topology \mathcal{J}_I^{φ} coincides with the mixed topology $\gamma[\mathcal{J}_{\varphi}, \mathcal{J}_{\mu|L^{\varphi}}]$.*

Proof. In view of Theorem 2.6 it follows from [24, 2.2.1 and 2.2.2].

3. SOME PROJECTIVE TOPOLOGY ON ORLICZ SPACES

In [5], [6] H.W. Davis, F.J.Murray and J. Weber studied the spaces

$$L^P(S) = \bigcap_{p \in S} L^p (S \subset [1, \infty))$$

endowed with the appropriate projective topology.

There are some results concerning a representation of an Orlicz space L^φ as the intersection of some family of other Orlicz spaces (see [10], [17], [18]). In this section we examine the appropriate projective topology on L^φ . In section 4 we shall show that this projective topology coincides with the generalized mixed topology \mathcal{J}_I^φ .

We start with some equalities among Orlicz spaces, proved in [17] and [18], which are of key importance in this section. At the very beginning we distinguish some classes of Orlicz functions.

An Orlicz function φ continuous for all $u \geq 0$, taking only finite values, vanishing only at zero, and not bounded is usually called a φ -function. By Φ we will denote the collection of all φ -functions.

A Young function φ vanishing only at zero and taking only finite values is called an N -function whenever $\varphi(u)/u \rightarrow 0$ as $u \rightarrow 0$ and $\varphi(u)/u \rightarrow \infty$ as $u \rightarrow \infty$. By Φ_N we will denote the collection of all N -functions.

Let Φ_1 be the set of all Orlicz functions φ vanishing only at zero and such that $\varphi(u) \rightarrow \infty$ as $u \rightarrow \infty$. Denote by

$$\Phi_{11} = \{\varphi \in \Phi_1 : \varphi(u) < \infty \text{ for } u > 0\},$$

$$\Phi_{12} = \{\varphi \in \Phi_1 : \varphi \text{ jumps to } \infty\}.$$

Then $\Phi_1 = \Phi_{11} \cup \Phi_{12}$. In view of [17, Theorem 3.1, 3.2, 3.7 and 3.8] we get

Theorem 3.1. *Let $\varphi \in \Phi_{1i}$ ($i = 1, 2$). Then the following equalities hold:*

$$L^\varphi = \bigcap \{L^\psi : \psi \in \Psi_{1i}^\varphi\} = \bigcap \{E^\psi : \psi \in \Psi_{1i}^\varphi\}$$

where

$$\Psi_{11}^\varphi = \{\psi \in \Phi : \psi \overset{a}{\ll} \varphi\}, \quad \Psi_{12}^\varphi = \{\psi \in \Phi : \psi \overset{s}{\ll} \varphi\}.$$

Moreover, if μ is an atomless measure or the counting measure on \mathcal{N} , then the strict inclusion $L^\varphi \subsetneq E^\psi$ holds for each $\psi \in \Psi_{1i}^\varphi$.

Next let Φ_1^c be the set of all Young functions φ vanishing only at zero and such that $\varphi(u)/u \rightarrow \infty$ as $u \rightarrow \infty$.

Denote by

$$\Phi_{11}^c = \{\varphi \in \Phi_1^c : \varphi(u) < \infty \text{ for } u > 0 \text{ and } \varphi(u)/u \rightarrow 0 \text{ as } u \rightarrow 0\},$$

$$\Phi_{12}^c = \{\varphi \in \Phi_1^c : \varphi \text{ jumps to } \infty \text{ and } \varphi(u)/u \rightarrow 0 \text{ as } u \rightarrow 0\},$$

$$\Phi_{13}^c = \{\varphi \in \Phi_1^c : \varphi(u) < \infty \text{ for } u > 0 \text{ and } \varphi(u)/u \rightarrow a \text{ as } u \rightarrow 0, a > 0\},$$

$$\Phi_{14}^c = \{\varphi \in \Phi_1^c : \varphi \text{ jumps to } \infty \text{ and } \varphi(u)/u \rightarrow a \text{ as } u \rightarrow 0, a > 0\}.$$

Then $\Phi_1^c = \bigcup_{i=1}^4 \Phi_{1i}^c$, where the sets Φ_{1i}^c are pairwise disjoint. It is seen that $\Phi_{11}^c = \Phi_N$. According to [18, Theorems 2.1-2.4] we get

Theorem 3.2. *Let $\varphi \in \Phi_{1i}^c$ ($i = 1, 2, 3, 4$). Then the following equalities hold:*

$$L^\varphi = \bigcap \{L^\psi : \psi \in \Psi_{1i}^\varphi(N)\} = \bigcap \{E^\psi : \psi \in \Psi_{1i}^\varphi(N)\}$$

where

$$\Psi_{11}^\varphi(N) = \{\psi \in \Phi_N : \psi \overset{a}{\ll} \varphi\}, \quad \Psi_{12}^\varphi(N) = \{\psi \in \Phi_N : \psi \overset{s}{\ll} \varphi\},$$

$$\Psi_{13}^\varphi(N) = \{\psi \in \Phi_N : \psi \overset{l}{\ll} \varphi\}, \quad \Psi_{14}^\varphi(N) = \Phi_N.$$

Next, let Φ_2^c be the set of all Young functions φ taking only finite values and such that $\varphi(u)/u \rightarrow 0$ as $u \rightarrow 0$.

Denote by

$$\Phi_{21}^c = \{\varphi \in \Phi_2^c : \varphi(u) > 0 \text{ for } u > 0 \text{ and } \varphi(u)/u \rightarrow \infty \text{ as } u \rightarrow \infty\},$$

$$\begin{aligned}\Phi_{22}^c &= \{\varphi \in \Phi_2^c : \varphi(u) > 0 \text{ for } u > 0 \text{ and } \varphi(u)/u \rightarrow a \text{ as } u \rightarrow \infty, a > 0\}, \\ \Phi_{23}^c &= \{\varphi \in \Phi_2^c : \varphi(u) = 0 \text{ near zero and } \varphi(u)/u \rightarrow \infty \text{ as } u \rightarrow \infty\}, \\ \Phi_{24}^c &= \{\varphi \in \Phi_2^c : \varphi(u) = 0 \text{ near zero and } \varphi(u)/u \rightarrow a \text{ as } u \rightarrow \infty, a > 0\}.\end{aligned}$$

Then $\Phi_2^c = \bigcup_{i=1}^4 \Phi_{2i}^c$, where the sets Φ_{2i}^c are pairwise disjoint. It is seen that $\Phi_{21}^c = \Phi_N$. According to [18, Theorems 1.1-1.4] we have

Theorem 3.3. *Let $\varphi \in \Phi_{2i}^c$ ($i = 1, 2, 3, 4$). Then the following equalities hold*

$$E^\varphi = \bigcup \{E^\psi : \psi \in \Psi_{2i}^\varphi(N)\} = \bigcup \{L^\psi : \psi \in \Psi_{2i}^\varphi(N)\}$$

where

$$\begin{aligned}\Psi_{21}^\varphi(N) &= \{\psi \in \Phi_N : \varphi \overset{a}{\prec} \psi\}, \quad \Psi_{22}^\varphi(N) = \{\psi \in \Psi_N : \varphi \overset{s}{\prec} \psi\}, \\ \Psi_{23}^\varphi(N) &= \{\psi \in \Phi_N : \varphi \overset{l}{\prec} \psi\}, \quad \Psi_{24}^\varphi(N) = \Phi_N.\end{aligned}$$

At last, in view of [18, Lemma 3.1 and Theorem 3.3] we get

Theorem 3.4 *Let φ_1 and φ_2 be a pair of complementary Young functions. Then $\varphi_1 \in \Phi_{1i}^c$ if and only if $\varphi_2 \in \Phi_{2i}^c$ ($i = 1, 2, 3, 4$), and moreover, the sets $\Psi_{1i}^\varphi(N)$ and $\Psi_{2i}^\varphi(N)$ are mutually related in such a way that*

$$(\Psi_{1i}^{\varphi_1}(N))^* = \Psi_{2i}^{\varphi_2}(N) \text{ and } (\Psi_{2i}^{\varphi_2}(N))^* = \Psi_{1i}^{\varphi_1}(N).$$

We shall need the following

Corollary 3.5. *Let $\varphi \in \Phi_{1i}^c$ ($i = 1, 2, 3, 4$). Then*

$$E^{\varphi^*} = \bigcup \{L^{\psi^*} : \psi \in \Psi_{1i}^\varphi(N)\}.$$

Proof. Since $\varphi \in \Phi_{2i}^c$ and $(\Psi_{1i}^\varphi(N))^* = \Psi_{2i}^{\varphi^*}(N)$ (see Theorem 3.4) by Theorem 3.3 we get

$$\begin{aligned} \bigcup \{L^{\psi^*} : \psi \in \Psi_{1i}^\varphi(N)\} &= \bigcup \{L^\psi : \psi \in (\Psi_{1i}^\varphi(N))^*\} \\ &= \bigcup \{L^\psi : \psi \in \Psi_{2i}^{\varphi^*}(N)\} = E^{\varphi^*}. \end{aligned}$$

We are now ready to define our projective topology on L^φ .

Definition 3.1. Let $\varphi \in \Phi_{1i}$ ($i = 1, 2$). By \mathcal{J}_P^φ we will denote the projective topology on L^φ with respect to the family $\{(E^\psi, \mathcal{J}_{\psi|E^\psi}) : \psi \in \Psi_{1i}^\varphi\}$, i.e., \mathcal{J}_P^φ is defined to be the coarsest of all linear topologies ξ on L^φ for which $\mathcal{J}_{\psi|L^\varphi} \subset \xi$ holds for every $\psi \in \Psi_{1i}^\varphi$. Thus

$$\mathcal{J}_P^\varphi = \sup \{\mathcal{J}_{\psi|L^\varphi} : \psi \in \Psi_{1i}^\varphi\}.$$

For φ being a φ -function the topology \mathcal{J}_P^φ has been examined in [14], [15], [16]. It is easy to verify that all properties of \mathcal{J}_P^φ which are obtained in [14], [15], [16] for φ being a φ -function remain valid for $\varphi \in \Phi_{11}$. In this section we extend results from [14], [15], [16] to the case of φ belonging to Φ_1 .

From the definition of \mathcal{J}_P^φ we have

Theorem 3.6. Let $\varphi \in \Phi_1$. Then $\mathcal{J}_{\mu|L^\varphi} \subset \mathcal{J}_P^\varphi \subset \mathcal{J}_\varphi$.

Theorem 3.7. Let $\varphi \in \Phi_1$ and let μ be an infinite atomless measure. Then there exists a sequence (x_n) in L^φ such that $x_n \rightarrow 0$ for \mathcal{J}_P^φ and $m_\varphi(x_n) = 1$ for $n \in \mathcal{N}$. Hence the strict inclusion $\mathcal{J}_P^\varphi \subsetneq \mathcal{J}_\varphi$ holds.

Proof. For $\varphi \in \Phi_{11}$ this fact is proved in [13, Theorem 2.5]. Now let $\varphi \in \Phi_{12}$, i.e., $\varphi(u) < \infty$ for $u \leq a$ and $\varphi(u) = \infty$ for $u > a$. Let (u_n) be a sequence of positive numbers such that $u_n \downarrow 0$ and $u_1 < a$. Let (Ω_n) be a sequence of measurable subsets of Ω such that $\mu(\Omega_n) = 1/\varphi(u_n)$. Define

$$x_n(t) = \begin{cases} u_n & \text{for } t \in \Omega_n, \\ 0 & \text{for } t \notin \Omega_n. \end{cases}$$

We shall show that $x_n \rightarrow 0$ for \mathcal{J}_P^φ , i.e., $\|x_n\|_\psi \rightarrow 0$ for each $\psi \in \Psi_{12}^\varphi$. Indeed, let $\psi \overset{s}{\prec} \varphi$ and let $\varepsilon > 0$ be given. Then there exists $u_0 > 0$ such that $\psi(u/\varepsilon) \leq \varepsilon\varphi(u)$ for $u \leq u_0$. Let $n_0 \in \mathcal{N}$ be such that $u_n \leq u_0$ for $n \geq n_0$. Then for $n \geq n_0$ we have $m_\psi(x_n/\varepsilon) = \psi(u_n/\varepsilon)/\varphi(u_n) \leq \varepsilon$ i.e., $\|x_n\|_\psi \leq \varepsilon$. On the other hand, $m_\varphi(x_n) = \varphi(u_n)/\varphi(u_n) = 1$.

Arguing as in the proof of [13, Theorem 1.2] we get

Theorem 3.8. *Let $\varphi \in \Phi_{1i}$ ($i = 1, 2$). Then the topology \mathcal{J}_P^φ has a base of neighbourhoods of zero consisting of all sets of the form:*

$$B_\psi(r) \cap L^\varphi$$

where $\psi \in \Psi_{1i}^\varphi$ and $r > 0$.

Repeating the arguments of the proof of [13, Theorem 5.1] and using the equalities from Theorem 3.1 we get

Theorem 3.9. *Let $\phi \in \Phi_1$. Then the space $(L^\varphi, \mathcal{J}_P^\varphi)$ is complete.*

Since the space $(L^\varphi, \mathcal{J}_\varphi)$ is complete, from Theorems 3.6 and 3.7, in view of the Open Mapping Theorem it follows

Theorem 3.10. *Let $\varphi \in \Phi_1$ and let μ be an infinite atomless measure. Then the space $(L^\varphi, \mathcal{J}_P^\varphi)$ is not metrizable.*

To the end of this section we will assume that $\varphi \in \Phi_1^c$. We start with the following lemma.

Lemma 3.11. *Let $\varphi \in \Phi_{1i}^c$ ($i = 1, 2, 3, 4$) and let ψ be a φ -function such that $\psi \overset{a}{\prec} \varphi$ if $i = 1$ (resp. $\psi \overset{s}{\prec} \varphi$ if $i = 2$, $\psi \overset{a}{\prec} \varphi$ if $i = 3$, $\psi \overset{s}{\prec} \varphi$ if $i = 4$). Then there exists an N -function ψ_0 such that $\psi(u) \leq \psi_0(2u)$ for $u \geq 0$ and $\psi_0 \overset{a}{\prec} \varphi$ if $i = 1$ (resp. $\psi_0 \overset{s}{\prec} \varphi$ if $i = 2$, $\psi_0 \overset{l}{\prec} \varphi$ if $i = 3$).*

Proof. Let ψ_1 be an arbitrary N -function such that $\psi_1 \overset{a}{\prec} \varphi$ if $i = 1$ (resp. $\psi_1 \overset{s}{\prec} \varphi$ if $i = 2$, $\psi_1 \overset{l}{\prec} \varphi$ if $i = 3$, $\psi_1 \overset{s}{\prec} \varphi$ if $i = 4$). Let $\psi_2 = \psi \vee \psi_1$. Next, let us put $p(0) = 0$ and $p(s) = \sup_{0 < t \leq s} (\psi_2(t)/t)$ for $s > 0$. Let

$$\psi_0(u) = \int_0^u p(s) ds \text{ for } u \geq 0.$$

It is seen that ψ_0 is an N -function. Arguing as in the proof of [13, Lemma 1.4] we can verify that ψ_0 satisfies the desired properties.

Theorem 3.12. *Let $\varphi \in \Phi_{1i}^c$ ($i = 1, 2, 3, 4$). Then the topology \mathcal{J}_P^φ is generated by the family of B -norms $\{|||\cdot|||_{\psi|L^\varphi} : \psi \in \Psi_{1i}^\varphi(N)\}$.*

Proof. For example, let $\varphi \in \Phi_{13}^c$. Then $\varphi \in \Phi_{11}$. Given $\psi \in \Psi_{11}^\varphi$ and $r > 0$, in view of Lemma 3.11 there exists $\psi_0 \in \Psi_{13}^\varphi(N)$ and such that $\psi(u) \leq \psi_0(2u)$ for $u \geq 0$. Hence

$$||x||_\psi \leq ||2x||_{\psi_0} \text{ for all } x \in L^{\psi_0}. \quad (1)$$

On the other hand, since the F -norms $||\cdot||_{\psi_0}$ and $|||\cdot|||_{\psi_0}$ are equivalent on L^{ψ_0} , there exists $r_1 > 0$ such that

$$B_{(\psi_0)}(r_1) \subset B_{\psi_0}(r). \quad (2)$$

We shall show that $B_{(\psi_0)}(r_1/2) \cap L^\varphi \subset B_\psi(r)$. Indeed, let $x \in B_{(\psi_0)}(r_1/2) \cap L^\varphi$. Then $|||2x|||_{\psi_0} \leq r_1$, hence by (2) we get $||2x||_{\psi_0} \leq r$ and next, by (1) we see that $||x||_\psi \leq r$.

For $i=1,2,4$ the proof is similar.

Now we are ready to establish the general form of \mathcal{J}_P^φ -continuous linear functionals on L^φ .

Theorem 3.13. *Let $\varphi \in \Phi_1^c$ and let μ be a σ -finite measure. Then for a linear functional f on L^φ the following statements are equivalent:*

(i) f is continuous for \mathcal{J}_P^φ .

(ii) *There exists a unique $y \in E^{\varphi^*}$ such that*

$$f(x) = f_y(x) = \int_{\Omega} x(t)y(t) d\mu \quad \text{for } x \in L^{\varphi}.$$

Proof. (i) \Rightarrow (ii). Let $\varphi \in \Phi_{1i}^c$ ($i = 1, 2, 3, 4$). In view of Theorem 3.12 there exist $\psi \in \Psi_{1i}^{\varphi}(N)$ and $\tau > 0$ such that f is bounded on $B_{(\psi)}(\tau) \cap L^{\varphi}$. This means that f is continuous on the linear subspace $(L^{\varphi}, \mathcal{J}_{\psi|L^{\varphi}})$ of the normed space $(E^{\psi}, \mathcal{J}_{\psi|E^{\psi}})$. Hence, by the Hahn-Banach theorem there exists a $\mathcal{J}_{\psi|E^{\psi}}$ -continuous linear functional \bar{f} on E^{ψ} such that $\bar{f}(x) = f(x)$ for $x \in L^{\varphi}$. According to [11, p. 56] there exists $y \in L^{\psi^*} \subset E^{\varphi^*}$ such that

$$\bar{f}(x) = \int_{\Omega} x(t)y(t) d\mu \quad \text{for } x \in E^{\psi}.$$

Hence

$$f(x) = f_y(x) = \int_{\Omega} x(t)y(t) d\mu \quad \text{for } x \in L^{\varphi}. \quad (1)$$

Now assume that there exists another $y' \in E^{\varphi^*}$ such that

$$f(x) = f_{y'}(x) = \int_{\Omega} x(t)y'(t) d\mu \quad \text{for } x \in L^{\varphi}. \quad (2)$$

Then, for example, there exists a measurable set $A \subset \{t \in \Omega : y'(t) > y(t)\}$ such that $0 < \mu(A) < \infty$. Hence by (1) and (2) we get

$$\int_{\Omega} \chi_A(t) (y'(t) - y(t)) d\mu = \int_A (y'(t) - y(t)) d\mu = 0.$$

This contradiction establishes that there exists a unique $y \in E^{\varphi^*}$ such that (1) holds.

(ii) \Rightarrow (i) Let $\varphi \in \Phi_{1i}^c$ ($i = 1, 2, 3, 4$). According to Corollary 3.5 there exists $\psi \in \Psi_{1i}^{\varphi}(N)$ such that $y \in L^{\psi^*}$. Then $L^{\varphi} \subset E^{\psi} \subset L^{\psi}$ and using the Holder's inequality we get that $|f_y(x)| \leq \|y\|_{\psi^*}^0 \|x\|_{\psi}$ for

$x \in L^\varphi$ (here $\|\cdot\|_{\psi^*}^0$ denotes the Orlicz norm on L^{ψ^*}). This means that f_y is $\mathcal{J}_{\psi|L^\varphi}$ -continuous, so f_y is \mathcal{J}_P^φ -continuous, because $\mathcal{J}_{\psi|L^\varphi} \subset \mathcal{J}_P^\varphi$.

4. THE IDENTITY OF THE TOPOLOGIES \mathcal{J}_I^φ AND \mathcal{J}_P^φ ON L^φ

In this section we shall prove that the topologies \mathcal{J}_I^φ and \mathcal{J}_P^φ coincide on L^φ . We start with the following lemma.

Lemma 4.1. *Let $\varphi \in \Phi_{12}$ and ψ be a φ -function. Then the following statements hold:*

(i) *For every $r > 0$ and $\varepsilon > 0$ there exists $\delta > 0$ such that*

$$\sup \{ \|x_A\|_\psi : x \in B_\varphi(r) \} < \varepsilon \quad \text{for } A \in \Sigma, \mu(A) < \delta.$$

(ii) *If $\psi \not\prec \varphi$, then for every $r > 0$*

$$\mathcal{J}_{\mu|B_\varphi(r)} = \mathcal{J}_{\psi|B_\varphi(r)}.$$

Proof. (i) Assume that $\varphi(u) < \infty$ for $0 \leq u \leq a$ and $\varphi(u) = \infty$ for $u > a$, where $a > 0$. Given $x \in B_\varphi(r)$ we have $\int_\Omega \varphi(|x(t)|/r) d\mu \leq r$, so $|x(t)|/r \leq a$ a.e. on Ω . Given $\varepsilon > 0$ let $\delta = \varepsilon/\psi(ar/\varepsilon)$. Then for $A \in \Sigma$ with $\mu(A) < \delta$

$$\int_\Omega \psi(|x_A(t)|/\varepsilon) d\mu \leq \psi(ar/\varepsilon)\mu(A) \leq \varepsilon$$

i.e., $\|x_A\|_\psi \leq \varepsilon$.

(ii) Since the inclusion $\mathcal{J}_{\mu|L^\varphi} \subset \mathcal{J}_{\psi|L^\varphi}$ holds it is enough to show that $\mathcal{J}_{\psi|B_\varphi(r)} \subset \mathcal{J}_{\mu|B_\varphi(r)}$ holds for every $r > 0$. To this end we shall show that for any $x \in B_\varphi(r)$ and $\xi > 0$ there exists $\eta_0 > 0$ such that

$$B_\mu(x, \eta_0) \cap B_\varphi(r) \subset B_\psi(x, \xi)$$

where for $\eta > 0$

$$\begin{aligned} B_\mu(x, \eta) &= \{y \in L^{\varphi_0} : \|y - x\|_\mu \leq \eta\} \\ &= \{y \in L^{\varphi_0} : \mu(\{t \in \Omega : |y(t) - x(t)| > \eta\}) \leq \eta\} \end{aligned}$$

and

$$B_\psi(x, \eta) = \{y \in L^\psi : \|y - x\|_\psi \leq \eta\}.$$

Indeed, let $x \in B_\varphi(r)$ and $\xi > 0$ be given. For $\eta > 0$ and $y \in B_\varphi(r)$ let put

$$E(\eta, y) = \{t \in \Omega : |y(t) - x(t)| > \eta\}, \quad G(\eta, y) = \Omega \setminus E(\eta, y).$$

It is seen that

$$m_\varphi((y - x)/2r) \leq 2r. \quad (1)$$

Since $\psi \stackrel{s}{\prec} \varphi$, there exists $u_0 > 0$ such that

$$\psi\left(\frac{2u}{\xi}\right) \leq \frac{\xi}{4r} \varphi\left(\frac{u}{2r}\right) \quad \text{for } 0 \leq u \leq u_0. \quad (2)$$

Moreover, in view of (i) there exists $\delta > 0$ such that

$$\|(y - x)_A\|_\psi \leq \frac{1}{2}\xi \quad \text{for } A \in \Sigma \text{ with } \mu(A) < \delta. \quad (3)$$

Now let $\eta_0 = \min(u_0, \delta)$ and let $y \in B_\mu(x, \eta_0) \cap B_\varphi(r)$. Then $\mu(E(\eta_0, y)) \leq \eta_0 \leq \delta$, and hence from (3) we get

$$\|(y - x)_{E(\eta_0, y)}\|_\psi \leq \frac{1}{2}\xi. \quad (4)$$

On the other hand, since $\eta_0 \leq u_0$, from (2) and (1) we get

$$\begin{aligned} m_\psi\left(\frac{2}{\xi}(y-x)_{G(\eta_0,y)}\right) &= \int_{G(\eta_0,y)} \psi\left(\frac{2|y(t)-x(t)|}{\xi}\right) d\mu \\ &\leq \frac{\xi}{4r} \int_{\Omega} \varphi\left(\frac{|y(t)-x(t)|}{2r}\right) d\mu = \frac{\xi}{2}. \end{aligned}$$

Hence

$$\|(y-x)_{G(\eta_0,y)}\|_\psi \leq \frac{\xi}{2}. \quad (5)$$

Thus from (1), (4) and (5) we get

$$\|y-x\|_\psi \leq \|(y-x)_{E(\eta_0,y)}\|_\psi + \|(y-x)_{G(\eta_0,y)}\|_\psi \leq \xi$$

and this means that $y \in B_\psi(x, \xi)$.

As an application of Lemma 4.1 we get

Corollary 4.2. *Let $\varphi \in \Phi_1$. Then for every $r > 0$*

$$\mathcal{J}_{P|B_\varphi(r)}^\varphi = \mathcal{J}_{\mu|B_\varphi(r)}.$$

Proof. This equality is proved in [14, Theorem 1.4] for φ being a φ -function, but the proof can be applied for $\varphi \in \Phi_{11}$. For $\varphi \in \Phi_{12}$ our equality follows Lemma 4.1, because

$$\mathcal{J}_{P|B_\varphi(r)}^\varphi = \sup\{\mathcal{J}_{\psi|B_\varphi(r)} : \psi \in \Psi_{12}^\varphi\} = \mathcal{J}_{\mu|B_\varphi(r)}.$$

In view of Corollary 4.2 and (2.2) we have: $\mathcal{J}_P^\varphi \subset \mathcal{J}_I^\varphi$. Repeating the arguments of the proof of [14, Theorem 2.2] we get

Theorem 4.3. *Let $\varphi \in \Phi_1$. If a sequence (x_n) in L^φ is modular convergent to $x \in L^\varphi$ (i.e., $m_\varphi(\lambda(x_n - x)) \rightarrow 0$ for some $\lambda > 0$) then $x_n \rightarrow 0$ for \mathcal{J}_I^φ*

It is well known that the set of all simple integrable functions \mathfrak{S} is dense in L^φ in the sense of modular convergence. Therefore, in view of the previous theorem and the inclusion $\mathcal{J}_P^\varphi \subset \mathcal{J}_I^\varphi$ we get:

Theorem 4.4. *Let $\varphi \in \Phi_1$. The set of all simple integrable functions \mathfrak{S} is dense in L^φ with respect to \mathcal{J}_P^φ and \mathcal{J}_I^φ .*

Now we are in position to prove our main theorem.

Theorem 4.5. *Let $\varphi \in \Phi_1$. Then the equality*

$$\mathcal{J}_I^\varphi = \mathcal{J}_P^\varphi$$

holds, i.e., for $\varphi \in \Phi_{1i}$ ($i = 1, 2$) the generalized mixed topology \mathcal{J}_I^φ is generated by the family $\{\|\cdot\|_{\psi|L^\varphi} : \psi \in \Psi_{1i}^\varphi\}$.

Proof. For $\varphi \in \Phi_{11}$ this equality is proved in [14, Theorem 2.4].

Next let $\varphi \in \Phi_{12}$. It is enough to show that the inclusion $\mathcal{J}_I^\varphi \subset \mathcal{J}_P^\varphi$ holds. Since the spaces $(L^\varphi, \mathcal{J}_P^\varphi)$ and $(L^\varphi, \mathcal{J}_I^\varphi)$ are complete (see Theorems 2.1 and 3.9), in view of Theorem 4.4 and [4, Corollary of Lemma 4, p. 34] it suffices to show that

$$\mathcal{J}_{I|\mathfrak{S}}^\varphi \subset \mathcal{J}_{P|\mathfrak{S}}^\varphi.$$

To his end, in view of Definition 2.1, given a sequence of positive numbers $(\varepsilon_n : n \geq 0)$ we shall find $\psi_0 \in \Psi_{12}^\varphi$ (i.e., $\psi_0 \not\prec \varphi$) and $r_0 > 0$ such that

$$B_{\psi_0}(r_0) \cap \mathfrak{S} \subset \bigcup_{N=0}^{\infty} \left(\sum_{n=0}^N (B_\varphi(2^n) \cap B_\mu(\varepsilon_n)) \right). \quad (1)$$

Without loss of generality we can assume that $\varepsilon_n \downarrow 0$, $\varepsilon_0 < 1$ and $\varepsilon_0 \varphi(1) < 1$. Moreover, for the reasons of convenience we can assume that $\varphi(u) < \infty$ for $u \leq 1$ and $\varphi(u) = \infty$ for $u > 1$.

Let us choose subsequence (ε_{k_n}) of (ε_n) in such a way that:

(a) $k_0 = 0$.

(b) k_1 is the smallest natural number such that

$$\varphi(\varepsilon_{k_1}) < \frac{1}{\varepsilon_{k_1}}.$$

(c) Given k_n ($n \geq 1$) we take $k_{n+1} > k_n$ such that

$$\varphi(\varepsilon_{k_n})/2 > \varphi(\varepsilon_{k_{n+1}}) \quad \text{and} \quad \varepsilon_{k_n} < \varepsilon_{n+1}/2.$$

Let

$$N(t) = \sup\{n \in \mathcal{N} : \varepsilon_{k_n} \geq t\} \text{ for } t \in \varepsilon_{k_1}.$$

We shall now define a φ -function ψ_0 such that $\psi_0 \stackrel{s}{\prec} \varphi$ and

$$\psi_0(u) \geq \frac{1}{N(u)} \varphi\left(\frac{u}{N(u)}\right) \text{ for } 0 \leq u \leq \varepsilon_{k_2}, \quad (2)$$

$$\psi_0(n) \geq \frac{1}{\varepsilon_{n+1}} \text{ for } n \geq 1. \quad (3)$$

Let us denote by:

$$A'_n = \{t > 0 : \varepsilon_{k_{n+1}} < t \leq \varepsilon_{k_n}\}, \quad n = 1, 2, \dots,$$

$$A = \{t > 0 : \varepsilon_{k_1} < t < 1\},$$

$$A''_n = \{t > 0 : n \leq t < n + 1\}, \quad n = 1, 2, \dots,$$

and

$$B'_n = \{s > 0 : \varphi(\varepsilon_{k_{n+1}}) < s \leq \varphi(\varepsilon_{k_n})\}, \quad n = 1, 2, \dots,$$

$$B = \{s > 0 : \varphi(\varepsilon_{k_1}) < s < \varphi(1)/2\},$$

$$B''_n = \{s > 0 : \varphi(1) n/2 \leq s < \varphi(1)(n+1)/2\}, \quad n = 1, 2, \dots.$$

Let us put

$$p(t) = \begin{cases} 0 & \text{for } t = 0, \\ \frac{2}{n-1} & \text{for } t \in A'_n, n \geq 2, \\ 2 & \text{for } t \in A'_1 \cup A \cup A''_1, \\ n & \text{for } t \in A''_n, n \geq 2, \end{cases}$$

and

$$q(s) = \begin{cases} 0 & \text{for } s = 0, \\ \frac{2}{n-1} & \text{for } s \in B'_n, n \geq 3 \\ 1 & \text{for } s \in B'_2 \cup B'_1 \cup B, \\ \frac{2}{\varphi(1)\varepsilon_{n+1}} & \text{for } s \in B''_n, n \geq 1. \end{cases}$$

Next, define for $u \geq 0$ and $v \geq 0$

$$\xi(u) = \int_0^u p(t) dt \quad \text{and} \quad \eta(v) = \int_0^v q(s) ds.$$

Let us put

$$\varphi_0(u) = \begin{cases} \varphi(u) & \text{for } u \leq 1 \\ \varphi(1)u & \text{for } u > 1 \end{cases}$$

At last let

$$\psi_0(u) = \eta(\varphi_0(\xi(u))) = \int_0^{\varphi_0(\xi(u))} q(s) ds \quad \text{for } u \geq 0.$$

Similarly as in the proof of [14, Theorem 2.4] we can show that $\psi_0 \overset{s}{\prec} \varphi$ and that the conditions (2) and (3) hold.

Now let us put

$$r_0 = \min\left(\frac{1}{2}, d_0 \varepsilon_0 \varphi(\varepsilon_0)\right) \quad (4)$$

Where

$$\frac{1}{d_0} = \sup\left\{\frac{\varphi(u)}{\psi_0(u)} : \varepsilon_{k_2} < u \leq 1\right\}.$$

We shall now show that the inclusion (1) holds. Indeed, let

$$x = \sum_{i \in I} \lambda_i \chi_{H_i} \in B_{\psi_0}(r_0)$$

where I is a finite subset of \mathcal{N} , and $\mu(H_i) < \infty$. Denote by

$$K = \{i \in I : \varepsilon_{k_2} < |\lambda_i| \leq 1\},$$

$$L = \{i \in I : |\lambda_i| \leq \varepsilon_{k_2}\}, J = \{i \in I : |\lambda_i| > 1\}.$$

Let

$$x_1 = \sum_{i \in K} \lambda_i \chi_{H_i}, \quad x_2 = \sum_{i \in L} \lambda_i \chi_{H_i}, \quad x_3 = \sum_{i \in J} \lambda_i \chi_{H_i}.$$

Since $x \in B_{\psi_0}(r_0)$ and $r_0 < 1$ we have

$$m_{\psi_0}(x) = \sum_{i \in I} \psi_0(|\lambda_i|) \mu(H_i) = c \leq r_0.$$

Write

$$c_i = \psi_0(|\lambda_i|) \mu(H_i).$$

Arguing as in the proof of [14, Theorem 2.4] we get

$$x_1 \in B_\varphi(1) \cap B_\mu(\varepsilon_0) \quad (5)$$

and moreover, using (2) we can find $N_1 \in \mathcal{N}$ such that

$$x_2 \in \sum_{n=1}^{N_1} \left(B_\varphi\left(\frac{1}{2} 2^n\right) \cap B_\mu\left(\frac{1}{2} \varepsilon_n\right) \right). \quad (6)$$

Now we shall find $N_2 \in \mathcal{N}$ such that

$$x_3 \in \sum_{n=1}^{N_2} \left(B_\varphi\left(\frac{1}{2} 2^n\right) \cap B_\mu\left(\frac{1}{2} \varepsilon_n\right) \right). \quad (7)$$

Let

$$n_i = \sup\{n \in \mathcal{N} : n \leq |\lambda_i|\} \text{ for } i \in J.$$

Then $n_i \geq 1$ and $n_i \leq |\lambda_i| < n_i + 1$. Let j_1, \dots, j_{m_0} be the different numbers in the set $\{n_i : i \in J\}$ and let us assume that $j_1 < \dots < j_{m_0}$. Write

$$J_l = \{i \in J : n_i = j_l\} \quad \text{for } 1 \leq l \leq m_0.$$

Then

$$x_3 = \sum_{i \in J} \lambda_i \chi_{H_i} = \sum_{l=1}^{m_0} \left(\sum_{i \in J_l} \lambda_i \chi_{H_i} \right).$$

and let

$$y_l = \sum_{i \in J_l} \lambda_i \chi_{H_i} \text{ for } 1 \leq l \leq m_0.$$

Then $j_l \leq |\lambda_i| < j_l + 1$ for $i \in J_l$ and using (4) we get

$$\begin{aligned} m_\varphi(y_l/(j_l + 1)) &= \sum_{i \in J_l} \varphi(|\lambda_i|/(j_l + 1)) \mu(H_i) \\ &\leq d_0^{-1} \sum_{i \in J_l} \psi_0(|\lambda_i|/(j_l + 1)) \mu(H_i) \leq d_0^{-1} \tau_0 < j_l + 1. \end{aligned}$$

Thus

$$y_l \in B_\varphi(j_l + 1) \subset B_\varphi\left(\frac{1}{2} 2^{j_l+1}\right). \quad (8)$$

Let

$$E_{y_l}(\varepsilon) = \{t \in \Omega : |y_l(t)| > \varepsilon\} \text{ for any } \varepsilon > 0.$$

Then

$$E_{y_l}\left(\frac{1}{2} \varepsilon_{j_l+1}\right) = \bigcup_{i \in J_l} H_i.$$

Hence, using (3) we get

$$\begin{aligned} \mu(E_{y_l}(\frac{1}{2} \varepsilon_{j_l+1})) &\leq \sum_{i \in J_l} \mu(H_i) \leq \sum_{i \in J_l} c_i / \psi_0(|\lambda_i|) \\ &\leq \sum_{i \in J_l} \frac{c_i / \psi_0(j_l)}{\psi_0(j_l)} \leq \left(\sum_{i \in J_l} c_i \right) / \psi_0(j_l) \leq \frac{1}{2} \varepsilon_{j_l+1} \end{aligned}$$

and this means that

$$y_l \in B_\mu\left(\frac{1}{2} \varepsilon_{j_l+1}\right). \quad (9)$$

Thus from (8) and (9) we have

$$y_l \in B_\varphi\left(\frac{1}{2} 2^{j_l+1}\right) \cap B_\mu\left(\frac{1}{2} \varepsilon_{j_l+1}\right).$$

Hence for $N_2 = j_{m_0} + 1$ we obtain

$$x_3 \in \sum_{l=1}^{m_0} \left(B_\varphi\left(\frac{1}{2} 2^{j_l+1}\right) \cap B_\mu\left(\frac{1}{2} \varepsilon_{j_l+1}\right) \right) \subset \sum_{n=1}^{N_2} \left(B_\varphi\left(\frac{1}{2} 2^n\right) \cap B_\mu\left(\frac{1}{2} \varepsilon_n\right) \right).$$

At last, using (5), (6) and (7), for $N_0 = \max(N_1, N_2)$ we get

$$\begin{aligned} x = x_1 + x_2 + x_3 &\in B_\varphi(1) \cap B_\mu(\varepsilon_0) + \sum_{n=1}^{N_0} \left(B_\varphi(2^n) \cap B_\mu(\varepsilon_n) \right) \\ &\subset \bigcup_{N=0}^{\infty} \left(\sum_{n=0}^N \left(B_\varphi(2^n) \cap B_\mu(\varepsilon_n) \right) \right). \end{aligned}$$

Thus the proof is completed.

5. A TOPOLOGICAL CHARACTERIZATION OF THE γ_φ -CONVERGENCE IN L^φ

In this section by applying of Theorem 4.5 we obtain a topological characterization of the γ_φ -convergence in L^φ .

Theorem 5.1. *Let $\varphi \in \Phi_{1i}$ ($i = 1, 2$). Then for a sequence (x_n) in L^φ and $x \in L^\varphi$ the following statements are equivalent:*

- (i) $x_n \rightarrow x$ for \mathcal{J}_i^φ .
- (ii) $\|x_n - x\|_\psi \rightarrow 0$ for every $\psi \in \Psi_{1i}^\varphi$.
- (iii) $x_n \rightarrow x$ ($\mu - \Omega$) and $\sup_n \|x_n\|_\varphi < \infty$.

Moreover, for $\varphi \in \Phi_{1i}^c$ ($i = 1, 2, 3, 4$) the above statements are equivalent to

- (iv) $\| \|x_n - x\| \|_\psi \rightarrow 0$ for every $\psi \in \Psi_{1i}^\varphi(N)$.

Proof. (i) \Leftrightarrow (ii) \Leftrightarrow (iii). It follows from Theorem 2.4. and Theorem 4.5.

(i) \Leftrightarrow (iv). It follows from Theorem 4.5 and Theorem 3.12.

Now we apply the above theorem to some classes of Orlicz spaces. Let

$$\chi_p(u) = u^p \text{ for } u \geq 0 \text{ and } \chi_\infty(u) = \begin{cases} 0 & \text{for } 0 \leq u \leq 1, \\ \infty & \text{for } u > 1. \end{cases}$$

Let $\|\cdot\|_p$ and $\|\cdot\|_\infty$ denote the usual norms in L^p ($p \geq 1$) and L^∞ respectively.

Examples

A. For $p \geq 1$ let

$$\varphi(u) = \begin{cases} u^p & \text{for } 0 \leq u \leq 1, \\ \infty & \text{for } u > 1. \end{cases}$$

Hence $\varphi(u) = \chi_p(u) \vee \chi_\infty(u)$ for $u \geq 0$, so $L^\varphi = L^p \cap L^\infty$ by Lemma 1.1. We see that $\varphi \in \Phi_{14}^c$ for $p = 1$ and $\varphi \in \Phi_{12}^c$ for $p > 1$. Hence by applying of Theorem 5.1 and Lemma 1.1 we get the following two theorems:

Theorem 5.2. *For a sequence (x_n) in $L^1 \cap L^\infty$ and $x \in L^1 \cap L^\infty$ the following statements are equivalent:*

$$(i) \ x_n \rightarrow x \ (\mu - \Omega) \text{ and } \sup_n \|x_n\|_1 < \infty, \sup_n \|x_n\|_\infty < \infty.$$

$$(ii) \ |||x_n - x|||_\psi \rightarrow 0 \text{ for every } N\text{-function } \psi.$$

Theorem 5.3. *Let $p > 1$. For a sequence (x_n) in $L^p \cap L^\infty$ and $x \in L^p \cap L^\infty$ the following statements are equivalent:*

$$(i) \ x_n \rightarrow x \ (\mu - \Omega) \text{ and } \sup_n \|x_n\|_p < \infty, \sup_n \|x_n\|_\infty < \infty.$$

(ii) $|||x_n - x|||_\psi \rightarrow 0$ for every N -function ψ such that $\psi(u)/u^p \rightarrow 0$ as $u \rightarrow 0$.

B. For $p > 1$ let

$$\varphi(u) = \begin{cases} u & \text{for } 0 \leq u \leq 1, \\ u^p & \text{for } u > 1. \end{cases}$$

Then $\varphi(u) = \chi_1(u) \vee \chi_p(u)$ for $u \geq 0$, so $L^\varphi = L^1 \cap L^p$. Then $\varphi \in \Phi_{13}^c$ and using Theorem 5.1 and Lemma 1.1 we get:

Theorem 5.4. *Let $p > 1$. For a sequence (x_n) in $L^1 \cap L^p$ and $x \in L^1 \cap L^p$ the following statements are equivalent:*

$$(i) \ x_n \rightarrow x \ (\mu - \Omega) \text{ and } \sup_n \|x_n\|_1 < \infty, \sup_n \|x_n\|_p < \infty.$$

(ii) $|||x_n - x|||_\psi \rightarrow 0$ for every N -function ψ such that $\psi(u)/u^p \rightarrow 0$ as $u \rightarrow \infty$.

References

- [1] Alexiewicz, A., *On two norm convergence*, Studia Math., 14 (1954), 49-56.

- [2] Alexiewicz, A., Semadeni, Z. *A generalization of two-norm spaces, Linear functionals*, Bull. Acad. Polon. Sci., 6 (1958), 135-139.
- [3] Cooper, J.B., *Saks spaces and applications to functional analysis*, North-Holland, Math. Studies 139, 1987.
- [4] Cristescu, R., *Topological vector spaces*, Leyden 1977.
- [5] Davis, H.W., Murray, F.J. and Weber, J.K., *Families of L_p -spaces with inductive and projective topologies*, Pacific J. Math., 34 (1970), 619-638.
- [6] Davis, H.W., Murray, F.J. and Weber, J.K., *Inductive and projective limits of L_p -spaces*, Portugal. Math., 38 (1972), 21-29.
- [7] Garling, D.J.H., *A generalized form of inductive-limit topology for vector spaces*, Proc. London Math. Soc., (3) 14 (1964). 1-28.
- [8] Hudzik, H., *Intersections and algebraic sums of Musielak-Orlicz spaces*, Portugal. Math., 40 (1985), 287-296.
- [9] Krasnosel'skii, M.A. and Rutickii, Ya., *Convex functions and Orlicz spaces*, P. Noordhoff, Groningen 1961.
- [10] Lesniewicz, R., *On two equalities for Orlicz spaces*, Bull. Acad. Polon. Sci., 27 (1979), 557-560.
- [11] Luxemburg, W.A., *Banach function spaces*, Delft 1955.
- [12] Matuszewska, W., Orlicz, W., *A note on the theory of s -normed spaces of φ -integrable functions*, Studia Math., 21 (1961), 107-115.
- [13] Nowak, M., *On two linear topologies on Orlicz spaces $L^{*\varphi}, I$* , Comment. Math., 23 (1983), 71-84.
- [14] Nowak, M., *Inductive limit of a sequence of balanced topological spaces in Orlicz spaces $L_E^{*\varphi}(\mu)$* , Comment. Math., 25 (1985), 295-313.
- [15] Nowak, M., *On some linear topology in Orlicz spaces $L_E^{*\varphi}(\mu), I$* , Comment. Math., 26 (1986), 51-68.
- [16] Nowak, M., *On some linear topology in Orlicz spaces $L_E^{*\varphi}(\mu), II$* , Comment. Math., 26 (1986), 69-74.
- [17] Nowak, M., *Some equalities among Orlicz spaces I*, Comment. Math., 29 (1990), 255-275.

- [18] Nowak, M., *Some equalities among Orlicz spaces II*, Bull. Acad. Polon. Sci., 34 (1986), 675-687.
- [19] Orlicz, W., *Linear operations in Saks spaces I*, Studia Math., 11, (1950), 237-272.
- [20] Orlicz, W., *A note on modular spaces I*, Bull. Acad. Polon. Sci., 9 (1961), 157-162.
- [21] Persson, A., *A generalization of two-norm spaces*, Ark. Math., 5 (1963), 27-36.
- [22] Turpin, P., *Convexités dans les espaces vectoriels topologiques généraux*, Dissertationes Math., 131 (1976).
- [23] Wiweger, A. *A topologization of Saks spaces*, Bull. Acad. Polon. Sci., 5 (1957), 773-777.
- [24] Wiweger, A. *Linear spaces with mixed topology*, Studia Math., 20 (1961), 47-68.
- [25] Wiweger, A. *Some applications of the mixed topology to the theory of two-norm spaces*, Bull. Acad. Polon. Sci., 9 (1961), 571-574.

Institute of Mathematics,
Adam Mickiewicz University
Matejki 48/49,
60-769 Poznan,
Poland.

Recibido: 9 de octubre de 1992