

Shape Theory of Maps

Zvonko ČERIN

ABSTRACT. We shall describe a modification of homotopy theory of maps which we call shape theory of maps. This is accomplished by constructing the shape category of maps \mathcal{HM}^b . The category \mathcal{HM}^b is built using multi-valued functions. Its objects are maps of topological spaces while its morphisms are homotopy classes of collections of pairs of multi-valued functions which we call multi-binets. Various authors have previously given other descriptions of shape categories of maps. Our description is intrinsic in the sense that we do not use any outside objects. It is a version of the author's extension to arbitrary topological spaces of Sanjurjo's approach to shape theory via small multi-valued functions adapted to maps.

1. INTRODUCTION

It is quite often in topology that a concept that was introduced for spaces is later also considered for maps. An example relevant to this paper is provided by the classical homotopy category \mathcal{H} on the one hand and the homotopy category of maps \mathcal{H}^b from E. H. Spanier's book on algebraic topology [10] on the other. The objects of \mathcal{H} are topological

spaces and morphisms are homotopy classes of maps between them while the objects of \mathcal{H}^b are maps of topological spaces and morphisms are homotopy classes of map pairs between them. In this way we can do homotopy theory of maps and talk about contractible maps, homotopy type of maps, and other notions from homotopy theory transferring them from spaces to maps.

The classical homotopy theory has been modified by the introduction of shape theory. The modification was invented by K. Borsuk [3] with the desire to handle more successfully spaces with bad local properties. The new improved homotopy theory that he named shape theory agrees with the old on spaces with nice local properties, for example on the absolute neighbourhood retracts. The key idea in Borsuk's approach was to replace homotopy classes of maps with homotopy classes of sequences of maps that he calls fundamental sequences.

The aim of this paper is to define shape theory for maps of topological spaces. In other words, we shall describe a shape category of maps \mathcal{HM}^b whose objects are maps of topological spaces and whose morphisms will be homotopy classes of pairs of collections of multi-valued functions that we call multi-binets. Therefore, our approach can be regarded as a version for maps of the author's description [4] of the shape category \mathcal{Sh} which is an extension of Sanjurjo's method [9] from compact metric spaces to arbitrary topological spaces.

Our construction extends shape theory from spaces to maps because the full subcategory of \mathcal{HM}^b with maps of topological spaces into a single-element space as objects is naturally isomorphic to the shape category \mathcal{Sh} [8].

The following features of our description of shape category of maps deserve to be emphasized. It is intrinsic in the sense that we do not use any outside objects (like ANR's, embeddings, resolutions, and expansions). It is simple and geometric because it is modelled on the original Borsuk's description of shape theory of compact metric spaces [3]. The only difficult part is in defining the composition of morphisms.

There are already two previous attempts to define shape theory of maps in the same generality as ours. In [6], David Edwards and Patricia McAuley use both Čech and Vietoris constructions to approximate a map $f : X \rightarrow Y$ by a tower of maps between CW-complexes. Each map

gives in this way an object of a suitable pro-category. The study of the shape properties of a map is thus reduced to the study of this associated pro-object. This approach is useful because one can apply homotopy invariant functors (like homology, cohomology, and homotopy groups functors) to get various pro-groups which reflect the geometry of the map.

The second attempt is similar and is due to V.H. Baladze. In [2], he exhibits an outline without any proofs of a shape theory of maps of arbitrary topological spaces following the inverse limit approach to shape theory based on the method of ANR-resolutions, i.e., resolutions of spaces consisting of absolute neighbourhood retracts. In this way a greater flexibility is achieved because we are no longer forced to use Čech and Vietoris constructions that are plagued by certain inconveniences. However, it is extremely complicated to find an $\text{ANR}(M_{maps})$ -expansion for a map, which is a prerequisite for a study of shape properties of the map by this method. Both [6] and [2] belong to the so called outer shape theory in which shape of spaces and maps is investigated with the help of some outside objects like expansions into absolute neighbourhood retracts.

The natural questions of relationship of our shape theory of maps with those mentioned above are deferred to another paper. As an illustration of our method, we shall discuss the problem of identifying maps on which shape theory of maps and homotopy theory of maps coincide. Some interesting classes of maps which we call internally movable, internally calm, and calm provide an unexpected solution to this question. We also give some applications of our approach to identify equivalences in the shape category of maps and maps with trivial shape.

Here is a brief outline of our organization. The principal goal of the paper is to describe with all details the category \mathcal{HM}^b . Homotopy theory of maps which we shall be improving is recalled first. Since our method relies on the idea of approximating a map by a generalized sequence (net over a cofinite directed set) of multi-valued functions with smaller and smaller images of points, we present facts on normal covers [1] which we use as a mean of measuring size of sets and from which our sets of indices for these nets are build. Connection between indexing sets of our nets provide increasing functions which are normally constructed by a well-known simple Lemma 1 (see [8, p. 9]).

Then we define multi-valued functions and multi-valued σ -functions (i.e. a multi-valued function of size less than a normal cover σ of the codomain). Our multi-valued functions must satisfy only the condition that they associate a non-empty subset of the codomain to every point of the domain.

The next step is to define all of these notions for maps. This is done by considering pairs and replacing exact commutativity with an approximate form of it up to a normal cover in the codomain. Since all concepts dealing with maps usually involve pairs we take the convention that they include prefix "bi" in their names. Thus maps themselves are also called bispaces and we define notions of bimap, multi-valued bifunction, normal bicover, and others. The multi-valued σ -functions for maps (or multi-valued σ -bifunctions) lead naturally to the relation of σ -bihomotopy. The important approximate transitivity of this relation is established in Lemma 2. In the first reading one can skip its tedious proof with many cases provided one is familiar with Dold's powerful lemma (see [5, p. 358]) on which our argument depends and which is used later over and over again.

After these preliminaries, we follow closely Borsuk's method from [3]. The role of fundamental sequences play multi-binets, i.e., multi-nets for maps. We define the relation of bihomotopy for multi-binets and take as objects of our category \mathcal{HM}^b maps and as morphisms bihomotopy classes of multi-binets. In order to prove that this is indeed a category we must describe how one composes morphisms and one must prove that this composition is associative. This is done in Claims 1, 2, and 3 and Theorem 1. The details are quite easy once the method of proof for relations (4) and (5) is understood. It would be unreal to expect that in the complete generality which we consider here one can get the shape category of maps without any effort.

The rest of the paper presents some applications of our approach to shape theory of maps. They are included with intention to convince the reader that the new method will be useful for some kind of problems. We intend to undertake further study of the category \mathcal{HM}^b . This paper is only the beginning and it gives basic facts about this approach to shape theory for maps.

2. HOMOTOPY THEORY OF MAPS

Since shape theory is just a modification of homotopy theory, we begin by recalling some notions from homotopy theory of maps.

We follow the convention that a map is a short name for a continuous single-valued function. In order to get closer analogies in our constructions, we shall also use *bispace* as another name for a map. More formally, a *bispace* is a triple (f, F', F) consisting of topological spaces F' and F and a continuous single-valued function $f : F' \rightarrow F$. Since we shall scrupulously follow the notation that a map is denoted by a small letter, it's domain and codomain with the same capital letter, and all domains have in addition a prime, we shall drop F' and F from our notation and use f to denote a bispace (f, F', F) . However, more important for our exposition will be the fact that a presence of a prefix "bi" and either a superscript or a subscript "b" indicates something referring to maps. Thus, x, X^b , and bispace X are all notations for the map (x, X', X) . With these convention enforced consistently we shall see that most results from [4] carry over to maps. In all statements and proofs in [4] we must only put bi and b at proper places to get shape theory for maps.

Following [10, p. 414], we now consider the category S^b whose objects are bispaces (i.e., maps $f : F' \rightarrow F$ between topological spaces) and whose morphisms will be called bimaps. Let F and G be bispaces. By a *bimap* from F into G we mean a pair (a', a) of maps $a' : F' \rightarrow G'$ and $a : F \rightarrow G$ such that $g \circ a' = a \circ f$. In other words we require that the following diagram commutes.

$$\begin{array}{ccc}
 & a' & \\
 F' & \longrightarrow & G' \\
 f \downarrow & & \downarrow g \\
 F & \longrightarrow & G \\
 & a &
 \end{array}$$

Just as with bispaces, we shall also simplify our notation for bispaces. Thus "... bimap $a : F \rightarrow G$..." means that we have a bimap (a', a) between bispaces F and G . The composition of bispaces (a', a) and (b', b)

is a bimap $(b' \circ a', b \circ a)$ and the identity bimap 1_X on a bispace X is a bimap $(1_{X'}, 1_X)$, where 1_X is the identity map on the space X .

The definition of bihomotopy involves the *bicylinder* $F \times I$ of a bispace F . Let I denote the unit closed segment and let I denote also the bispace $(1_I, I, I)$, where $1_I : I \rightarrow I$ is the identity map. The bicylinder is a special case of the biproduct defined for bispaces F and G to be the bispace $(f \times g, F' \times G', F \times G)$.

Let p and q be bimaps between bispaces X and Y . We shall say that p and q are *bihomotopic* and write $p \simeq_b q$ provided there is a bimap $m : X \times I \rightarrow Y$ called *bihomotopy* such that $m'(a', 0) = p'(a')$ and $m'(a', 1) = q'(a')$ for every $a' \in X'$ and $m(a, 0) = p(a)$ and $m(a, 1) = q(a)$ for every $a \in X$. The relation of bihomotopy is an equivalence relation and we denote the bihomotopy class of a bimap p by $[p]_b$. Since the bihomotopy relation \simeq_b is also compatible with the composition of bimaps, one can define the composition of bihomotopy classes of bimaps by composing representatives, i.e., $[p]_b \circ [q]_b = [p \circ q]_b$. In this way one obtains the bihomotopy category \mathcal{H}^b , whose objects are topological bispaces and whose morphisms are bihomotopy classes of bimaps. There is a bihomotopy functor H^b from S^b to \mathcal{H}^b which keeps the objects fixed and takes a bimap p into its bihomotopy class $[p]_b$.

3. NORMAL COVERS AND MULTI-VALUED FUNCTIONS

In this section we shall introduce notions and results on normal covers and multi-valued functions that are required for our theory.

Let $Cov(Y)$ denote the collection of all normal covers of a topological space Y [1]. With respect to the refinement relation \geq the set $Cov(Y)$ is a directed set. Two normal covers σ and τ of Y are equivalent provided $\sigma \geq \tau$ and $\tau \geq \sigma$. In order to simplify our notation we denote a normal cover and its equivalence class by the same symbol. Consequently, $Cov(Y)$ also stands for the associated quotient set.

If σ is a normal cover of a space Y , let σ^+ be the collection of all normal covers of Y which refine σ while σ^* denotes the set of all normal covers τ of Y such that the star $st(\tau)$ of τ refines σ . Similarly, for a natural number n , σ^{*n} denotes the set of all normal covers τ of Y such that the n -th star $st^n(\tau)$ of τ refines σ .

Let $Inc(Y)$ denote the collection of all finite subsets c of $Cov(Y)$ which have a unique (with respect to the refinement relation) maximal element which we denote by $[c]$. The notation $Inc(Y)$ comes from “indices of covers”. The set $Inc(Y)$ will be used as indexing set for multi-nets into Y . We consider $Inc(Y)$ ordered by the inclusion relation and regard $Cov(Y)$ as a subset of single-element subsets of $Cov(Y)$. Notice that $Inc(Y)$ is a cofinite directed set.

We shall repeatedly use the following lemma (see [8, p. 9]). Let us agree that an *increasing* function $f : P \rightarrow P$ of a partially ordered set (P, \leq) into itself is a function which satisfies $x \leq f(x)$ for every $x \in P$ and $x \leq y$ in P implies $f(x) \leq f(y)$. In the case when the domain and the codomain of a function f are different, the first requirement is dropped.

Lemma 1. *Let $\{f_1, \dots, f_n\}$ be functions from a cofinite directed set (M, \leq) into a directed set (L, \leq) . Then there is an increasing function $g : M \rightarrow L$ such that $g(x) \geq f_1(x), \dots, f_n(x)$ for every $x \in M$.*

Let X and Y be topological spaces. By a *multi-valued function* $F : X \rightarrow Y$ we mean a rule which associates a non-empty subset $F(x)$ of Y to every point x of X .

For our approach to shape theory the following notion of size for multi-valued functions will play the most important role.

Let $F : X \rightarrow Y$ be a multi-valued function and let $\alpha \in Cov(X)$ and $\sigma \in Cov(Y)$. We shall say that F is a *multi-valued (α, σ) -function* provided for every $A \in \alpha$ there is an $S_A \in \sigma$ with $F(A) \subset S_A$. On the other hand, F is a *multi-valued σ -function* provided there is an $\alpha \in Cov(X)$ such that F is a multi-valued (α, σ) -function.

Also important will be the following concept of closeness for two multi-valued functions.

Let $F, G : X \rightarrow Y$ be multi-valued functions, let $\sigma \in Cov(Y)$, and let $\alpha \in Cov(X)$. We shall say that F and G are (α, σ) -close and we write $F \stackrel{(\alpha, \sigma)}{=} G$ provided for every A in α there is an $S_A \in \sigma$ with $F(A) \cup G(A) \subset S_A$.

4. NORMAL BICOVERS AND MULTI-VALUED BIFUNCTIONS

In this section we shall define versions for maps of some notions from the previous section.

By a *normal bicover* of a bispace X we mean a pair (σ', σ) , where σ' is a normal cover of the space X' and σ is a normal cover of the space X such that σ' refines the cover $x^{-1}(\sigma)$. We shall again make a simplification of our notation by dropping σ' so that normal bicovers are denoted by small Greek letters which name a normal cover of the codomain while the part in the domain space has the prime.

Let $Cov^b(Y)$ denote the collection of all normal bicovers of a bispace Y . We order $Cov^b(Y)$ by the refinement relation \geq defined by $\sigma \geq \tau$ if and only if $\sigma \geq \tau$ and $\sigma' \geq \tau'$ for normal bicovers σ and τ of Y . With respect to the relation \geq the set $Cov^b(Y)$ is a directed set. Two normal bicovers σ and τ of a bispace Y are equivalent provided $\sigma \geq \tau$ and $\tau \geq \sigma$. In order to simplify our notation we denote a normal bicover and its equivalence class by the same symbol. Consequently, $Cov^b(Y)$ also stands for the associated quotient set.

If σ is a normal bicover of a bispace Y , let σ^+ be the collection of all normal bicovers of Y which refine σ while σ^* denotes the set of all normal bicovers τ of Y such that the star $st(\tau)$ of τ refines σ . Here, we define the star $st(\sigma)$ of a normal bicover (σ', σ) as a normal bicover $(st(\sigma'), st(\sigma))$. Similarly, for a natural number n , σ^{*n} denotes the set of all normal bicovers τ of Y such that the n -th star $st^n(\tau)$ of τ refines σ .

Let Y be a bispace. Let $Inc^b(Y)$ denote the collection of all finite subsets c of $Cov^b(Y)$ which have a unique (with respect to the refinement relation) maximal element which we denote by $[c]$. We consider $Inc^b(Y)$ ordered by the inclusion relation and regard $Cov^b(Y)$ as a subset a single-element subsets of $Cov^b(Y)$. Notice that $Inc^b(Y)$ is a cofinite directed set.

For our approach to shape theory of maps the following class of multi-valued bifunctions will play the most important role.

Let X and Y be bispaces. By a *multi-valued bifunction* from X into Y we mean a pair (A', A) of multi-valued functions $A' : X' \rightarrow Y'$ and $A : X \rightarrow Y$. We shall drop A' and use A to denote the multi-valued

bifunction (A', A) . Of course, we shall utilize the functional notation $A : X \rightarrow Y$ to indicate that A is a multi-valued bifunction between bispaces X and Y . The multi-valued bifunctions A and B are composed by the rule $B \circ A = (B' \circ A', B \circ A)$ and multiplied by the rule $A \times B = (A' \times B', A \times B)$.

Let X and Y be bispaces. Let α and β be normal bicovers of X and Y . By a *multi-valued (α, β) -bifunction* from X into Y we mean a multi-valued bifunction $A : X \rightarrow Y$ such that A' is a multi-valued (α', β') -function, A is a multi-valued (α, β) -function and the compositions $y \circ A'$ and $A \circ x$ are (α', β) -close.

On the other hand, A is a *multi-valued β -bifunction* provided there is a normal bicover $\alpha \in \text{Cov}^b(X)$ such that A is a multi-valued (α, β) -bifunction.

5. THE RELATION OF σ -BIHOMOTOPY

Now we are ready to introduce an important notion of σ -bihomotopy for multi-valued bifunctions of bispaces. We shall also prove in Lemma 2 a useful technical result.

Let F and G be multi-valued bifunctions between bispaces X and Y and let σ be a normal bicover of Y . We shall say that F and G are *σ -bihomotopic* and write $F \overset{\sigma}{\simeq}_b G$ provided there is a multi-valued σ -bifunction H from the bicylinder $X \times I$ into Y such that $F'(a') = H'(a', 0)$ and $G'(a') = H'(a', 1)$ for every $a' \in X'$ and $F(a) = H(a, 0)$ and $G(a) = H(a, 1)$ for every $a \in X$. We shall say that H is a *σ -bihomotopy that joins F and G* or that it *realizes* the relation (or σ -bihomotopy) $F \overset{\sigma}{\simeq}_b G$.

The following lemma is crucial because it provides an adequate substitute for the transitivity of the relation of σ -bihomotopy.

Lemma 2. *Let F, G , and H be multi-valued bifunctions between bispaces X and Y . Let σ be a normal bicover of Y and let $\tau \in \sigma^*$. If $F \overset{\tau}{\simeq}_b G$ and $G \overset{\tau}{\simeq}_b H$, then $F \overset{\sigma}{\simeq}_b H$.*

Proof. By assumption there are normal bicovers α and β of the bicylinder $X \times I$, a multi-valued (α, τ) -bifunction $K : X \times I \rightarrow Y$, and a multi-valued (β, τ) -bifunction $L : X \times I \rightarrow Y$ so that $F'(a') =$

$K'(a', 0)$, $G'(a') = K'(a', 1) = L'(a'; 0)$, and $H'(a') = L'(a', 1)$, for every $a' \in X'$, and $F(a) = K(a, 0)$, $G(a) = K(a, 1)$, $G(a) = L(a, 0)$, and $L(a, 1) = H(a)$ for every $a \in X$. Let a normal bicover γ of $X \times I$ be a common refinement of α and β . Observe first that K and L are both multi-valued (γ, τ) -bifunctions. Define a multi-valued bifunction $M : X \times I \rightarrow Y$ by the rule

$$M'(a', t) = \begin{cases} K'(a', 2t), & a' \in X', 0 \leq t \leq 1/2 \\ L'(a', 2t - 1), & a' \in X', 1/2 \leq t \leq 1, \end{cases}$$

and

$$M(a, t) = \begin{cases} K(a, 2t), & a \in X, 0 \leq t \leq 1/2 \\ L(a, 2t - 1), & a \in X, 1/2 \leq t \leq 1, \end{cases}$$

Since $F'(a') = M'(a', 0)$ and $H'(a') = M'(a', 1)$ for every $a' \in X'$, and $F(a) = M(a, 0)$ and $H(a) = M(a, 1)$ for every $a \in X$, it remains to see that M is a multi-valued σ -bifunction.

By [5, p. 358], there are normal covers δ' and δ of X' and X and functions $\tau' : \delta' \rightarrow \{2, 3, 4, \dots\}$ and $\tau : \delta \rightarrow \{2, 3, 4, \dots\}$ such that the set $D' \times D'_i$ is contained in a member $C'_{D', i}$ of γ' for every $D' \in \delta'$ and every $i = 1, \dots, \tau' D' - 1$, where $D'_i = [\frac{i-1}{\tau' D'}, \frac{i+1}{\tau' D'}]$ ($i = 1, \dots, \tau' D' - 1$) and the set $D \times D_i$ is contained in a member $C_{D, i}$ of γ for every $D \in \delta$ and every $i = 1, \dots, \tau D - 1$, where $D_i = [\frac{i-1}{\tau D}, \frac{i+1}{\tau D}]$ ($i = 1, \dots, \tau D - 1$).

We can assume that δ' refines $x^{-1}(\delta)$. Then for every $D' \in \delta'$ there is a $D(D') \in \delta$ such that D' is a subset of $x^{-1}(D(D'))$. Let $sD' = (\tau' D')(\tau D(D'))$ for every $D' \in \delta'$ and let

$$E_i = \begin{cases} [0, \frac{1}{2\tau D}], & i = 0, \\ (\frac{i}{4\tau D}, \frac{i+2}{4\tau D}), & i = 1, \dots, 4\tau D - 3, \\ (1 - \frac{1}{2\tau D}, 1], & i = 4\tau D - 2, \end{cases}$$

and similarly

$$E'_i = \begin{cases} \left[0, \frac{1}{2sD'}\right), & i = 0, \\ \left(\frac{i}{4sD'}, \frac{i+2}{2sD'}\right), & i = 1, \dots, 4sD' - 3, \\ \left(1 - \frac{1}{2sD'}, 1\right], & i = 4sD' - 2. \end{cases}$$

The collections

$$\varepsilon' = \{D' \times E'_i \mid D' \in \delta', i = 0, \dots, 4sD' - 2\}$$

and

$$\varepsilon = \{D \times E_i \mid D \in \delta, i = 0, 1, \dots, 4rD - 2\}$$

are normal covers of $X' \times I$ and $X \times I$. Moreover, we claim that the pair $(\varepsilon', \varepsilon)$ is a normal bicover of the bicylinder $X \times I$.

Indeed, let $T = D' \times E'_i$ be a member of ε' . Since the set $R = D \times E$, where $D = D(D')$ and $E = E_j$, is a member of ε for every $j = 0, 1, \dots, 4rD - 2$, the preimage $(x \times 1_I)^{-1}(R)$ is a set $x^{-1}(D) \times E$, and $D' \subset x^{-1}(D)$, it suffices to show that there exists an index j such that $E'_i \subset E_j$.

When $i = 0$, then we can take $j = 0$ because

$$E'_0 = \left[0, \frac{1}{2sD'}\right) = \left[0, \frac{1}{2r'D'rD(D')}\right) \subset \left[0, \frac{1}{2rD}\right) = E_0.$$

For a similar reason, when $i = 4sD' - 2$, then we can take $j = 4rD - 2$. Finally, when $i \in \{1, \dots, 4sD' - 3\}$, the required inclusion will hold provided $\frac{i}{4sD'} \geq \frac{j}{4rD}$ and $\frac{i+2}{4sD'} \leq \frac{j+2}{4rD}$. Thus, the index j must satisfy conditions $\frac{i+2}{r'D'} - 2 \leq j \leq \frac{i}{r'D'}$. But, if we write $i = r'D'm + n$, for integers m and n with $0 \leq n < r'D'$, then these conditions become $j \geq m - 2 + \frac{n+2}{r'D'}$ and $j \leq m + \frac{n}{r'D'}$. It is clear that $j = m$ always satisfies them. Another possibility is $j = m - 1$ in some situations. Therefore, $E'_i \subset E_m$ and our claim has been verified.

In order to prove that M is a multi-valued σ -bifunction, we shall show that it is a multi-valued (ε, σ) -bifunction. In other words, we shall prove that

- (1) M' is a multi-valued (ε', σ') -function,
- (2) M is a multi-valued (ε, σ) -function, and
- (3) the compositions $y \circ (M')$ and $M \circ (x \times 1_I)$ are (ε', σ) -close.

Add (1). The argument for (1) is similar to the argument for (2) given below. Of course, the case (2) is notationally simpler because there are no primes.

Add (2). Let $E = D \times E_i$ be a member of ε . We must find an $S_E \in \sigma$ such that $M(E) \subset S_E$.

Case I ($i = 2k$ for $0 \leq k \leq \tau D - 1$). Then $E_i = E_{2k}$ is below $1/2$ so that we get $M(E) = K(D \times 2E_i) = K(D \times D_k) \subset K(C_{D,k}) \subset T \subset S_E$, where the open set T is a member of τ which we obtain with respect to $C_{D,k}$ from the fact that K is a multi-valued (γ, τ) -function and S_E is a member of σ which contains T .

Case II ($i = 2k$ for $\tau D \leq k \leq 2\tau D - 1$). Then $E_i = E_{2k}$ is above $1/2$ so that we get $M(E) = L(D \times (2E_i - 1)) \subset L(D \times D_{k-\tau D}) \subset L(C_{D,k-\tau D}) \subset T \subset S_E$, where the open set T is a member of τ which we obtain with respect to $C_{D,k-\tau D}$ from the fact that L is a multi-valued (γ, τ) -function, and S_E is a member of σ which contains the set T .

Case III ($i = 2k + 1$ for $0 \leq k \leq \tau D - 2$). Then $E_i = E_{2k+1}$ is below $1/2$ so that we get

$$\begin{aligned} M(E) &= K(D \times 2E_i) \subset K(D \times (D_k \cup D_{k+1})) \\ &\subset K(C_{D,k}) \cup K(C_{D,k+1}) \subset T_1 \cup T_2 \subset S_E, \end{aligned}$$

where the sets T_1 and T_2 are members of τ which we obtain with respect to $C_{D,k}$ and $C_{D,k+1}$ from the fact that K is a (γ, τ) -function, and S_E is a member of σ which contains the union $T_1 \cup T_2$. Such an S_E exists because $D \times \{k/\tau D\} \subset C_{D,k} \cap C_{D,k+1}$ so that $\emptyset \neq K(D \times \{k/\tau D\}) \subset T_1 \cap T_2$.

Case IV ($i = 2k + 1$ for $\tau D \leq k \leq 2\tau D - 1$). This case is analogous to the case III. This time E_i is above $1/2$ and we must use L instead of K .

Case V ($i = 2k + 1$ and $k = \tau D - 1$). Then $E_i = E^{-1} \cup E^+$, where $E^- = (\frac{2\tau D - 1}{4\tau D}, \frac{1}{2}]$ and $E^+ = [\frac{1}{2}, \frac{2\tau D + 1}{4\tau D})$ so that we get

$$M(E) = K(D \times 2E^-) \cup L(D \times (2E^+ - 1)) \subset K(C_{D, \tau D - 1}) \cup L(C_{D, 1}) \\ \subset T_1 \cup T_2,$$

where T_1 and T_2 are members of τ which we obtain with respect to $C_{D, \tau D - 1}$ and $C_{D, 1}$ from the fact that K and L are multi-valued (γ, τ) -functions.

Observe that $K(D \times \{1\}) = G(D) \subset T_1$ and $L(D \times \{0\}) = G(D) \subset T_2$. We conclude that $T_1 \cap T_2 \neq \emptyset$ so that there is a member S_E of σ which contains both T_1 and T_2 . It follows that $M(E) \subset S_E$.

Add (3). Let $E' = D' \times E'_i$ be a member of ε' . We must find a member $S_{E'}$ of σ such that both $M \circ (x \times 1_I)(E')$ and $y \circ (M')(E')$ are contained in the set $S_{E'}$.

Once again we shall distinguish five cases considered above.

Case I ($i = 2k$ for $0 \leq k \leq \tau' D' - 1$). Let A' be a member of α' which contains the set $C'_{D', k}$. Since $K \circ (x \times 1_I)$ and $y \circ (K')$ are (α', τ) -close, there is a member T of τ which contains both $K \circ (x \times 1_I)(A')$ and $y \circ (K')(A')$. Since τ is a star-refinement of σ , there is an $S_E \in \sigma$ with $T \subset S_E$. Our choices imply

$$y \circ (M')(E') \subset y \circ (K')(C'_{D', k}) \subset y \circ (K')(A')$$

and

$$M \circ (x \times 1_I)(E') \subset M(x(D') \times E'_i) = K(x(D') \times 2E'_i) \\ \subset K \circ (x \times 1_I)(D' \times D'_k) \subset K \circ (x \times 1_I)(A')$$

because $E' \subset C'_{D', k} \subset A'$. It follows that $M \circ (x \times 1_I)(E')$ and $y \circ (M')(E')$ are both subsets of $S_{E'}$.

Case II ($i = 2k$ for $\tau' D' \leq k \leq 2\tau' D' - 1$). This is similar to the previous case. We have to deal with L instead of with K .

Case III ($i = 2k + 1$ for $0 \leq k \leq \tau' D' - 2$). The set $y \circ (M')(E')$ is now a subset of the union of sets $y \circ (K')(C'_{D', k})$ and $y \circ (K')(C'_{D', k+1})$.

Let A'_1 and A'_2 be members of α' which contain $C'_{D',k}$ and $C'_{D',k+1}$, respectively. Since $K \circ (x \times 1_I)$ and $y \circ (K')$ are (α', τ) -close, there are members T_1 and T_2 of τ such that T_1 contains $K \circ (x \times 1_I)(A'_1)$ and $y \circ (K')(A'_1)$ while T_2 contains $K \circ (x \times 1_I)(A'_2)$ and $y \circ (K')(A'_2)$. The sets T_1 and T_2 both contain the set $y \circ (K')(D' \times \{\frac{k}{r'D'}\})$. It follows that some member $S_{E'}$ of σ contains their union. This is the required open set.

Case IV. This is similar to the previous case.

Case V. The set $y \circ (M')(E')$ is a subset of the union of sets $y \circ (K')(C'_{D',r'D'-1})$ and $y \circ (L')(C'_{D',1})$. Let A'_1 be a member of α' and let B'_2 be a member of β' such that A'_1 contains $C'_{D',r'D'-1}$ and B'_2 contains $C'_{D',1}$. Since $K \circ (x \times 1_I)$ is (α', τ) -close to $y \circ (K')$ and $L \circ (x \times 1_I)$ is (β', τ) -close to $y \circ (L')$, there are members T_1 and T_2 of τ such that T_1 contains $K \circ (x \times 1_I)(A'_1)$ and $y \circ (K')(A'_1)$ while T_2 contains $L \circ (x \times 1_I)(B'_2)$ and $y \circ (L')(B'_2)$. But, the sets T_1 and T_2 both contain the non-empty set

$$y \circ (K')(D' \times \{1\}) = y \circ (G')(D') = y \circ (L')(D' \times \{0\}),$$

so that some member $S_{E'}$ of σ contains their union. It follows that $M \circ (x \times 1_I)(E')$ and $y \circ (M')(E')$ are both subsets of $S_{E'}$. ■

Since we shall be using [5, p. 358] quite often, for a space Y and a normal cover α of the product $Y \times I$, we let $D(Y, \alpha)$ denote all normal covers β of Y such that some stacked normal cover over β refines α .

6. MULTI-BINETES

The following two definitions correspond to Borsuk's definitions of fundamental sequence and homotopy for fundamental sequences [3].

Let X and Y be bispaces. By a *multi-binnet* from X into Y we shall mean a collection $\varphi = \{F_c \mid c \in Inc^b(Y)\}$ of multi-valued bifunctions $F_c : X \rightarrow Y$ such that for every $\sigma \in Cov^b(Y)$ there is a $c \in Inc^b(Y)$ with $F_d \simeq_b^\sigma F_c$ for every $d \geq c$. We use functional notation $\varphi : X \rightarrow Y$ to indicate that φ is a multi-binnet from X into Y . Let $M^b(X, Y)$ denote all multi-binnets $\varphi : X \rightarrow Y$ from X into Y .

Multi-binnets $\varphi = \{F_c\}$ and $\psi = \{G_c\}$ between bispaces X and Y are said to be *bihomotopic* and we write $\varphi \simeq_b \psi$ provided for every $\sigma \in Cov^b(Y)$ there is a $c \in Inc^b(Y)$ such that $F_d \simeq_b^\sigma G_d$ for every $d \geq c$.

It follows from Lemma 2 that the relation of bihomotopy is an equivalence relation on the set $M^b(X, Y)$. The bihomotopy class of a multi-binet φ is denoted by $[\varphi]_b$ and the set of all bihomotopy classes by $\mathcal{H}M^b(X, Y)$.

7. COMPOSITION OF BIHOMOTOPY CLASSES OF MULTI-BINETS

Our principal goal now is to define a composition for bihomotopy classes of multi-binets and to establish its associativity. This is the only tricky part in setting up the category $\mathcal{H}M^b$. Our idea is to associate to every multi-binet $\varphi : X \rightarrow Y$ two increasing functions $\varphi : Inc^b(Y) \rightarrow Inc^b(Y)$ and $\varphi^* : Inc^b(Y) \rightarrow Inc^b(X)$. The first function associates to an index $c \in Inc^b(Y)$ of the family $\varphi = \{F_c\}$ a much larger index $\varphi(c) \in Inc^b(Y)$ such that F_d and F_e are joined by a multi-valued $(\varphi(c, d, e), [c])$ -bihomotopy for a suitable normal bicover $\varphi(c, d, e)$ of the bicylinder $X \times I$ whenever $d, e \geq \varphi(c)$. The second function associates to a $c \in Inc^b(Y)$ an element $\varphi^*(c) \in Inc^b(X)$ such that the normal bicover $[\varphi^*(c)]$ refines some normal bicover of X which we get from the normal bicover $\varphi(c, \varphi(c), \varphi(c))$ by [5, p. 358]. This is a rough description of these functions and now we proceed with the details.

Let X and Y be bispaces. Let $\varphi = \{F_c\}_{c \in Inc^b(Y)} : X \rightarrow Y$ be a multi-binet. Let $\varphi : Inc^b(Y) \rightarrow Inc^b(Y)$ be an increasing function such that for every $c \in Inc^b(Y)$ the relation $d, e \geq \varphi(c)$ implies the relation $F_d \stackrel{[c]}{\simeq}_b F_e$, where $[c]$ denotes the unique maximal element of c .

Let $C_\varphi = \{(c, d, e) \mid c \in Inc^b(Y), d, e \geq \varphi(c)\}$. Then C_φ is a subset of

$$Inc^b(Y) \times Inc^b(Y) \times Inc^b(Y)$$

that becomes a cofinite directed set when we define that $(c, d, e) \geq (c', d', e')$ if and only if $c \geq c'$, $d \geq d'$, and $e \geq e'$ for (c, d, e) and (c', d', e') in C_φ .

We shall use the same notation φ for an increasing function $\varphi : C_\varphi \rightarrow Cov^b(X \times I)$ such that F_d and F_e are joined by a multi-valued $(\varphi(c, d, e), [c])$ -bihomotopy whenever $(c, d, e) \in C_\varphi$.

Let $\bar{\varphi} : C_\varphi \rightarrow Inc^b(X)$ be an increasing function such that the normal cover $[(\bar{\varphi}(c, d, e))']$ belongs to the set $D(X, (\varphi(c, d, e))')$ and the

normal cover $[\bar{\varphi}(c, d, e)]$ belongs to the set $D(X, \varphi(c, d, e))$ for every $(c, d, e) \in C_\varphi$.

Claim 1. *There is an increasing function $\varphi^* : Inc^b(Y) \rightarrow Inc^b(X)$ such that*

- (1) $\varphi^*(c) \geq \bar{\varphi}(c, \varphi(c), \varphi(c))$ for every $c \in Inc^b(Y)$, and
- (2) φ^* is cofinal in $\bar{\varphi}$, i.e., for every $(c, d, e) \in C_\varphi$ there is an $m \in Inc^b(Y)$ with $\varphi^*(m) \geq \bar{\varphi}(c, d, e)$.

Proof. Let $\mathcal{D} = \{\bar{\varphi}(c, d, e) \mid (c, d, e) \in C_\varphi\}$.

If $Inc^b(Y)$ is a finite set, then \mathcal{D} is a finite collection of elements of $Inc^b(X)$. Let $a \in Inc^b(X)$ be greater than all members of \mathcal{D} . Let $\varphi^* : Inc^b(Y) \rightarrow Inc^b(X)$ be a constant function into a .

If $Inc^b(Y)$ is an infinite set, then the cardinality of \mathcal{D} does not exceed the cardinality of $Inc(Y)$. Hence, there is a surjection $g : Inc^b(Y) \rightarrow \mathcal{D}$. Let $\varphi^* : Inc^b(Y) \rightarrow Inc^b(X)$ be an increasing function such that $\varphi^*(c) \geq g(c)$, $\bar{\varphi}(c, \varphi(c), \varphi(c))$ for every $c \in Inc^b(Y)$. ■

The above discussion shows that every multi-binnet $\varphi : X \rightarrow Y$ determines two increasing functions $\varphi : Inc^b(Y) \rightarrow Inc^b(Y)$ and $\varphi^* : Inc^b(Y) \rightarrow Inc^b(X)$. With the help of these functions we shall define the composition of bihomotopy classes of multi-binnets as follows.

Let X, Y , and Z be bispaces and let $\varphi = \{F_c\} : X \rightarrow Y$ and $\psi = \{G_s\} : Y \rightarrow Z$ be multi-binnets. Let $\chi = \{H_s\}$, where $H_s = G_{\psi(s)} \circ F_{\varphi(\psi^*(s))}$ for every $s \in Inc^b(Z)$. Observe that each H_s is a multi-valued bifunction because the composition of two multi-valued bifunctions is a multi-valued bifunction.

Claim 2. *The collection χ is a multi-binnet from X into Z .*

Proof. Let $\sigma \in Cov^b(Z)$. We must find a $u \in Inc^b(Z)$ such that

$$H_v \overset{\sigma}{\simeq}_b H_u \quad \text{for every } v \geq u. \quad (3)$$

Let $\tau \in \sigma^{*2}$ and $\xi \in \tau^*$. Let $u = \{\xi\} \in Inc^b(Z)$.

Consider an index $v \geq u$. We shall find an index $e \in Inc^b(Y)$ so that

$$H_v \overset{\tau}{\simeq}_b G_q \circ F_e, \quad (4)$$

$$G_q \circ F_e \overset{\tau}{\simeq}_b G_p \circ F_e, \quad (5)$$

and

$$G_p \circ F_e \overset{\tau}{\simeq}_b H_u, \quad (6)$$

where $p = \psi(u)$, $q = \psi(v)$, $a = \psi^*(u)$, $b = \psi^*(v)$, $c = \varphi(a)$, and $d = \varphi(b)$. Repeated use of Lemma 2 will give (3) from the relations (4)-(6).

Add (4). Since $(v, q, q) \in \mathcal{C}_\varphi$, we see that there is a multi-valued (α, ξ) -bihomotopy $K : Y \times I \rightarrow Z$ such that $\alpha = \psi(v, y, y)$, $K'_0 = G'_q$, $K'_1 = G'_q$, $K_0 = G_q$, and $K_1 = G_q$. Let $s = \bar{\psi}(v, q, q)$ and $\pi = [s]$. Observe that $\pi' \in D(Y', \alpha')$ and $\pi \in D(Y, \alpha)$. We claim that G_q is a multi-valued (π, ξ) -bifunction from Y into Z . In other words, we must show that

- (a) G'_q is a multi-valued (π', ξ') -function,
- (b) G_q is a multi-valued (π, ξ) -function,
- (c) the compositions $z \circ G'_q$ and $G_q \circ y$ are (π', ξ) -close.

Add (a). Let P' be a member of π' . Then there is a $t \geq 0$ such that the product $P' \times [0, t)$ lies in a member A' of α' . Since K' is an (α', ξ') -function, there is a $T' \in \xi'$ such that $K'(A') \subset T'$. It follows that $G'_q(P') = K'_0(P') = K'(P' \times \{0\}) \subset K'(A') \subset T'$ which proves that G'_q is a (π', ξ') -function.

Add (b). The proof of (b) is analogous to the proof for (a) given above.

Add (c). Since the compositions $K \circ (y \times 1_I)$ and $z \circ (K')$ are (α', ξ) -close so that their restrictions $K \circ (y \times 1_I)|_{Y' \times \{0\}}$ and $z \circ (K')|_{Y' \times \{0\}}$ on $Y' \times \{0\}$ will be $(\alpha'|_{Y' \times \{0\}}, \xi)$ -close. But, if we identify the space Y' with $Y' \times \{0\}$ in $Y' \times I$ we see that the normal cover π' refines $\alpha'|_{Y'}$, and that the restrictions above are $y \circ G_q$ and $z \circ G'_q$. Hence, $z \circ G'_q$ and $y \circ G_q$ are (π', ξ) -close.

Once we know that G_q is a multi-valued (π, ξ) -bifunction, we see that it suffices to take $e \geq d$ because then F_d and F_e are joined by

a multi-valued $(\varepsilon, [b])$ -bihomotopy $L : X \times I \rightarrow Y$, for some normal bicover ε of $X \times I$, so that $G_q \circ L$ is a multi-valued (ε, τ) -bihomotopy which realizes the relation (4).

(a) Indeed, by construction, the normal bicover $[b]$ refines π (i.e., $[b]'$ refines π' and $[b]$ refines π) so that $G'_q \circ L'$ is a multi-valued (ε', ξ') -function and $G_q \circ L$ is a multi-valued (ε, ξ) -function. On the other hand, since the compositions $y \circ (L')$ and $L \circ (x \times 1_I)$ are $(\varepsilon', [b])$ -close, the function G_q is a multi-valued (π, ξ) -bifunction, and the normal cover $[b]$ refines π , it follows that $G_q \circ y \circ (L')$ and $G_q \circ L \circ (x \times 1_I)$ are (ε', ξ) -close. But, the compositions $G_q \circ y$ and $z \circ G'_q$ are (π', ξ) -close, L' is a multi-valued $(\varepsilon', [b]')$ -bifunction, and the normal cover $[b]'$ refines π' so that $G'_q \circ y \circ (L')$ and $z \circ G'_q \circ (L')$ are (ε', ξ) -close. Since the cover $st(\xi)$ refines the cover τ , we conclude that the compositions $z \circ G'_q \circ (L')$ and $G'_q \circ L' \circ (x \times 1_I)$ are (ε', τ) -close.

Add (5). Since $(u, p, q) \in \mathcal{C}_\psi$, it follows that G_p and G_q are joined by a multi-valued (α, ξ) -bihomotopy $K : Y \times I \rightarrow Z$, where α denotes the normal bicover $\psi(u, p, q)$ of $Y \times I$. Choose normal covers β' of Y' and β of Y and functions $r' : \beta' \rightarrow \{2, 3, 4, \dots\}$ and $r : \beta \rightarrow \{2, 3, 4, \dots\}$ such that sets $B' \times [\frac{i-1}{r'B'}, \frac{i+1}{r'B'}]$ and $B \times [\frac{i-1}{rB}, \frac{i+1}{rB}]$, where $B' \in \beta'$, $B \in \beta$, $i = 1, \dots, r'B' - 1$, and $j = 1, \dots, rB - 1$ are contained in a member $A'_{B',i}$ of α' and in a member $A_{B,j}$ of α , respectively. We can assume that β' refines $y^{-1}(\beta)$. It follows that a pair (β', β) is a normal bicover of the bispaces Y . Let $k = \{\beta\}$ and $e = \varphi(k)$. Since $(k, e, e) \in \mathcal{C}_\varphi$, the function F_e is a multi-valued (π, β) -bifunction from X into Y for some normal bicover π of X . It follows that for every $P' \in \pi'$, every $P \in \pi$, and every $B' \in \beta'$ there is a $B'_{P'}$ from β' and a $[B']$, a $B_{P'}$, and a B_P in β such that $B' \subset y^{-1}([B'])$, $F'_e(P') \subset B'_{P'}$, $F_e(P) \subset B_P$, and $B_{P'}$ contains both $y \circ F'_e(P')$ and $F_e \circ x(P')$.

For every $P' \in \pi'$ and every $P \in \pi$, let $s'P' = r'B'_{P'} \tau B_{P'} \tau [B'_{P'}]$,

$$V_{P'} = \left\{ \left[0, \frac{2}{s'P'} \right), \left(\frac{s'P' - 2}{s'P'}, 1 \right] \right\}$$

$$\cup \left\{ \left(\frac{i}{s'P'}, \frac{i+2}{s'P'} \right) \mid i = 1, \dots, s'P' - 3 \right\},$$

and

$$\nu_P = \left\{ \left[0, \frac{2}{\tau B_P} \right), \left(\frac{\tau B_P - 2}{\tau B_P}, 1 \right] \right\}$$

$$\bigcup \left\{ \left(\frac{i}{\tau B_P}, \frac{i+2}{\tau B_P} \right) \mid i = 1, \dots, \tau B_P - 3 \right\}.$$

Put $\varrho' = \{P' \times N' \mid P' \in \pi', N' \in \nu'_{P'}\}$ and $\varrho = \{P \times N \mid P \in \pi, N \in \nu_P\}$. Observe that ϱ' and ϱ are normal covers of $X' \times I$ and $X \times I$, respectively. Let ω be a normal bicover of $X \times I$ such that the normal cover ω' refines ϱ' and the normal cover ω refines ϱ . We claim that the composition $H = K \circ (F_e \times 1_I)$ is a multi-valued (ω, τ) -bihomotopy which joins $G_p \circ F_e$ and $G_q \circ F_e$. Thus, we must show that

- (a) H' is a multi-valued (ω', τ') -function,
- (b) H is a multi-valued (ω, τ) -function, and
- (c) the compositions $z \circ H'$ and $H \circ (x \times 1_I)$ are (ω', τ') -close.

Add (a). Let S' be a member of ω' . Pick a member $R' = P' \times N'$ of ϱ' which contains S' , where $P' \in \pi'$ and $N' \in \nu'_{P'}$. Then

$$H'(S') \subset H'(R') = K'(F'_e(P') \times N') \subset K(B'_{P'} \times N') \subset K'(A'_{B'_{P'}, j}),$$

where j is such that N' lies in the interior of the segment $\left[\frac{1}{\tau B'_{P'}}, \frac{j+2}{\tau B'_{P'}} \right]$. Such an index can surely be found because the length of N' is less than $\frac{1}{\tau B'_{P'}}$. Since K' is a multi-valued (α', ξ') -homotopy, we obtain that the last set in the chain of inclusions above is a subset of a member of ξ' . Hence, H' is a multi-valued (ω', τ') -function.

Add (b). The proof of (b) is analogous to the proof of (a).

Add (c). Let S' be a member of ω' . Choose sets P', N' , and R' as above. Observe that the sets $y \circ F'_e(P')$ and $F_e \circ x(P')$ both lie in a member $B_{P'}$ of β . Since the length of the segment N' is less than $\frac{1}{\tau B_{P'}}$, the product $B_{P'} \times N'$ lies in a member $A_{B_{P'}, j}$ of α , for some $j \in \{1, \dots, \tau B_{P'} - 1\}$. This implies that some member T_1 of ξ contains sets $A = K(y \circ F'_e(P') \times N')$ and $B = K(F_e \circ x(P') \times N')$ because K is a multi-valued (α, ξ) -function. Notice that $B = K \circ (F_e \times 1_I) \circ (x \times 1_I)(R')$. On the other hand, the inclusions $F'_e(P') \subset B'_{P'}$ and $B'_{P'} \subset y^{-1}([B'_{P'}])$

imply that $y \circ F'_e(P') \times N' \subset [B'_{P'}] \times N'$ and therefore that A is a subset of the set $D = K([B'_{P'}] \times N')$. Moreover, since the length of N' is less than $\frac{1}{r[B'_{P'}]}$, the product $L = [B'_{P'}] \times N'$ lies in a member $A_{[B'_{P'}],k}$ of α for some $k \in \{1, \dots, r[B'_{P'}] - 1\}$. Hence, some member T_2 of ξ contains the set $K(L)$. The inclusion $y(B'_{P'}) \subset [B'_{P'}]$ now gives that $J = K \circ (y \times 1_I)(B'_{P'} \times N')$ is a subset of $K(L)$. Once again, since the length of N' is less than $\frac{1}{r'B'_{P'}}$, the set $B'_{P'} \times N'$ lies in a member $E' = A'_{B'_{P'},m}$ of α' for some $m \in \{1, \dots, r'B'_{P'}\}$. We conclude that J is a subset of $K \circ (y \times 1_I)(E')$. Now we use the assumption that $z \circ (K')$ and $K \circ (y \times 1_I)$ are (α', ξ) -close to find a member T_3 of ξ which contains the sets $K \circ (y \times 1_I)(E')$ and $z \circ (K')(E')$. It remains to observe that the inclusion $F'_e(P') \subset B'_{P'}$ implies

$$\begin{aligned} z \circ (K') \circ (F'_e \times 1_I)(R') &= z \circ (K')(F'_e(P') \times N') \subset \\ & z \circ (K')(B'_{P'} \times N') \subset z \circ (K')(E'). \end{aligned}$$

Finally, it follows that the star of T_2 with respect to ξ , and therefore also some member of τ , contains sets $z \circ (K') \circ (F'_e \times 1_I)(S')$ and $K \circ (F_e \times 1_I) \circ (x \times 1_I)(S')$.

Add (6). The proof of (6) is analogous to the proof of (4). \blacksquare

We now define the composition of bihomotopy classes of multi-binets by the rule $[\{G_s\}] \circ [\{F_c\}] = [\{G_{\psi(s)} \circ F_{\varphi(\psi^*(s))}\}]$.

Claim 3. *The composition of bihomotopy classes of multi-binets is well-defined.*

Proof. Let $\kappa = \{K_c\}$ and $\lambda = \{L_s\}$ be multi-binets bihomotopic to φ and ψ , respectively, and let $\mu = \{M_s\}$, where $M_s = L_{\lambda(s)} \circ K_{\kappa(\lambda^*(s))}$ for every $s \in \text{Inc}^b(Z)$. We must show that multi-binets χ and μ are bihomotopic. In other words, that for every $\sigma \in \text{Cov}^b(Z)$ there is an $s \in \text{Inc}^b(Z)$ such that

$$H_t \overset{\sigma}{\simeq}_b M_t \quad \text{for every } t \geq s. \quad (7)$$

Let $\sigma \in \text{Cov}^b(Z)$. Let $\tau \in \sigma^{*4}$ and $\xi \in \tau^*$. Let $s = \{\xi\} \in \text{Inc}^b(Z)$. In order to prove (7), we shall argue that for every $t \geq s$ we can find indices $e \in \text{Inc}^b(Y)$ and $u \in \text{Inc}^b(Z)$ such that

$$H_t \simeq_b^\tau G_p \circ F_e, \quad (8)$$

$$G_p \circ F_e \simeq_b^\tau G_u \circ F_e, \quad (9)$$

$$G_u \circ F_e \simeq_b^\tau L_u \circ F_e, \quad (10)$$

$$L_u \circ F_e \simeq_b^\tau L_u \circ K_e, \quad (11)$$

$$L_u \circ K_e \simeq_b^\tau L_q \circ K_e, \quad (12)$$

$$L_q \circ K_e \simeq_b^\tau M_t, \quad (13)$$

where we put $p = \psi(t)$, $a = \psi^*(t)$, $b = \varphi(a)$, $q = \lambda(t)$, $c = \lambda^*(t)$, and $d = \kappa(c)$. From the relations (8)-(13) with the help of Lemma 2 we shall get (7).

We shall now describe how big e and u must be chosen for relations (8), (9), (10), and (11) to hold separately. The relations (12) and (13) are analogous to relations (9) and (8), respectively. We leave to the reader the task of making a cumulative choice for e and u which accomplishes our goal. It is important to notice that u is selected first while e is selected only once u is already known.

Add (8). We know from the proof of Claim 2 that G_p is a multi-valued (π, ξ) -bifunction from the bispaces Y into the bispaces Z , where $\pi = [s]$ and $s = \bar{\psi}(t, p, p)$. Since $a \geq s$ by the property (1) of Claim 1, it suffices to take $e \geq b$.

Add (9). If $u \geq p$, then G_p and G_u are joined by a multi-valued (α, ξ) -bihomotopy $K : Y \times I \rightarrow Z$, where $\alpha = \bar{\psi}(u, m, m)$ and $m = \psi(u)$. Choose normal covers β' of Y' and β of Y and functions $r' : \beta' \rightarrow \{2, 3, 4, \dots\}$ and $r : \beta \rightarrow \{2, 3, 4, \dots\}$ such that the sets $B' \times [\frac{i-1}{r'B'}, \frac{i+1}{r'B'}]$ and $B \times [\frac{i-1}{rB}, \frac{i+1}{rB}]$, where $B' \in \beta'$, $B \in \beta$, $i = 1, \dots, r'B' - 1$, and $j = 1, \dots, rB - 1$, are contained in a member $A'_{B',i}$ of α' and in a member $A_{B,j}$ of α , respectively. We can assume that β' refines $y^{-1}(\beta)$ so that the pair (β', β) is a normal bicover of Y . Let $k = \{\beta\}$ and $e \geq \varphi(k)$. Just as in the proof of (5) we can see that $K \circ (F_e \times 1_I)$ is

a multi-valued τ -bihomotopy joining the left and the right side of the relation (9).

Add (10). Since multi-binets ψ and λ are bihomotopic, there is a $u \in Inc^b(Z)$, a normal bicover α of $Y \times I$, and a multi-valued (α, ξ) -bihomotopy $S : Y \times I \rightarrow Z$ joining G_u and L_u . Choose a normal bicover β and an index e as above. Then $S \circ (F_e \times 1_I)$ is a multi-valued τ -bihomotopy joining compositions which appear in (10).

Add (11). Let $u \geq q$. Then L_u is a multi-valued (α, ξ) -bifunction from Y into Z , where $\alpha = [s]$, $s = \bar{\lambda}(u, w, w)$, and $w = \lambda(u)$. Since multi-binets φ and k are bihomotopic, there is an index $e \in Inc^b(Y)$ so that F_e and K_e are joined by a multi-valued α -bihomotopy $T : X \times I \rightarrow Y$. The composition $L_u \circ T$ realizes the relation (11). ■

Theorem 1. *The composition of bihomotopy classes of multi-binets is associative.*

Proof. Let $\varphi = \{F_c\}$, $\psi = \{G_s\}$, and $\chi = \{H_p\}$ be multi-binets from X into Y , from Y into Z , and from Z into W , respectively. Let $\mu = \{M_s\}$, $\nu = \{N_p\}$, $\kappa = \{K_p\}$, and $\lambda = \{L_p\}$, where $M_s = G_{\psi(s)} \circ F_{\varphi(\psi^*(s))}$ for every $s \in Inc^b(Z)$ and $N_p = H_{\chi(p)} \circ G_{\psi(\chi^*(p))}$, $K_p = H_{\chi(p)} \circ M_{\mu(\chi^*(p))}$, and $L_p = N_{\nu(p)} \circ F_{\varphi(\nu^*(p))}$, for every $p \in Inc^b(W)$. We must show that multi-binets κ and λ are bihomotopic, i.e., that for every normal bicover $\pi \in Cov^b(W)$ there is an index $p \in Inc^b(W)$ such that

$$K_q \overset{\pi}{\simeq}_b L_q \quad \text{for every } q \geq p. \quad (14)$$

Let $\pi \in Cov^b(W)$. Let $\varrho \in \pi^{**4}$, $\xi \in \varrho^*$, and $\eta \in \xi^*$. Let $p = \{\eta\} \in Inc^b(W)$. In order to prove (14), we shall show that for every $q \geq p$ we can find indices $e \in Inc^b(Y)$ and $s \in Inc^b(Z)$ such that

$$K_q \overset{\varrho}{\simeq}_b H_m \circ G_n \circ F_e, \quad (15)$$

$$H_m \circ G_n \circ F_e \overset{\xi}{\simeq}_b H_m \circ G_s \circ F_e, \quad (16)$$

$$H_m \circ G_s \circ F_e \overset{\eta}{\simeq}_b H_k \circ G_s \circ F_e, \quad (17)$$

$$H_k \circ G_s \circ F_e \simeq_b^e N_l \circ F_e, \tag{18}$$

and

$$N_l \circ F_e \simeq_b^e L_q, \tag{19}$$

where $m = \chi(q)$, $n = \psi(\mu(\chi^*(q)))$, $k = \chi(\nu(q))$, and $l = \nu(q)$. Repeated use of Lemma 2 will give (14) from the relations (15)-(19).

The method of proof is similar to the proof of Claim 3. We shall only describe for each of the relations (15)-(19) how large the indices u and e must be in order that this multi-valued ρ -bihomotopy holds. An easy exercise of putting together all these selections is once again left to the reader. Since relations (18) and (19) are analagous with relations (16) and (15), respectively, it suffices to consider only relations (15)-(17).

Add (15). Observe that H_m is a multi-valued (θ, η) -bifunction, where $\theta = [a]$, $a = \bar{\chi}(q, m, m)$. Let $i = \chi^*(q)$, $j = \mu(i)$, $h = \psi(j)$, $d = \psi(j, n, n)$, $c = \psi^*(j)$, and $\omega = [d]$. Then G_n is a multi-valued $(\omega, [j])$ -bifunction from Y into Z . Since $j \geq i$ and $i \geq a$ by the property (1) from Claim 1, we obtain that $[j]$ refines θ . Let $b = \varphi(c)$. If $e \geq b$, then F_e and F_b are joined by a multi-valued $[c]$ -bihomotopy P . But, $c \geq d$ so that $[c]$ refines ω . Hence, $H_m \circ G_n \circ P$ is a multi-valued ρ -bihomotopy between K_q and $H_m \circ G_n \circ F_e$.

Add (16). As above, H_m is a multi-valued (θ, η) -bifunction from Z into W . If we take $s \geq n$, then $(j, n, s) \in C_\psi$ so that G_n and G_s are joined by a multi-valued $(\alpha, [j])$ -bihomotopy $Q : Y \times I \rightarrow Z$, where $\alpha = \psi(j, n, s)$. But, since $i \geq a$, we see that $[n]$ refines θ . Choose normal covers β' of Y' and β of Y and functions $r' : \beta' \rightarrow \{2, 3, 4, \dots\}$ and $r : \beta \rightarrow \{2, 3, 4, \dots\}$ such that sets $B' \times [\frac{i-1}{r'B'}, \frac{i+1}{r'B'}]$ and $B \times [\frac{i-1}{rB}, \frac{i+1}{rB}]$, where $B' \in \beta'$, $B \in \beta$, $i = 1, \dots, r'B' - 1$, and $j = 1, \dots, rB - 1$, are contained in a member $A'_{B',i}$ of α' and in a member $A_{B,j}$ of α , respectively. We can assume that β' refines $y^{-1}(\beta)$ so that the pair (β', β) is a normal bicover for the bispaces Y . Let $g = \{\beta\}$ and $e \geq \varphi(g)$. Just as in the proofs of (5) and (9), we can see that $H_m \circ Q \circ (F_e \times 1_I)$ realizes the relation (16).

Add (17). Since $\nu(r) \geq r$ for every $r \in Inc^b(W)$, we get $k \geq m$ so that $(q, m, k) \in C_\chi$ and H_m and H_k are joined by a multi-valued (α, η) -bihomotopy $T : Z \times I \rightarrow W$, where $\alpha = \chi(q, m, k)$. Let $v = \chi^*(k)$ and let $s \geq \psi(v)$. Then G_s is a multi-valued $([t], [v])$ -bifunction from Y into

Z , where $t = \bar{\psi}(s, \psi(s), \psi(s))$. Let $u = \psi^*(t)$ and take $e \geq \varphi(u)$. The composition $T \circ ((G_s \circ F_e) \times 1_I)$ realizes the relation (17). ■

8. THE SHAPE CATEGORY OF MAPS \mathcal{HM}^b

For a topological bispace X , let $i^X = \{I_a\} : X \rightarrow X$ be the identity multi-binet defined by $I_a = 1_X$ for every $a \in \text{Inc}^b(X)$. It is easy to show that for every multi-binet $\varphi : X \rightarrow Y$, the following relations hold:

$$[\varphi]_b \circ [i^X]_b = [\varphi]_b = [i^Y]_b \circ [\varphi]_b.$$

We can summarize the above with the following theorem which is the main result of this paper.

Theorem 2. *The topological bispaces as objects together with the bihomotopy classes of multi-binets as morphisms and the composition of bihomotopy classes form the category \mathcal{HM}^b .*

There are some natural functors into the category \mathcal{HM}^b . First of all, there is a functor $S^b : \mathcal{S}^b \rightarrow \mathcal{HM}^b$ from the category \mathcal{S}^b of bispaces and bimaps which leaves the objects unchanged and associates to a bimap $f : X \rightarrow Y$ between bispaces X and Y the bihomotopy class $[\varphi]_b$ of the multi-binet $\varphi = \{F_c\} : X \rightarrow Y$ generated by the bimap f , where $F_c = f$ for every $c \in \text{Inc}^b(Y)$. Then, there is an analogous functor $S_h^b : \mathcal{H}^b \rightarrow \mathcal{HM}^b$ from the category \mathcal{H}^b of bispaces and bihomotopy classes of bimaps. Finally, there is a functor $M^b : \mathcal{HM} \rightarrow \mathcal{HM}^b$ from the author's shape category [4] of spaces and homotopy classes of multi-nets which associates to a topological space X' a bispace $x : X' \rightarrow X$, where X is a fixed single-element space, and which associates to a homotopy class $[\varphi']$ of a multi-net $\varphi' : X' \rightarrow Y'$ between topological spaces X' and Y' the bihomotopy class $[\varphi]_b$ of the multi-binet $\varphi : X \rightarrow Y$ whose domains part is the multi-net φ' . The functor M^b is a full embedding. Since in [4] the author proved that the category \mathcal{HM} is isomorphic to the shape category \mathcal{Sh} , we get the following corollary.

Corollary 1. *Topological spaces X and Y have the same shape if and only if the maps of X and Y into a single-element space are equivalent in the category \mathcal{HM}^b .*

9. SPECIAL MULTI-BINETES

In this section we shall be looking for conditions under which a given bihomotopy class of multi-binets has a representative of a special kind. As a corollary of these results we identify a class of bispaces on which bishape theory and bihomotopy theory coincide.

A bispace X is *internally movable* provided for every $\sigma \in Cov^b(X)$ there is a $\tau \in Cov^b(X)$ such that every multi-valued τ -bifunction into X is σ -bihomotopic to a bimap.

A bispace X is *internally calm* provided there is a $\sigma \in Cov^b(X)$ such that bimaps into X which are σ -bihomotopic are bihomotopic.

A bispace X is *calm* provided there is a $\sigma \in Cov^b(X)$ such that for every τ in $Cov^b(X)$ there is a ϱ in $Cov^b(X)$ with the property that multi-valued ϱ -bifunctions into X which are σ -bihomotopic are also τ -bihomotopic.

A multi-binet $\varphi = \{F_c\}_{c \in Inc^b(Y)}$ from a bispace X into a bispace Y is *regular* provided each bifunction F_c is a bimap. It is called *simple* when there is a bimap f such that $f = F_c$ for every $c \in Inc^b(Y)$.

Theorem 3. *If a bispace Y is internally movable, then every multi-binet φ from a bispace X into Y is bihomotopic to a regular multi-binet.*

Proof. Since Y is internally movable, for every $c \in Inc^b(Y)$ there is a $\chi(c) \in [c]^+$ such that every multi-valued $\chi(c)$ -bifunction into Y is $[c]$ -bihomotopic to a bimap. Let $\sigma_c \in \chi(c)^*$. Let $\lambda : Inc^b(Y) \rightarrow Inc^b(Y)$ be an increasing function such that $\lambda(c) \geq \varphi(\{\sigma_c\})$ for every $c \in Inc^b(Y)$. Then $F_{\lambda(c)}$ is a multi-valued $\chi(c)$ -bifunction so that we can select a bimap $g_c : X \rightarrow Y$ with $g_c \stackrel{[c]}{\simeq}_b F_{\lambda(c)}$, for every $c \in Inc^b(Y)$.

In order to verify that $\psi = \{g_c\}_{c \in Inc^b(Y)}$ is a multi-binet from X into Y , let a $\sigma \in Cov^b(Y)$ be given. Let $\mu \in \sigma^*$ and put $c = \{\mu\}$. For every $d \geq c$ we have $g_d \stackrel{\mu}{\simeq}_b F_{\lambda(d)} \stackrel{\chi(c)}{\simeq}_b F_{\lambda(c)} \stackrel{\mu}{\simeq}_b g_c$. Hence, $g_d \stackrel{\sigma}{\simeq}_b g_c$ for every $d \geq c$.

It remains to check that multi-binets φ and ψ are bihomotopic. Let a normal bicover $\sigma \in Cov^b(Y)$ be given. Let $\mu \in \sigma^*$. Choose an index $c_0 \in Inc^b(Y)$ such that $F_d \stackrel{\mu}{\simeq}_b F_e$ for all $d, e \geq c_0$. Let $c \geq c_0$, $\{\mu\}$.

For every $d \geq c$, we get $g_d \simeq_b^\mu F_{\lambda(d)}$ by construction, while $F_{\lambda(d)} \simeq_b^\mu F_d$ because $\lambda(d) \geq d \geq c_0$. Hence, $g_d \simeq_b^\sigma F_d$ for every $d \geq c$. ■

Theorem 4. *If a bispace Y is both internally movable and calm, then every multi-binnet φ into Y is bihomotopic to a simple multi-binnet.*

Proof. Since Y is calm, there is a normal bicover $\gamma \in \text{Cov}^b(Y)$ such that for every $\sigma \in \text{Cov}^b(Y)$ there is a $\tau \in \text{Cov}^b(Y)$ with the property that γ -bihomotopic multi-valued τ -bifunctions into Y are in fact σ -bihomotopic.

Let $\delta \in \gamma^*$. Since the bispace Y is also internally movable, there is an $\eta \in \delta^+$ such that multi-valued η -bifunctions into Y are δ -bihomotopic to bimaps. Let $c = \varphi(\{\eta\})$. Then F_c is a multi-valued η -bifunction so that it is δ -bihomotopic to a bimap g . Let ψ denote the simple multi-binnet determined by the bimap g .

In order to check that φ and ψ are bihomotopic, let a normal bicover σ of Y be given. Choose a $\tau \in \text{Cov}^b(Y)$ as above. Since φ is a multi-binnet, there is an index $d \geq c$ such that F_e is a multi-valued τ -bifunction for every $e \geq d$. Thus, for every $e \geq d$ we get $F_e \simeq_b^\eta F_c \simeq_b^\delta g$. Hence, $F_e \simeq_b^\gamma g$ so that $F_e \simeq_b^\sigma g$. ■

Theorem 5. *Let Y be a internally calm bispace and let $\varphi = \{f\}$ and $\psi = \{g\}$ be simple multi-binnets into Y . If φ and ψ are bihomotopic, then the bimaps f and g are bihomotopic.*

Proof. Since the bispace Y is internally calm, there is a $\sigma \in \text{Cov}^b(Y)$ such that σ -bihomotopic bimaps into Y are in fact bihomotopic. But, the assumption that the multi-binnets φ and ψ are bihomotopic gives $f \simeq_b^\sigma g$. Hence, the bimaps f and g are bihomotopic. ■

The last three theorems imply that the functor S_h^b is an isomorphism of categories when we restrict to bispaces that have the above properties.

Let \mathcal{A} denote the collection of all bispaces that are at the same time internally movable, internally calm, and calm.

Let $\mathcal{H}_{\mathcal{A}}^b$ be the full subcategory of \mathcal{H}^b with objects precisely the members of the collection \mathcal{A} . The category $\mathcal{HM}_{\mathcal{A}}^b$ is defined similarly.

Let $T_h^b : \mathcal{H}_{\mathcal{A}}^b \rightarrow \mathcal{HM}_{\mathcal{A}}^b$ be the restriction of the functor S_h^b to the category $\mathcal{H}_{\mathcal{A}}^b$.

Theorem 6. *The functor T_h^b is an isomorphism of categories.*

Proof. We shall construct a functor $K_h^b : \mathcal{HM}_{\mathcal{A}}^b \rightarrow \mathcal{H}_{\mathcal{A}}^b$ which satisfies the relations $T_h^b \circ K_h^b = Id$ and $K_h^b \circ T_h^b = Id$. The functor K_h^b leaves the objects unchanged and on morphisms it is defined as follows. Let C be a bihomotopy class of multi-binets between two members X and Y of \mathcal{A} . Let φ be a representative of C and let $g : X \rightarrow Y$ be a bimap such that the simple multi-binet ψ determined by g is bihomotopic to φ . The functor K_h^b associates to C the bihomotopy class of the bimap g . It follows from the above results that this definition is correct and that K_h^b has the required properties. ■

Corollary 2. *On bispaces which are at the same time internally movable, calm, and internally calm, the bihomotopy theory and the bishape theory coincide.*

10. CHARACTERIZATION OF BISHAPE EQUIVALENCES

In this section we shall use classes of internally movable and internally calm bispaces to obtain an analogue for bimaps of G . Kozłowski's characterization of shape equivalences in terms of induced functions of homotopy classes.

For bispaces X and Y , let $[X, Y]_b$ denote the set of all bihomotopy classes of bimaps from X into Y . Every bimap $f : X \rightarrow Y$ induces for every bispace Z a set function $f_Z^\# : [Y, Z]_b \rightarrow [X, Z]_b$ defined by the rule $f_Z^\#([a]_b) = [a \circ f]_b$, for every bimap $a : Y \rightarrow Z$.

Recall that a morphism φ of the bishape category \mathcal{HM}^b is *bishape left invertible* provided there is another bishape morphism ψ with the composition $\psi \circ \varphi$ equal to the identity morphism. The *bishape right invertible morphisms* (or shortly *bishape dominations*) are defined similarly. A bishape morphism is a bishape equivalence iff it is both bishape left invertible and bishape right invertible. When applied to bimaps this notions mean that the bishape morphism determined by this bimap has them.

The following four theorems are similar to the characterization of shape equivalences from [7]. We get information on conditions which imply that functions $f_Z^\#$ induced by a bimap $f : X \rightarrow Y$ and a bispace Z are surjections and injections and we get an information on what kind of bispaces Z one can use.

Theorem 7. *If a bimap $f : X \rightarrow Y$ between topological bispaces is bishape left invertible, then for every bispace Z which is at the same time internally movable and internally calm the induced function $f_Z^\#$ is a surjection.*

Proof. Let $h : X \rightarrow Z$ be a bimap. The assumption about the bispace Z imply the existence of normal bicovers $\sigma \in Cov^b(Z)$, $\tau \in \sigma^*$, and $\mu \in \tau^+$ such that σ -bihomotopic bimaps into Z are bihomotopic and every multi-valued μ -bifunction into Z is τ -bihomotopic to a bimap. Let $\varrho = h^{-1}(\mu)$. Observe that ϱ is a normal bicover of X . Since f is bishape left invertible, there is a multi-binet $\psi : Y \rightarrow X$ with $\psi \circ f$ bihomotopic to the identity multi-binet on X . In particular, there is an index $a \geq \{\varrho\}$ in $Inc^b(X)$ such that $G_p \circ f \simeq_b^e 1_X$, where $p = \psi(a)$. Let $d = \psi^*(a)$.

The bifunction G_p is a multi-valued $([d], \varrho)$ -bifunction so that there is a bimap $k : Y \rightarrow Z$ which is τ -bihomotopic to the composition $h \circ G_p$. It follows that $k \circ f$ and h are σ -bihomotopic and therefore bihomotopic. In other words, $f_Z^\#([k]_b) = [h]_b$. ■

Theorem 8. *If a bimap $f : X \rightarrow Y$ between topological bispaces is bishape right invertible, then for every internally calm bispace Z the induced function $f_Z^\#$ is an injection.*

Proof. Let $h, k : Y \rightarrow Z$ be bimaps and assume that $f_Z^\#([h]_b) = f_Z^\#([k]_b)$, i.e., that $h \circ f$ and $k \circ f$ are joined by a bihomotopy $m : X \times Y \rightarrow Z$. Since Z is internally calm, there is a $\sigma \in Cov^b(Z)$ such that σ -bihomotopic bimaps into Z are bihomotopic. Let $\mu \in \sigma^*$. Let ϱ be a common refinement of $h^{-1}(\mu)$ and $k^{-1}(\mu)$. Let $\pi \in \varrho^*$. Since f is a bishape domination, there is a multi-binet $\psi : Y \rightarrow X$ with the composition $f \circ \psi$ bihomotopic to the identity multi-binet on Y . Hence, there is a multi-valued π -bihomotopic joining 1_Y and $f \circ G_p$, where $\lambda = f^{-1}(\pi)$, $n = \{\lambda\}$, and $p = \psi(n)$.

Let $\alpha \in \text{Cov}^b(X)$ be such that $\alpha' \in D(X', (m')^{-1}(\rho'))$ and $\alpha \in D(X, m^{-1}(\rho))$. Pick an index $q \geq p$ such that G_q is a multi-valued α -bifunction. Then $G_q \simeq_b^\lambda G_p$ so that

$$f \circ G_q \simeq_b^\pi f \circ G_p \simeq_b^\pi 1_Y.$$

It follows that $h \circ f \circ G_q \simeq_b^\mu h$ and similarly $k \circ f \circ G_q \simeq_b^\mu k$. On the other hand $m \circ (G_q \times 1_I)$ is a multi-valued μ -bihomotopy joining $h \circ f \circ G_q$ and $k \circ f \circ G_q$. Hence, $h \simeq_b^\sigma k$ and therefore $h \simeq_b k$. ■

In order to formulate a partial converse to theorem 7 and for some other results below we shall need the following notion.

Let \mathcal{P} be a class of bispaces. By a \mathcal{P} -exposition of a bispace X we mean a family $\{X_a\}_{a \in \text{Inc}^b(X)}$ of members of \mathcal{P} , a family $\{p_b^a \mid a, b \in \text{Inc}^b(X), a \leq b\}$ of bimaps $p_b^a : X_b \rightarrow X_a$, a family $\{p^a \mid a \in \text{Inc}^b(X)\}$ of bimaps $p^a : X \rightarrow X_a$, and a family $\{J_a \mid a \in \text{Inc}^b(X)\}$ of multi-valued $[a]$ -bifunctions such that

- (1) The relation $a \leq b$ in $\text{Inc}^b(X)$ implies $p^a \simeq_b p_b^a \circ p^b$,
- (2) For every $a \in \text{Inc}^b(X)$, we have $J_a \circ p^a \simeq_b^{[a]} 1_X$, and
- (3) The relation $a \leq b$ in $\text{Inc}^b(X)$ implies $J_b \simeq_b^{[a]} J_a \circ p_b^a$.

Theorem 9. *Let \mathcal{P} be a class a bispaces. If a bispace X has a \mathcal{P} -exposition and a bimap $f : X \rightarrow Y$ is such that the induced function $f_Z^\#$ is a bijection for every $Z \in \mathcal{P}$, then f is bishape left invertible.*

Proof. By assumption the bispace X has a \mathcal{P} -exposition formed by bispaces X_a from \mathcal{P} , bimaps p_b^a and p^a , and multi-valued $[a]$ -bifunctions J_a . We shall define a multi-binet $\psi : Y \rightarrow X$ such that $\psi \circ f$ is bihomotopic to the identity multi-binet on X .

Let $a \in \text{Inc}^b(X)$. Since the function $f_{X_a}^\#$ is a surjection, there is a map $r^a : Y \rightarrow X_a$ with $p^a \simeq_b r^a \circ f$. Put $G_a = J_a \circ r^a$ and $\psi = \{G_a\}_{a \in \text{Inc}^b(X)}$.

In order to verify that ψ is a multi-binet, let a normal bicover $\sigma \in \text{Cov}^b(X)$ be given. Let $\tau \in \sigma^*$ and put $a = \{\tau\}$. For every $b \geq a$, we have

$$r^a \circ f \simeq_b p^a \simeq_b p_b^a \circ p^b \simeq_b p_b^a \circ r^b \circ f.$$

Since $f_{X_a}^\#$ is also an injection, we obtain $r^a \simeq_b p_b^a \circ r^b$. It follows that

$$G_a = J_a \circ r^a \simeq_b^{\tau} J_a \circ p_b^a \circ r^b \simeq_b^{\tau} J_b \circ r^b = G_b$$

and therefore $G_a \simeq_b^{\sigma} G_b$.

We shall now check that $\psi \circ f$ is bihomotopic to i^X , i.e., that for every $\sigma \in \text{Cov}^b(X)$ there is an $a \in \text{Inc}^b(X)$ such that $G_{\psi(b)} \circ f \simeq_b^{\sigma} 1_X$ for every $b \geq a$.

Let $\sigma \in \text{Cov}^b(X)$ and $\tau \in \sigma^*$. Put $a = \{\tau\}$. For every $b \geq a$, we get $\psi(b) \geq \psi(a) \geq a$ so that $G_{\psi(b)} \simeq_b^{\tau} G_{\psi(a)}$. Hence,

$$G_{\psi(b)} \circ f \simeq_b^{\tau} G_{\psi(a)} \circ f = J_{\psi(a)} \circ r^{\psi(a)} \circ f \simeq_b^{\tau} J_{\psi(a)} \circ p^{\psi(a)} \simeq_b^{\tau} 1_X.$$

Thus, $G_{\psi(b)} \circ f \simeq_b^{\sigma} 1_X$. ■

Theorem 10. *Let \mathcal{P} and \mathcal{R} be classes of bispaces such that each member of \mathcal{R} is both internally movable and internally calm. If a bispace X has a \mathcal{P} -exposition, a bispace Y has an \mathcal{R} -exposition, and a bimap $f : X \rightarrow Y$ is such that the function $f_Z^\#$ is a bijection for every $Z \in \mathcal{P}$ and an injection for every $Z \in \mathcal{R}$, then f is a bishape equivalence.*

Proof. First we shall show that for every $\sigma \in \text{Cov}^b(Y)$ there is a $\tau \in \text{Cov}^b(Y)$ such that the relation $U \circ f \simeq_b^{\tau} V \circ f$ for multi-valued τ -bifunctions $U, V : Y \rightarrow Y$ implies the relation $U \simeq_b^{\sigma} V$.

Let a $\sigma \in \text{Cov}^b(Y)$ be given. Let $\mu \in \sigma^*$ and $\xi \in \mu^*$. Put $c = \{\xi\}$. By assumption the bispace Y has an \mathcal{R} -exposition formed by bispaces Y_d from \mathcal{R} , bimaps q_c^d and q^d , and multi-valued $[d]$ -bifunctions K_d .

Since the bispace Y_c is internally calm, there is a normal bicover α of Y_c such that α -bihomotopic bimaps into Y_c are bihomotopic. Let $\beta \in \alpha^*$ be such that $K_c : Y_c \rightarrow Y$ is a multi-valued (β, ξ) -bifunction.

Since Y_c is also internally movable, there is a normal bicover $\gamma \in \beta^+$ of Y_c with the property that every multi-valued γ -bifunction into Y_c is β -bihomotopic to a bimap. Let $\tau = (q^c)^{-1}(\gamma)$.

Let $U, V : Y \rightarrow Y$ be multi-valued τ -bifunctions and assume that $U \circ f \simeq_b^\tau V \circ f$. Then $q^c \circ U$ and $q^c \circ V$ are multi-valued γ -bifunctions so that there are bimaps a and b from Y into Y_c with $a \simeq_b^\beta q^c \circ U$ and $b \simeq_b^\beta q^c \circ V$. Our choices imply the following chain of relations:

$$a \circ f \simeq_b^\beta q^c \circ U \circ f \simeq_b^\tau q^c \circ V \circ f \simeq_b^\beta b \circ f.$$

It follows that $a \circ f \simeq_b b \circ f$ and therefore $a \simeq_b b$ because $f_{Y_c}^\#$ is an injection. Now, we have

$$U \simeq_b^\mu K_c \circ q^c \circ U \simeq_b^\mu K_c \circ a \simeq_b^\mu K_c \circ b \simeq_b^\mu K_c \circ q^c \circ V \simeq_b^\mu V.$$

Hence, $U \simeq_b^\sigma V$.

Now we are ready to prove the theorem. Let ψ be a multi-binet constructed in the proof of Theorem 9. We shall show that $f \circ \psi$ is bihomotopic to i^Y , i.e., that for every $\sigma \in Cov^b(Y)$ there is an index $a \in Inc^b(Y)$ such that $f \circ G_{\psi(\varphi^*(b))} \simeq_b^\sigma 1_Y$ for every $b \geq a$. Since we can take $\varphi^*(b) = \{f^{-1}([b])\}$, we must prove that $f \circ G_{\lambda(b)} \simeq_b^\sigma 1_Y$ for every $b \geq a$, where $\lambda(b) = \psi(\{f^{-1}([b])\})$.

Let a $\sigma \in Cov^b(Y)$ be given. Choose a normal bicover τ of Y with the property described above. Let $\varrho \in \tau^*$ and put $a = \{\varrho\}$. For every $b \geq a$, $f \circ G_{\lambda(b)}$ and 1_Y are multi-valued τ -bifunctions of Y into itself and

$$f \circ G_{\lambda(b)} \circ f = f \circ J_{\lambda(b)} \circ \tau^{\lambda(b)} \circ f \simeq_b^\varrho f \circ J_{\lambda(b)} \circ p^{\lambda(b)} \simeq_b^\varrho f = 1_Y \circ f.$$

It follows that $f \circ G_{\lambda(b)} \simeq_b^\sigma 1_Y$. ■

11. APPROXIMATE CONTRACTIBILITY OF MAPS AND TRIVIAL BISHAPE

As an illustration of the new insight offered in shape theory of maps by approach via multi-binets we will characterize bispaces of trivial bishape as approximately contractible bispaces.

Let us agree that a bispace T is *trivial* provided both the codomain and the domain of T are single-element spaces. A bispace X has *trivial bishape* provided it is equivalent in the bishape category \mathcal{HM}^b to a trivial bispace.

As we shall see shortly, bispaces of trivial bishape coincide with approximately contractible bispaces which we define next. A bispace X is called *approximately contractible* provided for every normal bicover σ of X there is a normal bicover τ of X such that every two multi-valued τ -bifunctions into X are σ -bihomotopic.

Theorem 11. *A bispace X is approximately contractible if and only if X has trivial bishape.*

Proof. (\implies). Let T be a trivial bispace and let $c : X \rightarrow T$ be the obvious bimap. Let $a \in X$ and $a' \in x^{-1}(a)$. Define a bimap $g : T \rightarrow X$ by $g(T) = a$ and $g'(T') = a'$. Let $\gamma : X \rightarrow T$ and $\psi : T \rightarrow X$ be simple multi-binets determined by bimaps c and g , respectively. Then $\gamma \circ \psi = i^T$. In order to see that X and T are bishape equivalent, it remains to check that $\psi \circ \gamma \simeq_b i^X$, i.e., that $g \circ c \simeq_b^\sigma 1_X$ for every normal bicover σ of X .

Let a $\sigma \in \text{Cov}^b(X)$ be given. Since X is approximately contractible, there is a $\tau \in \text{Cov}^b(X)$ such that multi-valued τ -bifunctions into X are σ -bihomotopic. But, $g \circ c$ and 1_X are both multi-valued τ -bifunctions of X into itself. Hence, $g \circ c \simeq_b^\sigma 1_X$.

(\impliedby). It suffices to observe that every trivial bispace T is approximately contractible and that approximate contractibility is preserved under bishape dominations and bishape equivalence. ■

Let us call a bimap $f : X \rightarrow Y$ *constant* provided both $f'(X')$ and $f(X)$ are single points (i.e., maps f' and f are constant). The following

notion is a version of path-connectedness for maps. A bispace X is *path-connected* provided every two constant bimaps into X are bihomotopic.

Theorem 12. *If a bispace X has trivial bishape, then every two bimaps from X into a path-connected bispace which is both internally movable and internally calm are bihomotopic.*

Proof. Let Z be a path-connected bispace which is both internally movable and internally calm. Let $a, b : X \rightarrow Z$ be bimaps. Let T be a trivial bispace. Since X has trivial bishape, the obvious bimap $c : X \rightarrow T$ is a bishape equivalence. It follows from Theorem 7 that there are bimaps $f, g : T \rightarrow Z$ with $a \simeq_b f \circ c$ and $b \simeq_b g \circ c$. But, f and g are constant bimaps into Z so that $f \simeq_b g$ because Z is path-connected. Hence, $a \simeq_b b$. ■

Theorem 13. *Let \mathcal{P} be a class of path-connected bispaces. If a bispace X has a \mathcal{P} -exposition and every two bimaps of X into a member of \mathcal{P} are bihomotopic, then X has trivial bishape.*

Proof. Let $c : X \rightarrow T$ be the obvious bimap from X into a trivial bispace T . We shall show that c is bishape left invertible. Then X will be bishape dominated by the approximately contractible bispace T and will therefore have trivial bishape by Theorem 11. According to Theorem 9, we must in fact check that the induced function $c_Z^\#$ is a bijection for every $Z \in \mathcal{P}$.

In order to see that $c_Z^\#$ is a surjection for every $Z \in \mathcal{P}$, let Z be from \mathcal{P} and let $f : X \rightarrow Z$ be a bimap. Let $g : T \rightarrow Z$ be a constant bimap. Then f and $g \circ c$ are two bimaps of X into Z . By assumption, they are bihomotopic so that $[f]_b = c_Z^\#([g]_b)$ and $c_Z^\#$ is onto.

Finally, in order to see that $c_Z^\#$ is an injection for every $Z \in \mathcal{P}$, let Z be from \mathcal{P} and let $f, g : T \rightarrow Z$ be bimaps and assume that $c_Z^\#([f]_b) = c_Z^\#([g]_b)$, i.e., that $f \circ c \simeq_b g \circ c$. But, the bimaps f and g are constant bimaps into Z . Since Z is path-connected, we obtain $f \simeq_b g$ so that $c_Z^\#$ is indeed one-to-one. ■

References

- [1] Aló, R.A. and Shapiro, M.L., *Normal Topological Spaces*, Cambridge Univ. Press, Cambridge, 1972.
- [2] Baladze, V.H., *Fiber shape theory of maps and resolutions*, Bull. Acad. Sci. of Georgian SSR **141** (1991), 489-492.
- [3] Borsuk, K., *Concerning homotopy properties of compacta*, Fund. Math. **62** (1968), 223-254.
- [4] Čerin, Z., *Shape theory intrinsically*, Publicacions Mathématiques **37** (1993), 317-334.
- [5] Dold, A., *Lectures on Algebraic Topology*, Springer-Verlag, Berlin, 1972.
- [6] Edwards, D.A. and McAuley, P.T., *The shape of a map*, Fund. Math. **96** (1977), 195-210.
- [7] Kozłowski, G., *Images of ANR's*, Trans. Amer. Math. Soc. (to appear).
- [8] Mardešić, S. and Segal, J., *Shape Theory*, North Holland, Amsterdam, 1982.
- [9] Sanjurjo, J., *An intrinsic description of shape*, Trans. AMS **329** (1992), 625-636.
- [10] Spanier, E.H., *Algebraic Topology*, McGraw-Hill, New York, 1966.

Kopernikova 7, 41020 Zagreb,

Croatia, Europe

E-mail address: zcerin@X400.SRCE.HR

Recibido: 30 de Noviembre de 1993