

A remark on the blow-up of the solutions of the equation $u_t + f(x) a(u) u_x = h(x, u)$

João-Paulo DIAS and Mário FIGUEIRA

Abstract

We consider the Cauchy problem for the equation $u_t + f(x)a(u)u_x = h(x, u)$ where f, a and h are real C^1 functions, $f \geq \theta > 0$, $a' > 0$, $h_u \geq 0$ and $h_x \leq 0$. Following the ideas of Lax [4] and Klainerman-Majda [3], we prove a blow-up result for the solutions with special data corresponding, in certain cases, to the development of a singularity in u_x .

1 Introduction and statement of the result

Let us consider the scalar conservation law

$$u_t + a(u) u_x = 0, \quad a \in C^1(\mathbf{R}), \quad (x, t) \in \mathbf{R}^2 \quad (1.1)$$

The study of the development of singularities for the solution of the Cauchy problem for the equation (1.1) has been treated by Lax [4] and Majda [6] by proving the appearance of shocks if we impose some conditions to the function a and to the initial data u_0 . In this paper we extend some of these results to the equation

$$u_t + f(x) a(u) u_x = h(x, u), \quad f, a \in C^1(\mathbf{R}), \quad h \in C^1(\mathbf{R}^2) \quad (1.2)$$

by using a method similar to the one employed by Klainerman - Majda [3] for a system of conservation laws.

1991 Mathematics Subject Classification: 35L65, 35L67

Servicio Publicaciones Univ. Complutense. Madrid, 1996.

For the special cases of the equation (1.2) related to the generalised Burgers equation, Natalini - Tesei [8] gave some conditions for the initial data in order to obtain blow-up results for the L^∞ norm of the solution. In the last section we give some applications to a class of equations arising in physics.

We assume

$$\begin{aligned} f(\xi) \geq \theta > 0, \quad a'(\xi) \geq \rho > 0, \quad \forall \xi \in \mathbf{R}, \quad h_u \geq 0, \quad h_x \leq 0 \quad (1.3) \\ \text{and } f \in W^{1,\infty}(\mathbf{R}), \quad h(\cdot, \xi) \in W^{1,\infty}(\mathbf{R}) \quad \text{for each } \xi \in \mathbf{R}. \end{aligned}$$

Following Douglis [1] and Li-Yu [5], ch.1, if we take the initial data $u_0 \in C^1(\mathbf{R}) \cap W^{1,\infty}(\mathbf{R})$ then there exists a unique local solution

$$u \in C^1(\mathbf{R} \times [0, T_0]) \cap C^1([0, T_0]; L^\infty(\mathbf{R})) \cap C([0, T_0]; W^{1,\infty}(\mathbf{R})) \quad (1.4)$$

of the equation (1.2) such that $u(\alpha, 0) = u_0(\alpha), \forall \alpha \in \mathbf{R}$. We will denote by $[0, T']$ the corresponding maximal interval of existence where the sharp continuation principle (cf. [6], 2.3), can be applied.

For such a solution let us consider the equation of the characteristics

$$\frac{dx}{dt}(t) = f(x(t)) - a(u(x(t), t)) \quad \text{with } x(0) = \alpha, \alpha \in \mathbf{R}. \quad (1.5)$$

Along this characteristic curve the solution u satisfies the differential equation

$$\frac{d}{dt} u(x(t), t) = h(x(t), u(x(t), t)) \quad \text{with } u(x(0), 0) = u(\alpha, 0) = u_0(\alpha). \quad (1.6)$$

We can now state our result which extends previous results of Lax [4] and Majda [6] for conservation laws.

Theorem 1 *Under the above assumption (1.3) consider the unique local solution u of (1.2) for the initial data $u_0 \in C^1(\mathbf{R}) \cap W^{1,\infty}(\mathbf{R})$ and assume that $u \in L^\infty(\mathbf{R} \times [0, T'])$ where $[0, T']$ is the corresponding maximal interval of existence. Let $\alpha_0 \in \mathbf{R}$ be such that $u'_0(\alpha_0) < 0$ and let $x(t) = x(t; \alpha_0)$ be the corresponding characteristic curve starting in $x(0) = \alpha_0$. Then, or $\limsup_{t \rightarrow T'} (\|u_x(\cdot, t)\|_{L^\infty} + \|u_t(\cdot, t)\|_{L^\infty}) = +\infty$*

$$\text{or } \liminf_{t \rightarrow T'} \int_0^t a'(u(x(\tau), \tau)) u_x(x(\tau), \tau) d\tau = -\infty$$

and hence $\liminf_{t \rightarrow T'} u_x(x(t), t) = -\infty$.

Moreover $T' \leq T^* = (-\rho f(\alpha_0) u'_0(\alpha_0))^{-1}$.

Remark. Since we suppose $u \in L^\infty(\mathbf{R} \times [0, T'])$, it is enough to assume $a' > 0$ to prove the blow-up result.

The authors are indebted to Prof. L. Sanchez for stimulating discussions and to Prof. Li Ta-tsien who has pointed out a mistake in a first redaction of this note and suggested some other improvements.

This research was partially supported by the European Project HCM N^o CHRX/CT93/0407.

2. Proof of Theorem 1.

Following an idea of Klainerman - Majda [3] we have, along the characteristic curve defined by (1.5),

$$\frac{d}{dt} \left(\frac{\partial x}{\partial \alpha} \right) = \frac{\partial}{\partial \alpha} \left(\frac{dx}{dt} \right) = f' \frac{\partial x}{\partial \alpha} a(u) + f a'(u) \frac{\partial u}{\partial \alpha}$$

Hence, by the theory of linear ordinary differential equations, we have

$$\frac{\partial x}{\partial \alpha}(t) = \left(\exp \int_0^t f' a(u) d\tau \right) \left\{ 1 + \int_0^t f a'(u) \frac{\partial u}{\partial \alpha} \left[\exp \left(- \int_0^s f' a(u) d\tau \right) \right] ds \right\} \tag{2.1}$$

On the other hand, by the results on the derivative of the solution of an ordinary differential equation in order to the initial data (cf. Petrovski [9], for example) we obtain, from (1.5),

$$\frac{\partial x}{\partial \alpha}(t; \alpha) = \exp \int_0^t \frac{\partial}{\partial x} (fa(u)) (x(\tau), \tau) d\tau \tag{2.2}$$

Also, we have, from (1.6), if $u'_0(\alpha) \neq 0$,

$$\frac{d}{dt} \left(\frac{\partial u}{\partial u_0} \right) = h_u \frac{\partial u}{\partial u_0} + h_x \frac{\partial x}{\partial u_0} = h_u \frac{\partial u}{\partial u_0} + h_x \frac{\partial x}{\partial \alpha} (u'_0(\alpha))^{-1} \tag{2.3}$$

Hence, in a neighborhood of $\alpha_0 \in \mathbf{R}$ such that $u'_0(\alpha_0) < 0$ we have (since $h_u \geq 0$, $h_x \leq 0$ and $\frac{\partial x}{\partial \alpha} \geq 0$), by (2.3),

$$\frac{\partial u}{\partial u_0}(t; u_0) \geq \frac{\partial u}{\partial u_0}(0; u_0) = 1$$

and so

$$\frac{\partial u}{\partial \alpha}(t; \alpha) = \frac{\partial u}{\partial u_0}(t; u_0) u'_0(\alpha) \leq u'_0(\alpha) < 0 \quad (2.4)$$

Furthermore, since $f > 0$, we obtain from (1.5), along the characteristic curve

$$f'a(u) = \frac{f'}{f} \frac{dx}{dt} = \frac{d}{dt} \log f$$

and so

$$\exp\left(\int_0^t f'a(u) d\tau\right) = \frac{f(x(t))}{f(\alpha)}. \quad (2.5)$$

Introducing (2.4) and (2.5) in (2.1) we obtain

$$\frac{\partial x}{\partial \alpha}(t; \alpha) = \frac{f(x(t))}{f(\alpha)} \left[1 + f(\alpha) \int_0^t a'(u(x(s), s)) \frac{\partial u}{\partial u_0}(s; u_0) u'_0(\alpha) ds \right] \quad (2.6)$$

Hence, since by (1.3) $a' \geq \rho > 0$, we obtain if $u'_0(\alpha_0) < 0$, by applying (2.4) and (2.6),

$$\frac{\partial x}{\partial \alpha}(t; \alpha_0) \leq \frac{f(x(t))}{f(\alpha_0)} (1 + \rho f(\alpha_0) u'_0(\alpha_0)t) \quad (2.7)$$

and the right hand side is equal to zero for $t = T^* = (-\rho f(\alpha_0) u'_0(\alpha_0))^{-1}$. Introducing (2.5) in (2.2) we derive

$$\frac{\partial x}{\partial \alpha}(t; \alpha) = \frac{f(x(t))}{f(\alpha)} \exp\left(\int_0^t (fa'(u)u_x)(x(\tau), \tau) d\tau\right) \quad (2.8)$$

and so, by (2.7) and (2.8), there exists a $T \leq T^*$ such that, or

$$\limsup_{t \rightarrow T^+} (\|u_x(\cdot, t)\|_{L^\infty} + \|u_t(\cdot, t)\|_{L^\infty}) = +\infty$$

or

$$\liminf_{t \rightarrow T^-} \frac{\partial x}{\partial \alpha}(t; \alpha_0) = \liminf_{t \rightarrow T^-} \frac{f(x(t))}{f(\alpha_0)} \exp\left(\int_0^t (fa'(u)u_x)(x(\tau), \tau) d\tau\right) = 0,$$

that is, since $f(x) \geq \theta > 0$,

$$\liminf_{t \rightarrow T^-} \int_0^t (a'(u)u_x)(x(\tau), \tau) d\tau = -\infty$$

and the theorem is proved.

3. Examples.

Our first example of application of theorem 1 is a semi-linear perturbation of the Burgers equation which can not be reduced by a suitable transformation to the Burgers equation in the framework of [2]:

$$u_t + u u_x = \lambda u^p, \quad \text{for odd } p > 1 \text{ and } \lambda > 0 \quad (3.1)$$

For this equation we obtain the following blow-up result under the assumptions of theorem 1:

$$\liminf_{t \rightarrow T} \int_0^t u_x(x(\tau), \tau) d\tau = -\infty$$

for a certain $T \leq T^* = (-u'_0(\alpha_0))^{-1}$ if $u'_0(\alpha_0) < 0$ and where $x(t)$ is the characteristic curve corresponding to $x(0) = \alpha_0$.

Other kind of results, concerning the blow-up of the L^∞ space norm of the solutions of (3.1) for suitable initial data can be found in [8].

Now, consider the more general equation

$$u_t + a(u) u_x + \lambda h(u) = 0, \quad (3.2)$$

with $\lambda < 0$, $a'(\xi) \geq \rho > 0$, $\forall \xi \in \mathbf{R}$ and $h' \geq 0$. These equations are introduced in [7] (with the supplementary condition $h'(\xi) > 0$ for $\xi > 0$) and appear in the study of the so called Gunn effect in semiconductors. The situation described in theorem 1 corresponds to the appearance of shocks pointed out in section 2.4 of [7] in the case of the existence of a negative dissipation term.

Finally, consider the equation (3.2) in the special case

$$u_t + u^k u_x - u^p = 0, \quad 0 < p < 1, k > 0, \quad (3.3)$$

for positive solutions (see [7] for the positive dissipation case). For this equation the theorem 1 can not be applied without modification. Take a smooth strictly positive initial data u_0 . Along the characteristic curve $x(t; \alpha)$ defined by (1.5) we easily obtain

$$u(x(t; \alpha), t) = \left[(1-p)t + u_0(\alpha)^{1-p} \right]^{\frac{1}{1-p}}, \quad t \geq 0.$$

Following the proof of theorem 1 we derive

$$\frac{\partial x}{\partial \alpha}(t; \alpha) = 1 + k u_0'(\alpha) u_0^{-p}(\alpha) \int_0^t [(1-p)s + u_0(\alpha)^{1-p}]^q ds, \quad (3.4)$$

where $q = \frac{k-1+p}{1-p} > -1$.

Hence, for α_0 such that $u_0'(\alpha_0) < 0$, the right hand side of (3.4) attains zero for a certain $T^* < +\infty$. Therefore we obtain, as in the proof of theorem 1,

$$\liminf_{t \rightarrow T^*} \int_0^t (u^{k-1} u_x)(x(\tau; \alpha_0), \tau) d\tau = -\infty$$

and hence $\liminf_{t \rightarrow T^*} u_x(x(t; \alpha_0), t) = -\infty$.

References

- [1] A. Douglis, *Some existence theorems for hyperbolic systems of partial differential equations in two independent variables*, Comm. Pure Appl. Math., 5, 119-154 (1952).
- [2] K.T. Joseph and P.L. Sachdev, *On the solution of the equation $u_t + u^n u_x + H(x, t, u) = 0$* , Quart. of Appl. Math. 52, 519-527 (1994).
- [3] S. Klainerman and A. Majda, *Formation of singularities for wave equations including the nonlinear vibrating string*, Comm. Pure Appl. Math. 33, 241-263 (1980).
- [4] P.D. Lax, *Development of singularities of solutions of nonlinear hyperbolic partial differential equations*, J. Math. Phys. 5, 611-613 (1964).
- [5] Li Ta-tsien and Yu Wen-ci, *Boundary value problems for quasilinear hyperbolic systems*, Duke University Mathematics Series, Vol.5, 1985.
- [6] A. Majda, *Compressible fluid flow and systems of conservation laws in several space variables*, Applied Math. Sciences, Vol. 53, Springer, 1984.
- [7] J.D. Murray, *On the Gunn effect and other physical examples of perturbed conservation equations*, J. Fluid Mech. 44, 315-346 (1970).

- [8] R.Natalini and A.Tesei, *Blow-up of solutions for a class of balance laws*, Comm. Part. Diff. Eq., 19, 417-453 (1994).
- [9] I.G.Petrovski, *Ordinary differential equations*, Prentice-Hall, 1966.

CMAF/University of Lisbon, Recibido: 23 de Octubre de 1995
2 Av. Prof. Gama Pinto,
1699 Lisboa Codex-PORTUGAL