

## A multiplier theorem for the Hankel transform.

Rafał KAPELKO

### Abstract

Riesz function technique is used to prove a multiplier theorem for the Hankel transform, analogous to the classical Hörmander-Mihlin multiplier theorem [6].

The celebrated Hörmander-Mihlin multiplier theorem [6] says that if a function  $m$  on  $R^n$  satisfies the following condition

$$\sup_{R>0} R^{-n} \sum_{|l|\leq k_0} \int_{R<|x|<2R} |R^{|l|} D^l m(x)|^2 dx < \infty \quad (1)$$

for some integer  $k_0 > \frac{n}{2}$  then the operator  $T_m$  defined by  $\widehat{(T_m g)} = m\widehat{g}$  is bounded on every  $L^p(R^n)$ ,  $1 < p < \infty$ .

Restriction of the theorem to the set of radial functions on  $R^n$  gives the multiplier theorem on spaces  $L^p(R_+, x^{2\alpha+1} dx)$ ,  $1 < p < \infty$  with  $\alpha = \frac{n-2}{2}$ . The ordinary Fourier transform on  $R^n$  has to be replaced by the Hankel transform

$$\widehat{f}(y) = 2^\alpha \Gamma(\alpha + 1) \int_0^\infty f(x) (yx)^{-\alpha} J_\alpha(xy) x^{2\alpha+1} dx, \quad (2)$$

where  $J_\alpha$  is the Bessel function of the first kind of order  $\alpha$ .

The assumption (1) gets even the simpler form

$$\sup_{R>0} \left( \int_R^{2R} |x^k m^{(k)}(x)|^2 \frac{1}{x} dx \right)^{\frac{1}{2}} < \infty,$$

where  $k = 0, 1, 2, \dots, k_0$  and  $k_0 > \alpha + 1$ .

It is quite natural to expect that the multiplier theorem should have an extension to all values  $\alpha \geq \frac{1}{2}$  of the real parameter. However the exact repetition of the Hörmander proof does not lead to effect, mainly because the Hankel transform of the derivative of a function has no representation in terms of the transformation of the function. In order to omit this difficulty there were developed two technics in the literature.

The first one, [2], is indirect, uses a relation between the Jacobi polynomials and the Bessel functions but the result obtained there is weaker then expected. The proof goes under stronger assumption

$$\sup_{R>0} R^{-1} \int_R^{2R} |x^{k_0} m^{(k_0)}(x)|^2 x^{-1} dx < \infty, k_0 = [\alpha] + 2.$$

The second one, [4], develops the original Hörmander's technique but instead of the ordinary derivative of a function it makes use of the powers of a Sturm-Liouville operator. The result is like the Hörmander one, but  $k_0 > \alpha + 1$  must be an even number.

The aim of the note is to prove the multiplier theorem in full generality. We assume that  $k_0$  is the least integer greater than  $\alpha + 1$ . In fact  $k_0$  may be a real number if one uses the Weyl fractional derivatives instead of ordinary derivatives. The main idea is based on the fact that the Hankel transform of Riesz function  $R_u^{k_0}(x^2)$  has especially simple form. Then we follow the arguments of Gosselin and Stempak [4].

For a bounded function  $m$  on  $R_+$  we define the multiplier operator  $T_m$  by  $(T_m g)^\wedge = m \hat{g}$ , where  $\hat{\phantom{g}}$  denotes the Hankel transform (2).

**Theorem 1.** Fix  $\alpha \geq \frac{1}{2}$  and let  $k_0$  denote the least integer greater than  $\alpha + 1$ . Assume that a bounded function  $m$  on  $R_+$  satisfies

$$\sup_{R>0} \left( \int_R^{2R} |x^k m^{(k)}(x)|^2 \frac{1}{x} dx \right)^{\frac{1}{2}} < \infty,$$

where  $k = 0, 1, \dots, k_0$ . Then the operator  $T_m$  is of weak-type (1,1) and, consequently is bounded on every  $L^p(R_+, x^{2\alpha+1} dx)$ ,  $1 < p < \infty$ .

In the proof we use the notion of the generalized convolution

$$f * g(x) = \int_0^\infty f(y) T_\alpha^y g(x) y^{2\alpha+1} dy,$$

where  $T_\alpha^y$  is the generalized translation operator

$$T_\alpha^y g(x) = b(\alpha) \int_0^\pi g((x, y)_\theta) \sin^{2\alpha}(\theta) d\theta,$$

$(x, y)_\theta = (x^2 + y^2 - 2xy \cos \theta)^{\frac{1}{2}}$ ,  $b(\alpha) = \pi^{-\frac{1}{2}} \Gamma(\alpha + 1) \left(\Gamma(\alpha + \frac{1}{2})\right)^{-1}$  and  $f, g$  are suitable functions on the half-line (cf [5]).

As usual we use  $C$  with subscripts or without subscripts for a constant which is not necessarily the same at each occurrence.

**Proof.** The main idea of the proof is based on the fact that the Hankel transform of the function

$$R(x) = \frac{1}{\Gamma(k_0)} (u - x^2)_+^{k_0-1}$$

has a very simple form

$$\widehat{R}(x) = \Gamma(\alpha + 1) 2^{\alpha+k_0-1} \left(\frac{\sqrt{u}}{x}\right)^{\alpha+k_0} J_{\alpha+k_0}(\sqrt{ux}). \tag{3}$$

(cf. [7, §4 Theorem 4.15]).

As usual we cut the function  $m$  into small pieces by using a fixed bump function. Let  $\Psi \in C_0^\infty(\mathbb{R}_+)$  with support in  $(1, 2)$  such that  $\sum_{-\infty}^\infty \Psi(2^{-j}x) = 1$  and  $m_j(x) = m(x)\Psi(2^{-j}x)$ . Define new family of functions  $h(x) = m(x^2)$ ,  $h_j(x) = m_j(x^2)$ . First using (3) and applying the method of [4], we will obtain the theorem for  $h$ . More precisely we will prove

$$\|T_h g\|_p \leq C_{1,p} \|g\|_p. \tag{4}$$

Then we will show how to deduce the thesis for the function  $m$  from the thesis for the function  $h$ .

For  $h_j$  we write the reproducing formula

$$h_j(x) = \frac{1}{\Gamma(k_0)} \int_{2^j}^{2^{j+1}} m_j^{(k_0)}(u) (u - x^2)_+^{k_0-1} du.$$

By (3) we have

$$\widehat{h}_j(x) = \Gamma(\alpha + 1) 2^{\alpha+k_0-1} \int_{2^j}^{2^{j+1}} m_j^{(k_0)}(u) \left(\frac{\sqrt{u}}{x}\right)^{\alpha+k_0} J_{\alpha+k_0}(\sqrt{ux}) du. \tag{5}$$

Then  $T_h = \sum_{j=-\infty}^{\infty} T_{h_j}$  where  $T_{h_j} g = \hat{h}_j * g$  and  $g \in L^1(\mathbb{R}_+, x^{2\alpha+1} dx)$ . In order to prove (4) it is sufficient to establish (cf. [4, p.659] and [1, p.75]) that

$$\sum_{j=-\infty}^{\infty} \int_{|x-y_0| > 2|y-y_0|} |T_{\alpha}^y \hat{h}_j(x) - T_{\alpha}^{y_0} \hat{h}_j(x)| x^{2\alpha+1} dx \leq C, \quad (6)$$

with  $C > 0$  independent of  $y, y_0$ .

An application of Leibniz formula yields

$$\left( \int_{2^j}^{2^{j+1}} |m_j^{(k_0)}(x)|^2 dx \right)^{\frac{1}{2}} \leq C(2^j)^{\frac{1}{2}-k_0}, \quad (7)$$

where  $C$  does not depend on  $j$ , and  $k_0 = \alpha + 1 + \epsilon$  for an  $\epsilon > 0$ . We prove the following estimates:

$$\int_t^{\infty} |\hat{h}_j(x)| x^{2\alpha+1} dx \leq C(\sqrt{2^j} t)^{-\epsilon}, \quad (8)$$

$$\int_0^{\infty} |\hat{h}_j(x)| x^{2\alpha+1} dx \leq C. \quad (9)$$

To prove (8) observe that by definition,  $\hat{h}_j(x)$  coincides with the Hankel transform of the function

$$H_j(y) = \frac{\Gamma(\alpha+1)}{\Gamma(\alpha+k_0+1)} \chi_{[\sqrt{2^j}, \sqrt{2^{j+1}}]}(y) m_j^{(k_0)}(y^2),$$

with respect to the measure  $d_1\mu(x) = x^{4\alpha+3+2\epsilon}$ .

Now Schwartz' inequality, the Plancherel formula applied to  $H_j$  and (7) give

$$\begin{aligned} \int_t^{\infty} |\hat{h}_j(x)| x^{2\alpha+1} dx &\leq \left( \int_0^{\infty} |\hat{h}_j(x)|^2 (x^{2\alpha+1+\frac{1}{2}+\epsilon})^2 dx \right)^{\frac{1}{2}} \left( \int_t^{\infty} \frac{1}{x^{1+2\epsilon}} dx \right)^{\frac{1}{2}} \\ &= \left( \int_0^{\infty} |\hat{h}_j(x)|^2 x^{4\alpha+3+2\epsilon} dx \right)^{\frac{1}{2}} t^{-\epsilon} \frac{1}{\sqrt{2\epsilon}} \\ &= C_{\alpha, k_0} \left( \int_{2^j}^{2^{j+1}} |m_j^{(k_0)}(p)|^2 p^{2\alpha+1+\epsilon} dp \right)^{\frac{1}{2}} t^{-\epsilon} \frac{1}{\sqrt{2\epsilon}} \\ &\leq C(2^j)^{\alpha+\frac{1}{2}+\frac{\epsilon}{2}} (2^j)^{\frac{1}{2}-k_0} t^{-\epsilon} = C(\sqrt{2^j} t)^{-\epsilon}. \end{aligned}$$

To prove (9) we use (8). Now changing the variable  $y = x\sqrt{u}$  in (5) we get

$$\int_0^{2^{-\frac{j}{2}}} |\widehat{h}_j(x)|x^{2\alpha+1}dx \leq C_3 \int_{2^j}^{2^{j+1}} |m_j^{(k_0)}(u)|u^{k_0-1}du \int_0^{\sqrt{2}} |J_{\alpha+k_0}(y)|\frac{y^{2\alpha+1}}{y^{\alpha+k_0}}dy.$$

But Schwarz' inequality and (7) yield

$$\int_{2^j}^{2^{j+1}} |m_j^{(k_0)}(u)|u^{k_0-1}du \leq C_1 \left( \int_{2^j}^{2^{j+1}} |m_j^{(k_0)}(u)|^2du \right)^{\frac{1}{2}} (2^j)^{k_0-\frac{1}{2}} \leq C_4.$$

Since  $J_{\alpha+k_0}(x)$  is  $x^{\alpha+k_0}$  asymptotically at  $0^+$  we have

$$\int_0^{2^{-\frac{j}{2}}} |\widehat{h}_j(x)|x^{2\alpha+1}dx \leq C_2.$$

Also, by (8)

$$\int_0^\infty |\widehat{h}_j(x)|x^{2\alpha+1}dx \leq C_2 + \int_{2^{-\frac{j}{2}}}^\infty |\widehat{h}_j(x)|x^{2\alpha+1}dx \leq C.$$

Finally, to get (6) we use inequality (8) with estimates of Gosselin and Stempak (cf. [4, p.661])

$$\begin{aligned} \int_{|x-y_0| \geq 2|y-y_0|} |T_\alpha^y \widehat{h}_j(x) - T_\alpha^{y_0} \widehat{h}_j(x)|x^{2\alpha+1}dx \\ \leq \int_{|y-y_0|}^\infty |\widehat{h}_j(x)|x^{2\alpha+1}dx + \int_{2|y-y_0|}^\infty |\widehat{h}_j(x)|x^{2\alpha+1}dx \\ \leq C_1(1+2^{-\epsilon})(\sqrt{2^j}|y-y_0|)^{-\epsilon}, \end{aligned}$$

which will work for  $\sqrt{2^j}|y-y_0| \geq 1$ .

Since  $h_j$  has support in  $(0, \sqrt{2^{j+1}})$  it follows from [4, Corollary 2.2] and (9) that

$$\begin{aligned} \int_{|x-y_0| \geq 2|y-y_0|} |T_\alpha^y \widehat{h}_j(x) - T_\alpha^{y_0} \widehat{h}_j(x)|x^{2\alpha+1}dx \\ \leq \|T_\alpha^y \widehat{h}_j - T_\alpha^{y_0} \widehat{h}_j\|_{L^1(\mathbb{R}_+, x^{2\alpha+1}dx)} \\ \leq C_1 \sqrt{2^{j+1}}|y-y_0| \|\widehat{h}_j\|_{L^1(\mathbb{R}_+, x^{2\alpha+1}dx)} \\ \leq \sqrt{2}C C_1 \sqrt{2^j}|y-y_0|, \end{aligned}$$

which will be enough whenever  $\sqrt{2^j}|y-y_0| < 1$ .

This completes the proof of (6) and, consequently for the function  $h$ . The result for the function  $m$  follows than from the lemma below.

**Lemma 1.** For  $\alpha > 0$  the transformation  $x \rightarrow x^\alpha$  of  $[0, \infty)$  induces the isomorphism  $m(x) \rightarrow m(x^\alpha)$  of the space of all functions for which

$$\|m\|_{2,k_0} = \sup_{R>0} \left( \int_R^{2R} |x^k m^{(k)}(x)|^2 \frac{1}{x} dx \right)^{\frac{1}{2}} < \infty$$

for  $k = 0, 1, 2, \dots, k_0$ .

**Proof.** This is a simple consequence of fact that space  $\|m\|_{2,k_0}$  is invariant under multiplication by  $x^\alpha$  and Leibniz formula.

**Remark.** The method of Riesz function works when we use the Weyl fractional derivatives instead of ordinary derivatives.

A function  $f$  on  $R_+$  has the Weyl fractional derivative of order  $v > 0$  if there exists a measurable function  $g$  on  $R_+$  such that

$$f(x) = \frac{1}{\Gamma(v)} \int_x^\infty (t-x)^{v-1} g(t) dt$$

for almost all  $x > 0$ . The function  $g$  is unique up to a set of measure zero. It is denoted  $f^{(v)}$  and called  $v$ -fractional derivative of order  $v$ .

The problem is that for a positive integer  $v$  there exist smooth functions in the ordinary sense but not in the Weyl sense.

**Theorem 2.** Let  $m$  be a bounded function on  $R_+$  satisfies the condition

$$\sup_{R>0} \left( \int_R^{2R} |x^v m^{(v)}(x)|^2 \frac{1}{x} dx \right)^{\frac{1}{2}} < \infty,$$

where  $v > \alpha + 1$ ,  $m^{(v)}$  is the Weyl fractional derivative. Then the operator  $T_m$  is of weak-type  $(1,1)$  and, consequently is bounded on every  $L^p(R_+, x^{2\alpha+1} dx)$ ,  $1 < p < \infty$ .

**Proof.** As in the proof of Theorem 1 we define  $h(x) = m(x^2)$  and obtain the theorem for function  $h$ . To do this we don't work with bump functions and define

$$h_j(x) = \frac{1}{\Gamma(v)} \int_{2^j}^{2^{j+1}} m^{(v)}(u) (u-x^2)_+^{v-1} du.$$

Clearly  $T_h = \sum_{-\infty}^{\infty} T_{h_j}$  where  $T_{h_j}g = \hat{h}_j * g$ . The rest is the exact repetition of the proof of Theorem 1. Finally the result for the function  $m$  follows from lemma below.

**Lemma 2.** For  $\alpha > 0$  the transformation  $x \rightarrow x^\alpha$  of  $[0, \infty)$  induces the isomorphism  $m(x) \rightarrow m(x^\alpha)$  of the space of all function for which

$$\|m\|_{2,v} = \sup_{R>0} \left( \int_{\frac{R}{2}}^R |x^v m^{(v)}(x)|^2 \frac{1}{x} dx \right)^{\frac{1}{2}} < \infty.$$

**Proof.** The lemma is a modification of [3, Proposition 3.9]. The only difference is the norm  $\|\cdot\|_{(\mu),2,1}$  is changed into the norm  $\|\cdot\|_{2,v}$  and the proof is essentially the same.

#### Acknowledgements.

I would like to thank Tadeusz Pytlik who suggested me the method I am used and Krzysztof Stempak for useful suggestions, and fruitful discussions.

#### References

- [1] R. Coifman and G. Weiss, *Analyse harmonique non commutative sur certains espaces homogènes*, Springer-Verlag, vol. 242, Berlin and New York, 1971.
- [2] W. C. Connett and A. L. Schwartz, Weak type multipliers for Hankel transforms, *Pac. Jour. of Math.*, vol.63, no. 1, 1976, 125-129.
- [3] J. Galé and T. Pytlik, Functional calculus for infinitesimal generators of holomorphic semigroups, *Jour. of Fun. Anal.*, vol.150, number 2, 1997, 307-355
- [4] J. Gosselin and K. Stempak, A weak-type estimate for Fourier-Bessel multipliers, *Proc. of the Amer. Math. Soc.*, vol.106, no. 3, 1989, 655-662.
- [5] I. Hirschman, Variation diminishing Hankel transforms, *Jour. d'Anal. Math.*, vol.8, 1960/61, 307-336.
- [6] L. Hörmander, Estimates for translation invariant operators in  $L^p$  spaces, *Acta Math.*, no. 104, 1960, 93-140.

- [7] E.M. Stein and G.Weiss, Introduction to Fourier analysis on Euclidean spaces, Princeton Univeristy Press, 1971.

Instytut Matematyczny  
Uniwersytet Wrocławski  
pl. Grunwaldzki 2/4  
50-384 Wrocław  
POLAND  
Rafał Kapelko  
*e-mail:* [kapelko@math.uni.wroc.pl](mailto:kapelko@math.uni.wroc.pl)

Recibido: 8 de Diciembre de 1997

Revisado: 15 de Abril de 1998